

AGROSOL™ FLOWABLE
Systemic Soybean Seed Treatment Fungicide
Contains MERTECT™

ACTIVE INGREDIENTS:	By Wt.
*Captan	29.37%
Related derivatives.	0.63%
**Thiabendazole.	0.53%
INERT INGREDIENTS:	<u>69.47%</u>
TOTAL	100.00%

*N-Thrichloromethylthio-4-cyclohexene-1,2-dicarboximide
 **2-(4-Thiazolyl) benzimidazole

This product contains: 2.77 lb. captan per gallon
 0.06 lb. related derivatives per gallon
 0.05 lb. thiabendazole per gallon

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention. Note to Physician: This product contains a petroleum distillate.

If in eyes, flush with plenty of water. Call a physician.

If on skin, wash with plenty of soap and water.


See side panel for additional precautionary information

EPA Est. No. 37429-GA-1

EPA Reg. No. 2935-471

NET CONTENTS 250 GALLONS

*AGROSOL is a trademark of C-I-L Inc., Canada
 **MERTECT is a registered trademark of Merck & Co., Inc.



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder before reuse. Wash with soap and water after handling and before eating, drinking or smoking.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Exposed treated seeds may be hazardous to birds and other wildlife. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil. Dispose of all excess treated seeds by burial away from bodies of water.

PRODUCT INFORMATION

AGROSOL Flowable is a ready-to-use seed treatment formulation prepared for commercial seed treaters. When used at the correct dosage rate, AGROSOL Flowable will protect seeds from soil and seed borne diseases that cause damping-off, seed decay, seedling blights and pod and stem blight (*Phomopsis* spp.) during germination and early seedling stages.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE OR STIR WELL BEFORE USING. Use at temperatures above 40°F. Use AGROSOL Flowable in liquid and slurry treaters at the rate of 1.5 fl. oz. per bushel of soybean seed (2.5 fl. oz. per 100 lbs. of soybean seed).

NOTE: The purchaser of this product is responsible for insuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any dye added must be cleared for use under 40 CFR, Part 180.1001.

Avoid contamination of feed, food and foodstuffs. Do not graze of cut soybean forage for feed.

Treated seed must not be used for a mixed with food or animal feed, or processed for oil. Seed commercially treated with

3:13

AGROSOL Flowable must be labeled in accordance with all applicable requirements of the Federal and State seed laws.

Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store above 32°F. Product stored at a lower temperature will reconstitute to satisfactory utility if warmed to approximately 70°F and thoroughly shaken or stirred before use. Store treated seed in a ventilated place away from water, food and feed.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use of handling of this product and accepts the product on that condition.

Wilbur-Ellis Company
P. O. Box 16458
Fresno, CA 93755