NET CONTENTS: 32 fl. oz.

AGROSOL™ Pour-on

1/1/21

Systemic Soybean Seed Treatment Fungicida

248-477 P\$105

Contains MERTECT®

EPA Hou to 82 Ches product contains 4.14% molybdenum from sodium molybdate

ACTIVE INGREDIENTS:

%

Thiram (tetramethyl thiuram disulfide) 11.400

2-(4-Thiazolyl)bendimidazole (thiabendazole) 0.314

INERT INGREDIENTS: 88.268

TOTAL 100.000

This product contains 1.09 lb thiram per gallon.

0.03 lb thiabendazole per gallon

This product contains the toxic inert rhodamine B

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional precautionary information

CHIPMAN, A Business Unit of C-I-L Inc.

Stoney Creek, Ontario, Canada

L8G 3Z1

EPA Est 8203-CN-1

EPA Reg No. 8203-41

AGROSOL is a trademark of C-I-L Inc., Canada MERTECT is a registered trademark of Merck & Co., Inc.

NET CONTENTS: 32 fl. oz.

AGROSOL™ Pour-on

Systemic Soybean Seed Treatment Fungicide

Contains MERTECT®

This product contains 4.14% molybdenum from sodium molybdate

ACTIVE INGREDIENTS:

1

Thiram (tetramethyl thiuram disulfide) 11.400

2-(4-Thiazolyl)benzimidazole (thiabendazole) 0.314

INEFT INGREDIENTS: 88.268

TOTAL 100.000

. .is product contains 1.09 lb thiram per gallon.

0.03 lb thiabendazole per gallon

This product contains the toxic inert rhodamine B

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary information

CHIPMAN CHEMICALS, A Business Unit of C-I-L Corporation of America River Rouge, MI 48218

EPA Est 8203-CN-1

EPA Reg No. 8203-41-44742

AGROSOL is a trademark of C-I-L Inc., Canada
MERTECT is a registered trademark of Merck & Co., Inc.

Victoria.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION:

Causes skin and eye irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid inhaling dust. Do not store near food or feed. During commercial exposure in seed treatment and loading operations, wear goggles, clean rubber gloves, protective clothing, and a pesticide respirator jointly approved by the Mine Safety and Health Administration (formerly the U.S. Bureau of Mines) and by the National Institute of Occupational Safety and Health under provisions of 30 CFR, Part II. Wash hands, arms and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work. Wash all clothing with soap and hot water before reusing. Do not use or store near food or feed. Consumption of alcoholic beverages increases the toxic effects of thiram.

STATEMENT OF PRACTICAL TREATMENT:

If swallowed, induce vomiting by drinking 1 or 2 large glasses of water and then touch back of throat with finger. Get medical attention. If in eyes, flush with clean water for at least 15 minutes and then get medical attention. If on skin, wash with soap and water. Get medical attention if skin irritation persists.

ENVIRONMENTAL HAZARDS:

PRODUCT INFORMATION:

AGROSOL Pour-on when used at the correct rate protects seeds from soil and seed borne diseases that cause damping-off, seed decay, seedling blights, and pod and stem blight (Phomopsis spp) during germination and early seedling stages. AGROSOL Pour-on provides essential molybdenum for plant growth where soils are found to be deficient in available molybdenum.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labelling.

SHAKE WELL BEFORE USING.

Apply AGROSOL Pour-on to soybean seed at a rate of 4 fl. oz. per bushel as a planter box application. Mix thoroughly with seed in the planter box as follows:

For Conventional Plate Planters and MAX-EMERGE® Plateless Planters by

John Deere - Pour one bushel of seed into each planter box. Pour enough

AGROSOL Pour-on on top of the seed to treat one bushel. Stir thoroughly

with an axe handle (purchased and used for this purpose only) until all

the seed is uniformly coated. (MAX-EMERGE® is a reg. T.M. of Deere

and Co.)

NOTE: Avoid very deep seeding and exceptionally early sowing under poor growing conditions for maximum benefits. Use only recommended rates.

Lower amounts may not give the desired control. Regularly inspect planter plates of seed drill and clean as required.

Clean-up for Electric Eye Population Monitors: At the end of each day of use, clean out the electric-eye monitors with brushes provided by the planter manufacturers, according to their directions.

BEST AVAILABLE COPY

Pros

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store above 32°F. Product stored at a lower temperature will reconstitute to satisfactory utility if warmed to approximately 70°F and thoroughly shaken or stirred before use. Keep container tightly closed when not in use. Store away from water, food and feed.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse, then puncture and dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT:

Seller's guarantee shall be limited to the terms of the label and subject thereto, buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product or that condition

0588

