



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

JUN 3 2010

Ms. Sandi Escamilla
Registration Specialist
Product Regulatory Department
Wilbur-Ellis Company
P.O. Box 1286
Fresno, CA 93715

Subject: VOLCK® SUPREME ~~OIL~~ SPRAY *WWS 10/13/10*
EPA Registration No. 2935-542
Submission of March 17, 2010

Dear Ms. Escamilla:

The proposed amended label submitted on March 17, 2010, for this product is accepted under the provisions of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, with the further stipulation that the changes listed below must be made to that label.

1. As this product is labeled only for terrestrial uses, "For terrestrial uses:" must be deleted from the "ENVIRONMENTAL HAZARDS:" section on page 2 of the label.
2. Move the third paragraph under "GROUND APPLICATION" in the "SPRAY DRIFT MANAGEMENT" text on page 3 (first column) to a position between the current second ("Do not apply at wind ... site") and third ("Apply as a medium ... nozzles") paragraphs under "SPRAY DRIFT MANAGEMENT". This change is needed because the text related to temperature inversions pertains to aerial as well as ground applications.
3. In the last cell of the "Crop" column in the table for "FOLIAR APPLICATIONS – DILUTE SPRAYS (continued)" (page 5), replace "Hawthorne" with "Hawthorn". Make the same change in the "Crop" column on the "DORMANT – DILUTE SPRAYS" table at the top of page 7.
4. In the "Container Disposal:" portion of the "STORAGE AND DISPOSAL" box (page 7), change the text pertaining to "Refillable container (110 gallons or greater):" as indicated below.
 - a. Delete "Container Disposal:" from the second paragraph (which could be run together with the first paragraph. The subheading "Container Disposal:" already appears in the "STORAGE AND DISPOSAL" box, and repeating it is likely to cause some confusion.

5. The part of the “**Container Disposal:**” text that pertains to “**30 gallon Nonrefillable container:**” is missing a sentence required for rinsing and does not indicate what is to be done with containers that cannot or are not to be recycled. To address these issues, modify the text as indicated below.
- Delete “**Container Disposal:**” from the second paragraph, which could be combined with the first paragraph.
 - Between the sentences “Stand the container ... times” and “Empty the rinsate ... disposal” in the “Triple rinse as follows:” passage, insert the following sentence:

Turn the container over on its other end and tip it back and forth several times.

- After the sentence “To find the nearest site ... rperkins@acrecycle.org”, insert text that indicates what to do if the container is not to be recycled. Following earlier labels for this product, that text would read as follows:
- If container is not to be recycled, offer it for reconditioning or puncture and dispose of it in a sanitary landfill; or by incineration; or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.
6. The part of the “**Container Disposal:**” text that pertains to “**5 gallon Nonrefillable container:**” does not indicate what is to be done with containers that cannot or are not to be recycled. To address these issues, modify the text as indicated below.

- Delete “**Container Disposal:**” from the second paragraph, which could be combined with the first paragraph.
- After the sentence “To find the nearest site ... rperkins@acrecycle.org”, insert text that indicates what to do if the container is not to be recycled. Following earlier labels for this product, that text would read as follows:

If container is not to be recycled, offer it for reconditioning or puncture and dispose of it in a sanitary landfill; or by incineration; or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.


A stamped copy of the label is enclosed for your records.

Submit two copies of final printed labeling before releasing this product for shipment under the labeling accepted via this letter.

If these conditions are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6(e).

If you have any questions regarding this letter, you may reach Dr. William Jacobs by phone at 703-305-6406, by FAX at 703-308-0029, or via e-mail at jacobs.bill@epa.gov.

Sincerely yours,

A handwritten signature in cursive script that reads "Venus Eagle". The signature is written in dark ink and has a long, horizontal flourish extending to the right.

Venus Eagle
Product Manager 01
Insecticide-Rodenticide Branch
Registration Division (7505P)

Volck® Supreme Spray

(INSECTICIDE - MITICIDE)

ACTIVE INGREDIENT	By Wt.
Mineral Oil*	97.95%
OTHER INGREDIENTS	2.05%
	100.00%

*Contains petroleum distillates
Minimum Unsulfonated Residue - 99%
Petroleum Oil Classification - Unclassified

ACCEPTED
With COMMENTS
In EPA Letter Dated:

JUN 3 2010

Under the Federal Insecticide, Fungicide
and Rodenticide Act, As amended, for the
Pesticide Registered under EPA Reg. No:

2935-542

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate and viton. If you want more options, follow the instructions for category G on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators, flaggers, and other handlers must wear:

- long-sleeve shirt and long pants
- shoes plus socks and
- chemical-resistant gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

ENGINEERING CONTROLS

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

FIRST AID	
If on skin or clothing:	-Take off contaminated clothing. -Rinse skin immediately with plenty of water for 15 to 20 minutes. -Call a poison control center or doctor for treatment advice.
If in eyes:	-Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. -Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. -Call a poison control center or doctor for treatment advice.
If swallowed:	-Call a poison control center or doctor for treatment advice. -Do not induce vomiting unless told to do so by a poison control center or doctor. -Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
If Inhaled:	-Move person to fresh air. -If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth- to-mouth, if possible. -Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-892-0099 for emergency medical treatment information. NOTE TO PHYSICIANS: This product contains petroleum distillates and may pose an aspiration pneumonia hazard.	

EPA Reg. No. 2935-542
EPA Est. No. 2935-CA-01^F, 2935-CA-19^R, 2935-CA-31^H
Superscript used is the first letter of the lot number.

NET CONTENTS _____ GALLONS

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as soil or water, is:

- coveralls
- shoes plus socks, and
- chemical-resistant gloves made of any waterproof material.

PRODUCT INFORMATION AND RESTRICTIONS

Volck Supreme Spray is an especially refined Phytonomic Oil Contact Insecticide. It is for use on plants in foliage or the dormant stage.

When using this product in combination with other pesticides as directed on this label, observe appropriate cautions, limitations for the product involved.

For foliage applications, the spray tank must be free of any sulfur residue before using Volck Supreme Spray. Do not combine sulfur sprays with Volck Supreme Spray on foliage and do not follow a previous application of sulfur or apply sulfur following Volck Supreme Spray until sufficient time has elapsed for safety (at least 30 days). If in doubt, spray a portion of one tree previously treated with sulfur, using 1 qt. to 25 gals. water. If sufficient time has not elapsed, burn will show in several days.

Do not use Volck Supreme Spray when temperatures are above 90° Fahrenheit. When temperatures drop below 32° Fahrenheit (usually nighttime lows), do not spray tree and vine crops until daytime temperatures have warmed to a point above 35° Fahrenheit and are free from moisture due to melting frost or heavy dews. Do not use on foliage closer than 60 days after or 90 days before using captan or folpet products.

Application to trees weakened by disease, drought, drying winds, or high nitrogen applications may result in oil injury.

Do not apply Volck Supreme Spray concentrate to Queen Ann variety or other oil sensitive varieties as phytotoxicity to buds and new wood will result.

Observe on all crops and uses - oil effectiveness in dormant and delayed dormant applications is determined by thoroughness of coverage. Concentrate applications may reduce coverage and effectiveness. Oil applications made to trees that are weak or suffering from moisture stress or when slow drying of spray occurs may cause bud, twig or leaf injury.

Phytotoxicity studies have not been made on all fruit varieties, or in all growing areas in California for concentrate applications. Concentrate spraying is a more exacting technique of spray application than dilute. Concentrate spray equipment must be properly calibrated and operated to obtain desired results.

Multiple concentrate applications may cause wood injury. Double applications can occur on the ends of the row during turning - turn off sprayer to avoid multiple applications. Equipment must be capable of giving thorough coverage for maximum insecticidal effectiveness. If San Joaquin Quale is a moderate to severe problem, use dilute application. Concentrate applications may delay bud development. The following conditions may cause oil injury; cool, damp, or extremely dry weather; broken emulsions and/or faulty application. Follow state recommendations and cautions.

Do not apply this product through any type of irrigation system.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof unfit for sale, use or consumption.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator and grower must evaluate all factors and make appropriate adjustments when applying this product.

Do not apply at wind speeds greater than 15 mph at the application site.

Apply as a medium or coarser spray (ASABE standard 572), and the minimum volume mean diameter (VMD) for spinning atomizer nozzles.

GROUND APPLICATION

Apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

Do not apply at wind speeds greater than 15 mph at the application site.

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Apply as a medium or coarser spray (ASABE standard 572), and the minimum volume mean diameter (VMD) for spinning atomizer nozzles.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying outer row. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

AERIAL APPLICATION

For aerial applications of agricultural products, do not release spray at a height greater than 10 feet above the ground, top of crops, or above the orchard canopy.

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Leave at least one swath unsprayed at the downwind edge of the treated field.

Do not apply at wind speeds greater than 15 mph at the application site.


Apply as a medium or coarser spray (ASABE standard 572), and the minimum volume mean diameter (VMD) for spinning atomizer nozzles.

ADDITIONAL PRECAUTIONS FOR AERIAL APPLICATIONS

Apply quantity of product shown for each listed crop for control of listed insects in sufficient water to make a minimum of 20 gals. dilute spray per acre by air application. Follow timing shown for each crop as shown on container label. **Use the aerial application method only when soil conditions do not permit regular ground application.**

MIXING DIRECTIONS

Check this product for proper emulsification by mixing a small quantity of it with water in a small container before filling the entire spray tank. Correct emulsification is indicated when a uniform mixture, with a blue color, results on shaking. If this mixture is not a milky emulsion after shaking, do not use the remaining product. With agitator running, start filling spray tank with water. When combining with wettable powder insecticide or fungicide products, add these first when tank is 1/3 filled. Add Volck Supreme Spray when tank is about 1/2 full and while there is sufficient agitation of the spray to ensure complete emulsification.



FOLIAR APPLICATIONS - DILUTE SPRAYS

Crop	Pests	Volck per 100 gals. Water	Further Use Instructions
Almonds, Apricots, Cherries, Nectarines, Peaches, Pistachio, Plums, Prunes	Brown Apricot Scale, San Jose Scale	1 gal.	For Summer stages of Brown Apricot and San Jose Scale.
	Pacific Spider Mites, Red Spider Mites	3 to 4 qts.	For Summer use on Red Spider Mites.
Apples	Rust Mites	1 to 1½ gals.	For Summer use.
Apples, Pears	Red spider Mites	3 to 4 qts.	For Summer use.
	Italian Pear Scale, San Jose Scale	1 to 1½ gals.	Apply Volck Supreme Spray to Summer forms of San Jose and Italian Pear Scale.
Grapes	Mealybug	See NOTE	For (Post Harvest), Fall spray use only. NOTE: Use 4 to 6 gals. in 200 to 300 gals. of water per acre.
Grapefruit, Lemons	Black Scale, Purple Scale, Red Scale, Red Spider Mites, Yellow Scale	1 to 1½ gals.	For best results, apply Volck Supreme Spray during Winter and Spring months when Red Scale are more easily controlled and trees generally are less reactive to oil sprays. Do not use within 8 weeks of harvest on lemons. Do not apply more than 1500 gallons of spray mix per acre in Florida and Texas. Do not apply more than 2000 gallons of spray mix per acre in California.
Navel Oranges	Black Scale, Purple Scale, Red Scale, Red Spider Mites, Yellow Scale	3 qts.	Do not use on Navel oranges after September 1. Do not apply more than 159 gallons of active ingredient or more than 1500 gallons of spray mix per acre in Texas or Florida. Do not apply more than 212 gallons of active ingredient or more than 2000 gallons of spray mix per acre in California.
Olives	Black Scale, Oleander Scale, Olive Scale (Parlatoria Scale), Spider Mites	1½ gals.	Apply Volck Supreme Spray in the prebloom or in postbloom up to buckshot growth stage.
Hops (Not registered for use in California)	Mites	1 to 2 gals.	Apply Volck Supreme Spray with ground equipment only. Thorough coverage is essential. Use 50 to 200 gals. of dilute spray per acre depending upon plant size. Make applications as needed.
	Powdery Mildew	1 to 2 gals.	Apply Volck Supreme Spray with ground equipment only. Thorough coverage is essential. Use 50 to 200 gals. of dilute spray per acre depending upon plant size. Make first application at early leaf stage. Repeat every 7 to 14 days depending upon new growth. Use shorter interval and/or the higher rate when the conditions for infection are more severe.
	Do not use Propargite (Ornite) with or following an oil spray. Do not spray wet foliage or if temperature is above 90° Fahrenheit or if plants are under heat or moisture stress.		

FOLIAR APPLICATIONS - DILUTE SPRAYS (continued)

Crop	Pests	Volck per 100 gals. Water	Further Use Instructions
Olives (California Only)	Black Scale	1 to 1½ gals.	Apply 400 to 800 gals. dilute spray per acre. Make applications between July 1 and August 31. Higher concentrations of Volck Supreme Spray than noted above may cause phytotoxicity.
Pears	Pear Psylla	1 to 1½ gals.	For Summer use.
	Blister Mite	1 to 1½ gals.	For Fall spray use only.
Valencia Oranges	Black Scale, Purple Scale, Red Scale, Red Spider Mites, Yellow Scale	1 gal.	Do not apply to Valencia oranges after September 1. Do not apply more than 159 gallons of active ingredient or more than 1500 gallons of spray mix per acre in Texas or Florida. Do not apply more than 212 gallons of active ingredient or more than 2000 gallons of spray mix per acre in California.
Walnuts	European Fruit Scale, Frosted Scale (summer stages, Red Spider Mites, San Jose Scale, Soft Shelled Scales	2 to 4 qts.	Apply Volck Supreme Spray during the late Spring to mid-Summer only. Do not apply after husk split.
Ornamentals and Shade Trees Apple, Birch, Conifers (see NOTE), Dogwood, Flowering Peach, Flowering Cherry, Hawthorne, Holly, Magnolia, Maple, Oak, Privet, Pyracantha, Sycamores	Black Scale, Brown Apricot Scale, Oyster Shell Scale, Red Spider Mites and Eggs, San Jose Scale, Soft Brown Scale	1 gal.	NOTE: Conifer applications to blue spruce may cause discoloration. Do not use on conifers or ferns not tolerant to oil sprays.

FOLIAR APPLICATIONS - CONCENTRATE SPRAYS

Crop	Pests	Volck per Acre	Further Use Instructions
Almonds, Pistachio	Brown Mite, European Red Mite, Pacific Spider Mite, Two-Spotted Spider Mite	4 to 6 gals.	For Summer use with concentrate sprayers. Apply Volck Supreme Spray in 60 to 150 gals. water per acre. Do not apply to trees lacking adequate moisture.
Grapefruit, Lemons, Valencia Oranges	Citrus Red Mites	5 to 15 gals.	Apply Volck Supreme Spray as a concentrate spray application (mist sprayer) in 50 to 100 gals. water per acre. Apply to Valencia oranges only during the period from mid-July through August. Do not apply to lemons within 60 days of harvest. Concentrate spray applications require that the spray nozzles be discharged into an air stream pattern of high velocity (above 100 mph) and with sufficient volume to carry the spray through the tree. Ground speed should not exceed 1½ mph. Observe all label notes, relative to the use of spray oils on citrus. Use only equipment capable of applying concentrate sprays.
Pears (California Only)	Red Spider Mites	4 to 6 gals.	For Summer use with concentrate sprayers.. Apply Volck Supreme Spray in 90 to 120 gals. water per acre.

DORMANT OR DELAYED DORMANT - DILUTE SPRAYS

Crop	Pests	Volck per 100 gals. Water	Further Use Instructions
Almonds, Apricots, Cherries, Nectarines, Peaches, Pistachio, Plums, Prunes (See NOTE)	Aphids (eggs of), Black Scale, Brown Apricot Scale, European Fruit Lecanium, Frosted Scale, Italian Pear Scale, Oleander Scale, Olive Scale (Parlatoria Scale), Parthenolecanium Soft Scale, Putnam Scale, Red Spider Mites, San Jose Scale, Scale Insects (certain other), Soft Brown Scale	1½ to 2 gals.	For dormant or delayed dormant use only. NOTE: For plums and prunes, spray only during the delayed dormant period. Oil sensitive varieties (such as, but not limited to Moyer or Queen Ann) may develop dead limbs or scaffolds, which will reduce yield.
Apples	Rust Mites	1½ to 2 gals.	For delayed dormant use only.
Apples, Pears	Aphids (eggs of), Italian Pear Scale, Red Spider Mites, San Jose Scale	1½ to 2 gals.	For dormant or delayed dormant use only.
Figs	Fig Scale	3 gals.	For dormant or delayed dormant use only.
Grapes	Mealybug	See NOTE	For dormant use only. NOTE: Use 4 to 6 gals. in 200 to 300 gals. of water per acre.
Pears	Pear Psylla	1 to 1½ gals.	For delayed dormant use only.
	Blister Mite	1 to 1½ gals.	For delayed dormant use only.

DORMANT OR DELAYED DORMANT - CONCENTRATE SPRAYS

Crop	Pests	Volck per Acre	Further Use Instructions
Almonds, Apricots, Cherries, Nectarines, Peaches (see NOTE), Pistachio	Brown Apricot Scale, Mite Eggs, Olive Scale (Parlatoria Scale), Parthenolecanium Soft Scale, Peach Twig Borer, San Jose Scale, Walnut Scale	4 to 6 gals.	For dormant or delayed dormant use only. Apply Volck Supreme Spray in 40 to 125 gals. of water per acre. NOTE: Do not apply to oil sensitive varieties as phytotoxicity to buds and new wood may result.
Apples, Pears (except California)	Aphids (eggs of), European Red Mites (eggs of), Italian Pear Scale, San Jose Scale	3 gals.	For dormant or delayed dormant use only. For Volck Supreme Spray applications in 25 to 50 gals. of water per acre.
		4 gals.	For dormant or delayed dormant use only. For Volck Supreme Spray applications in 50 to 100 gals. of water per acre.
		5 gals.	For dormant or delayed dormant use only. For Volck Supreme Spray applications in 100 or more gals. of water per acre.
Apples, Pears (see NOTE)	European Red Mites (eggs of), Pear Psylla, San Jose Scale	4 to 6 gals.	For dormant or delayed dormant use only. Apply Volck Supreme Spray in 40 to 125 gals. of water per acre. NOTE: Do not apply to oil sensitive varieties as phytotoxicity to buds and new wood may result.
Pears (Washington Only)	Pear Psylla	3 to 4 gals. by Air	For dormant use only. Apply by air in sufficient water to make a minimum of 10 gals. of dilute spray per acre.
Plums, Prunes (see NOTE)	Black Scale, Brown Apricot Scale, Brown Mite Eggs, European Red Mites (eggs of), Mealy Plum Aphid Eggs, Olive Scale (Parlatoria Scale), Plum Aphid Eggs, San Jose Scale	3 to 5 gals.	Apply Volck Supreme Spray in 40 to 125 gals. of water per acre. NOTE: For plums and prunes, spray only during the delayed dormant period. Oil sensitive varieties (such as, but not limited to Moyer or Queen Ann) may develop dead limbs or scaffolds, which will reduce yield.

DORMANT - DILUTE SPRAYS

Crop	Pests	Volck per 100 gals. Water	Further Use Instructions
Ornamentals Shade Trees	Black Scale, Brown Apricot Scale, Oyster Shell Scale, Red Spider Mites and Eggs, San Jose Scale, Soft Brown Scale.	2 gals.	For dormant sprays on deciduous varieties.
Apple, Birch, Conifers (see NOTE), Dogwood, Flowering Cherry, Flowering Peach, Hawthorne, Holly, Magnolia, Maple, Oak, Privet, Pyracantha, Sycamores		1 gal.	For dormant sprays on other varieties. NOTE: Conifer application to blue spruce may cause discoloration. Do not use on conifers or ferns not tolerant to oil sprays.

SEED CROP USES

Crop	Pests	Volck per Acre	Further Use Instructions
Alfalfa (Seed Alfalfa Only)	Spider Mites	1 qt.	For use on seed alfalfa only. Apply in 20 to 30 gals. of water by ground application or 10 to 15 gals. of water by air with recommended rates of Kelthane 7EC.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing or pouring.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Refillable Container (110 gallons or greater): Refill this container with pesticides only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning the container before refilling is the responsibility of the refiller.

Container Disposal: To clean the container before final disposal, empty the remaining contents from this container into the production mix tank. Triple rinse the interior of the container with water by using a suitable hose and nozzle attachment or pressure rinser. Collect and store the rinsate for proper disposal. Then, offer the container for recycling or reconditioning, or puncture and dispose of in a sanitary landfill.

30 gallon Nonrefillable container: Do not reuse or refill this container.

Container Disposal: Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact Ag Container Recycling Council (ACRC) at 877-952-2272 or rperkins@acrecycle.org.

5 gallon Nonrefillable container: Do not reuse or refill this container.

Container Disposal: Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact Ag Container Recycling Council (ACRC) at 877-952-2272 or rperkins@acrecycle.org.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW, AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN.

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label.

EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS", AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.


Copyright© 2006 by Wilbur-Ellis Company

WILBUR-ELLIS Logo® and IDEAS TO GROW WITH® are registered trademarks of WILBUR-ELLIS Company.

Kelthane® is a registered trademark of Dow Agro Science. Volck® is a registered trademark of OMS Investments, Inc.

F-082108-1

Manufactured in U.S.A. by:
WILBUR-ELLIS COMPANY
PO BOX 16458 -- FRESNO, CA 93755
(559) 442-1220



Batch Code _____