

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Mrs. Sandi Escamilla Registration Specialist Wilbur-Ellis Company P.O. Box 1286 Fresno, CA 93715

JUN 1 6 2010

RE: Notification to Correct the Zip Code 93715 to 93755

EPA Registration Number: 2935-546 Date of Submission: May 13, 2010

Dear Mrs. Escamilla:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated, May 13, 2010, for the above mentioned product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P)

Office of Pesticide Programs

	((-			Print F	orm
Please read instructions	on reverse before comple	ting form.			Form Ap	Droved	. OMB No.	2070-00	SO, Approv	oi expires 2-28
SEPA	Environmenta	United States I Protection Ington, DC 20	_	ncy		×	Registr Amend Other		OPP Ider	ntifier Number
		Application	on for l	estic	ide - Sec	tion	l			
1. Company/Product Nur 2935-546				1	Product Man	ager		\ <u></u>	roposed Cla	
4. Company/Product (Na PN All Sea	me) ason Spray Oil			PM#	21		-		None	Restricte
Wilbur-Ellis PO Box 128 Fresno CA	36	odej		(b)(i), i to: EPA Prodi	ny product Reg. No uct Name	is sim	ilar or iden	tical in co	omposition	ection 3(c)(3) and labeling
	· · · · ·		Sec	tion -	11					
X Notification - Exp Explanation: Use add	response to Agency letter		n I and Se	ction II.)	Final printe Agency lett "Me Too" / Other - Exp	ter dat Applica	ation.	NOTIF	FICATIO 1 6 2010	_
			Sect	ion - l	11			~		
1. Material This Product	Will Be Packaged In							÷ .		
Child-Resistant Packagin Yes No * Certification must be submitted	Unit Packaging Yes No	No. per . container	Water	Yes Jext	No. per contains	······································	2. Type of	Metal Plastic Glass Paper	r Specify)	
3. Location of Net Conte	nts Information Container	4. Size(s) Re	tail Contain	ner		5. Lo	cation of La On Label On Label		ons paning produ	oct
6. Manner in Which Labe	l is Affixed to Product	Lithor Paper Stenc	raph glued iled		Othe	·				
			Sect	ion - I	V					
1. Contact Point /Comp	lete items directly below	for identification	on of indivi	dual to b	e contected,	if nec	essary, to p	rocess this	s application).J
Name Sandi Escami	illa		Title	Registra	tion Speciali	st		Telephor 559-493		ide Area Code)
•	tatements I have made or t any knowlingfly false or ble law.		all attach				mprisonme	nt or	6. Date A Receive	• •
2. Signature	Lacamel.		3. Title R	egistrati	on Specialis	t	(((((((((((((((((((((((

5. Date

May 13, 2010

Sandi Escamilla

.....



May 13, 2010

CERTIFIED MAIL
Return Receipt Requested
70091410000051302719

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
1200 Pennsylvania Ave. NW
Washington DC 20460

Subject: Notification: PN All Season Spray Oil

EPA Reg. No. 2935-546

Dear Administrator:

Wilbur-Ellis Company is submitting this notification to EPA to correct the zip code 93715 to 93755.

Enclosed are four (4) finial printed labels.

Thank you for your attention to this matter.

Sincerely

Sandi Escamilla

Registration Specialist Wilbur-Ellis Company

Enclosure(s)



PN ALL SEASON SPRAY OIL

(Insecticide – Miticide) NOTIFICATION

Petroleum Oil Min. Unsulfonated Residue – 99% Petroleum Oil Classification – Unclassified

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Causes moderate eye irritation.

FIRST AID Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. lf on skin or clothing: Call a poison control center or doctor for treatment advice. If in eyes: - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If swallowed: - Immediately call a poison control center or doctor. - Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. If inhaled: Move person to fresh air - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-892-0099 for emergency medical treatment information.

NOTE TO PHYSICIANS: Contains petroleum distillate - vomiting may cause aspiration pneumonia.

EPA Reg. No. 2935-546

EPA Est. No. 2935-CA-1^F, 2935-CA-19^R, 2935-CA-31^H, 2935-TX-1^E

Superscript used is the first letter of the lot number.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: Long-sleeved shirt and long pants, and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS

Do not weld, heat or drill this container. Residues may ignite with explosive violence if heated sufficiently.

USER SAFETY REQUIR ENTS

Users should:

- (1) Wash hands before eating, drinking, chewing gum, using tobacco or using the tollet.
- (2) Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry

Interval (REI) of 4 hours.

PPE required for early entry to treated area that is permitted under the Worker

Protection Standard and that involves contact with anything that has been treated,
such as plants, soil, or water, is: coveralls, waterproof gloves, and shoes plus socks.

DIRECTIONS

PN All Season Spray Oil is an especially refined Phytonomic Oil Contact Insecticide. It is for use on plants in foliage or the dormant stage.

When using this product in combination with other pesticides as directed on this label observe appropriate cautions, limitations for the product involved.

For foliage applications, the spray tank must be free of any sulfur residue before using PN All Season Spray Oil. Do not combine sulfur sprays with PN All Season Spray Oil on foliage and do not follow a previous application of sulfur or apply sulfur following PN All Season Spray Oil until sufficient time has elapsed for safety (at least 30 days). If in doubt, spray a portion of one tree previously treated with sulfur, using 1 qt. to 25 gals. water. If sufficient time has not elapsed, burn will show in several days.

Do not use PN All Season Spray Oil when temperatures are above 90° Fahrenheit. When temperatures drop below 32° Fahrenheit (usually nighttime lows), do not spray tree and vine crops until daytime temperatures have warmed to a point above 35° Fahrenheit and are free from moisture due to melting frost or heavy dews. Do not use on foliage closer than 60 days after or 90 days before using captan or folpet products.

Application to trees weakened by disease, drought, drying winds, or high nitrogen applications may result in oil injury.

Do not apply this product through any type of irrigation system.

AIR APPLICATION

Apply quantity of product shown for each listed crop for control of listed insects in sufficient water to make a minimum of 20 gals, dilute spray per acre by air application. Follow timing shown for each crop as shown on container label. Use only as emergency application when soil conditions do not permit regular ground application.

NOTE: Do not apply PN All Season Spray Oil concentrate to Queen Ann variety or othe oil sensitive varieties as phytotoxicity to buds and new wood will result.

INFORMATION: Observe on all crops and uses — oil effectiveness in dormant and delayed dormant applications is determined by thoroughness of coverage. Concentrate applications may reduce coverage and effectiveness. Oil applications made to trees that are weak or suffering from moisture stress or when slow drying of spray occurs may cause bud, twig or leaf injury.

Phytotoxicity studies have not been made on all fruit varieties, or in all growing areas in California for concentrate applications. Concentrate spraying is a more exacting technique of spray application than dilute. Concentrate spray equipment must be properly calibrated and operated to obtain desired results.

MIXING DIRECTIONS

Check this product for proper emulsification by mixing a small quantity of it with water in a small container before filling the entire spray tank. Correct emulsification is indicated when a uniform mixture, with a blue color, results on shaking. If this mixture is not a milky emulsion after shaking, do not use the remaining product. With agitator running, start filling spray tank with water. When combining with wettable powder insecticide or fungicide products, add these first when tank is 1/3 filled. Add PN All Season Spray Oil when tank is about 1/2 full and while there is sufficient agitation of the spray to ensure complete emulsification.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof unfit for sale, use or consumption.

NOTE: Multiple concentrate applications may cause wood Injury. Double applications can occur on the ends of the row during turning – turn off sprayer to avoid multiple applications. Equipment must be capable of giving thorough coverage for maximum insecticidal effectiveness. If San Jose Scale is a moderate to severe problem, use dilute application. Concentrate applications may delay bud development. The following conditions may cause oil injury: cool, damp, or extremely dry weather; broken emulsions and/or faulty application. Follow state recommendations and cautions.

Apples, Pears Red Spider Italian Pear San Jose S Grapes Mealybug Grapefruit. Black Scale Purple Sca Lemons Scale, Red Mites Yellor Navel Oranges Black Scale Purple Sca Mites Vella Black Scale Olives Oleander S Olive Scale (Parlatoria Spider Mite Mitoe (Not registered for use in California) Powderv M Do not use Do not spra are under t Black Scale (California Only) Pear Psylla Pears Blister Mite Black Scale Purple Sca Scale, Red Mites, Yello Walnuts European i Scale, Fros Scale (summer si Red Spider San Jose S Soft Shelle Scales Ornamentals and Black Scali Shade Trees Apricot Sca Ovster She Apple, Birch, Conifers (see Red Spide and Eggs, NOTE), Dogwood, Jose Scale Flowering Peach, Brown Sca Flowering Cherry, Magnolia, Maple Oak, Privet,

Sycamores

FOLIAR AP

Pest

Brown Apri

Scale, San

Pacific Spk

Mites, Red

Rust Mites

Scale

Mites

Crop

Almonds,

Apricots,

Chemies.

Nectarines

Peaches, Pistachio, Plums.

Apples

Manufactured in U.S.A by: WILBUR-ELLIS COMPANY PO BOX 16458 – FRESNO, CA 93755

F-020408-3 • K-040110

USER SAFETY REQUIR

Users should:

interval (REI) of 4 hours.

Wt.

0% 0%

0%

N

tes.

o 20

าคก

rol

tvice

alling

ontact

ause

ι-19^R

TX-1E

ımber

ants.

such

where

Do not

olence

(1) Wash hands before eating, drinking, chewing gum, using tobacco or using the

ENTS

(2) Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry

PPE required for early entry to treated area that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, and shoes plus socks.

DIRECTIONS

PN All Season Spray Oil is an especially refined Phytonomic Oil Contact Insecticide. It is for use on plants in foliage or the dormant stage.

When using this product in combination with other pesticides as directed on this label, observe appropriate cautions, limitations for the product involved.

For foliage applications, the spray tank must be free of any sulfur residue before using PN All Season Spray Oil. Do not combine sulfur sprays with PN All Season Spray Oil on foliage and do not follow a previous application of sulfur or apply sulfur following PN All Season Spray Oil until sufficient time has elapsed for safety (at least 30 days). If in An obastri Spray a portion of one tree previously treated with suffur, using 1 qt. to 25 gals, water. If sufficient time has not elapsed, burn will show in several days.

Do not use PN All Season Spray Oil when temperatures are above 90° Fahrenheit. When temperatures drop below 32° Fahrenheit (usually nighttime lows), do not spray tree and vine crops until daytime temperatures have warmed to a point above 35° Fahrenheit and are free from moisture due to melting frost or heavy dews. Do not use on foliage closer than 60 days after or 90 days before using captan or folpet products.

Application to trees weakened by disease, drought, drying winds, or high nitrogen applications may result in oil injury.

Do not apply this product through any type of irrigation system.

AIR APPLICATION

Apply quantity of product shown for each listed crop for control of listed insects in sufficient water to make a minimum of 20 gals, dilute spray per acre by air application. Follow timing shown for each crop as shown on container label. Use only as emergency application when soil conditions do not permit regular ground

NOTE: Do not apply PN All Season Spray Oil concentrate to Queen Ann variety or other oil sensitive varieties as phytotoxicity to buds and new wood will result.

INFORMATION: Observe on all crops and uses - oil effectiveness in dormant and delayed dormant applications is determined by thoroughness of coverage. Concentrate applications may reduce coverage and effectiveness. Oil applications made to trees that are weak or suffering from moisture stress or when slow drying of spray occurs may cause bud, twig or leaf injury.

Phytotoxicity studies have not been made on all fruit varieties, or in all growing areas in California for concentrate applications. Concentrate spraying is a more exacting technique of spray application than dilute. Concentrate spray equipment must be properly calibrated and operated to obtain desired results.

MIXING DIRECTIONS

Check this product for proper emulsification by mixing a small quantity of it with water in a small container before filling the entire spray tank. Correct emulsification is indicated when a uniform mixture, with a blue color, results on shaking. If this mixture is not a milky emulsion after shaking, do not use the remaining product. With agitator running, start filling spray tank with water. When combining with wettable powder insecticide of fungicide products, add these first when tank is 1/3 filled, Add PN All Season Spray Oil when tank is about 1/2 full and while there is sufficient agitation of the spray to ensure complete emulsification.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof unfit for sale, use or

NOTE: Multiple concentrate applications may cause wood injury. Double applications can occur on the ends of the row during turning - turn off sprayer to avoid multiple applications. Equipment must be capable of giving thorough coverage for maximum insecticidal effectiveness. If San Jose Scale is a moderate to severe problem, use dilute application. Concentrate applications may delay bud development. The following conditions may cause oil injury: cool, damp, or extremely dry weather; broken emulsions and/or faulty application. Follow state recommendations and cautions.

A by: WILBUR-ELLIS COMPANY

58 – FRESNO, CA 93755

020408-3 • K-040110

	i i	PN AII	
Сгор	Pests	Season per 100 gallons water	Further Use Instructions
Almonds, Apricots, Cherries,	Brown Apricot Scale, San Jose Scale	1 gal.	For summer stages of Brown Apricot and San Jose Scale.
Nectarines, Peaches, Pistachio, Plums, Prunes	Pacific Spider Mites, Red Spider Mites	3 to 4 qts.	For summer use on Red Spider Mites.
Apples	Rust Mites	1 to 1½ ga/s.	For summer use.
Apples, Pears	Red Spider Mites	3 to 4 ats.	For summer use.
	Italian Pear Scale, San Jose Scale	1 to 1½ gals.	Apply PN All Season Spray Oil to summer forms of San Jose and Italian Pear Scale.
Grapes	Mealybug	See NOTE	For (Post Harvest), fall spray use only. NOTE: Use 4 to 6 gals. in 200 to 300 gals. of water per acre.
Grapetruit, Lemons	Black Scale, Purple Scale, Red Scale, Red Spider Mites, Yellow Scale	1 to 1¼ gals.	For best results, apply PN All Season Spray Oil during winter and spring months when Red Scale are more easily controlled and trees generally are less reactive to oil sprays. Do not use within 8 weeks of harvest on lemons.
Navel Oranges	Black Scale, Purple Scale, Red Scale, Red Spider Mites, Yellow Scale	3 qts.	Do not use on Navel oranges after September 1.
Olives	Black Scale, Oleander Scale, Olive Scale (Parlatoria Scale), Spider Mites	1½ gals.	Apply PN All Season Spray Oil in the prebloom or in postbloom up to buckshot growth stage.
Hops (Not registered for use in California)	Mites	1 to 2 gals.	Apply PN All Season Spray Oil with ground equipment only. Thorough cover- age is essential. Use 50 to 200 gals. of dilute spray per acre depending upon plant size. Make applications as needed.
	Powdery Mildew	1 to 2 gals.	Apply PN All Season Spray Oil with ground equipment only. Thorough coverage is essential. Use 50 to 200 gals, of dilute spray per acre depending upon plant size. Make first application at early leaf stage. Repeat every 7 to 14 days depending upon new growth. Use shorter interval and/or the higher rate when the conditions for infection are more severe.
		ige or if lempera	following an oll spray. fure is above 90° Fahrenheit or if plants
Olives (California Only)	Black Scale	1 to 1/2 gals.	Apply 400 to 800 gals, dilute spray per acre. Make applications between July 1 and August 31. Higher concentrations of PN All Season Spray Oil than noted above may cause phytotoxicity.
Pears	Pear Psylla	1 to 1½ gais.	For summer use.
	Blister Mite	1 to 1½ gals.	For fail spray use only.
Valencia Oranges	Black Scale, Purple Scale, Red Scale, Red Spider Mites, Yellow Scale	1 gal.	Do not apply to Valencia oranges after September 1.
Walnuts	European Fruit Scale, Frosted Scale (summer stages, Red Spider Mites, San Jose Scale, Soft Shelled Scales	2 to 4 qts.	Apply PN All Season Spray Oil during the late spring to mid-summer only. Do not apply after husk split.
Omamentals and Shade Trees Apple, Birch, Confibrs (see MOTE), Dogwood, Flowering Peach, Flowering Cherry, Hawthorne, Holly, Magnolia, Maple, Oak, Privet, Pyracantha, Sycamores	Black Scale, Brown Apricot Scale, Oyster Shell Scale, Red Spider Mites and Eggs, San Jose Scale, Soft Brown Scale	1 gal.	NOTE: Conifer applications to blue spruce may cause discoloration. Do not use on conifers or terms not tolerant to oil sprays.

_	FOLIAR	APPLIC S
	Сгор	Pests
	Almonds, Pistachio	Brown Mite, European Red Mite, Pacific Spi Mite, Two-Spotte Spider Mite
	Grapefruit, Lemons, Valencia Oranges	Citrus Red Mite

Red Spider Mit-(California Only)

DORMANT OR DEL

	Crop	Pests
	Almonds, Apricots, Apricots, Cherries, Nectarines, Peaches, Pistachio, Pilums, Prunes (See NOTE)	Aphids (eggs o Black Scale, Br Apricot Scale, European Fruit Lecanium, Fros Scale, Olive Sc (Parlatoria Scale, Olive Sc (Parlatoria Scale, Put Scale, Red Spi Mites, San Jos Scale, Scale Insects (certair other), Soft Brc Scale
	Apples	Rust Mites
	Apples, Pears	Aphids (eggs c Italian Pear Sc Red Spider Mil San Jose Scal
	Figs	Fig Scale
	Grapes	Mealybug
1	Pears	Pear Psylla
		Blister Mite
	Ī	ORMANI

CO

Crop	Pests
Almonds, Apricots, Cherries, Nectarines, Peaches (see NOTE), Pistachio	Brown Apricot Scale, Mite Eç Olive Scale (Parlatoria Sc Parthenolecai Soft Scale, Pe Twig Borer, Sa Jose Scale, W Scale
Apples, Pears (except California)	Aphids (eggs European Re Mites (eggs o Italian Pear S San Jose Sca

E OI TIATO
Further Use Instructions
mer stages of Brown Apricot 1 Jose Scale.
nmer use on Red Spider Mites.
ımer use.
ımer use.
N All Season Spray Oil to r forms of San Jose and Italian ale.

st Harvest), fall spray use only.
Use 4 to 6 gals. in 200 to 300
water per acre.

t results, apply PN All Season il during winter and spring when Red Scale are more easily ed and trees generally are less to oil sprays. Do not use within s of harvest on lemons.

use on Navel oranges after iber 1.

'N All Season Spray Oil in the m or in postbloom up to buckshot stage.

'N All Season Spray Oil with equipment only. Thorough coverissential. Use 50 to 200 gals. of pray per acre depending upon ze. Make applications as needed.

¹N All Season Spray Oil with equipment only. Thorough je is essential. Use 50 to 200 dilute spray per acre depending ant stze.

rst application at early leaf stage. every 7 to 14 days depending aw growth. Use shorter interval the higher rate when the condir infection are more severe.

ı an oil spray. ove 90° Fahrenheit or il plants

00 to 800 gals, dilute spray per ake applications between July 1 gust 31. Higher concentrations Il Season Spray Oil than noted nay cause phytotoxicity.

nmer use.

spray use only.

apply to Valencia oranges after

N All Season Spray Oil the late spring to mid-summer not apply after husk split.

Conifer applications to blue may cause discoloration. Do not conifers or ferns not tolerant to

FULIAN	APPLIC O	ICENTRALE SPRATS	
Сгор	Pests	PN All Season per Acre	Further Use Instructions
Almonds, Pistachio	Brown Mite, European Red Mite, Pacific Spider Mite, Two-Spotted Spider Mite	4 to 6 gals.	For summer use with concentrate sprayers. Apply PN All Season Spray Oil in 60 to 150 gals. water per acre. Do not apply to trees lacking adequate moisture.
Grapefruit, Lemons, Valencia Oranges	Citrus Red Mites	5 to 15 gals.	Apply PN All Season Spray Oil as a concentrate spray application (mist sprayer) in 50 to 100 gals. water per acre. Apply to Valencia oranges only during the period from mid-July through August. Do not apply to lemons within 60 days of harvest. Concentrate spray applications require that the spray droplets be discharged into an air stream pattern of high velocity (above 100 mph) and with sufficient volume to carry the spray through the tree. Ground speed should not exceed 1½ mph. Observe all label notes, relative to the use of spray oils on citrus. Use only equipment capable of applying concentrate sprays.
Pears (California Only)	Red Spider Mites	4 to 6 gals.	For summer use with concentrate sprayers. Apply PN All Season Spray Oil In 90 to 120 oals, water per acre.

DORMANT OR DELAYED DORMANT - DILUTE SPRAYS

Сгор	Pests	PN Ali Season per 100 gallons water	Further Use Instructions
Almonds, Apricots, Cherries, Nectarines, Peaches, Pistachio, Plums, Prunes (See NOTE)	Aphids (eggs of), Black Scale, Brown Apricot Scale, European Fruit Lecanium, Frosted Scale, Italian Pear Scale, Oleander Scale, Olive Scale (Parlatoria Scale), Parthenolecanium Soft Scale, Purlnam Scale, Red Spider Mites, San Jose Scale, Scale Insects (certain other), Soft Brown Scale)	1½ to 2 gals.	For dormant or delayed dormant use only. NOTE: For plums and prunes, spray only during the delayed dormant period. Oil sensitive varieties (such as, but not limited to Moyer or Queen Ann) may develop dead limbs or scaffolds, which will reduce yield.
Apples	Rust Mites	11½ to 2 gals.	For delayed dormant use only.
Apples, Pears	Aphids (eggs of), Italian Pear Scale, Red Spider Mites, San Jose Scale	1½ to 2 gals.	For dormant or delayed dormant use only.
Figs	Fig Scale	3 gals.	For dormant or delayed dormant use only.
Grapes	Mealybug	See NOTE	For dormant use only. NOTE: Use 4 to 6 gals. in 200 to 300 gals. of water per acre.
Pears	Pear Psylla	1 to 11/2 gais.	For delayed dormant use only.
	Blister Mite	1 to 11/2 gals.	For delayed dormant use only.

DORMANT OR DELAYED DORMANT -CONCENTRATE SPRAYS

Crop	Pests	PN All Season per acre	Further Use Instructions
Almonds, Apricots, Cherries, Nectarines, Peaches (see NOTE), Pistachio	Brown Apricot Scale, Mite Eggs, Olive Scale (Parlatoria Scale), Parthenolecanium Soft Scale, Peach Twig Borer, San Jose Scale, Walnut Scale	4 to 6 gals.	For dormant or delayed dormant use only, Apply PN All Season Spray Oil in 40 to 125 gals, of water per acre, NOTE: Do not apply to oil sensitive varieties as phytotoxicity to buds and new wood may result.
Apples, Pears (except California) Aphids (eggs of), European Red Mites (eggs of), Italian Pear Scale, San Jose Scale	3 gals.	For dormant or delayed dormant use only. For PN All Season Spray Oil applications in 25 to 50 gals. of water per acre.	
	4 gals.	For dormant or delayed dormant use only. For PN All Season Spray Oil applications in 50 to 100 gals. of water per acre.	
		5 gals.	For dormant or delayed dormant use only. For PN All Season Spray Oil applications in 100 or more gals. of

			<u> </u>
Apples, Pears (see NOTE)	European Red Mites (eggs of), Pear Psylla, San Jose Scate	4 to 6 gals.	For dormant or delayed dormant use only Apply PN All Season Spray Oil in 40 ½ 125 gals of water per acre. NOTE: Do not apply to oil sensitive varieties as phytotocics to buds and new wood may result.
Pears (Washington Only)	Pear Psylla	3 ta ∕gals. Sy air	For desmant, se only. Apply by air in sufficient water to make a minimum of 10 fals, of flinds spray per acre.
Plums, Prunes (see NOTE)	Black Scale, Brows Apricot Scale, Brown Mite Eggs, *, European Red Mites (eggs of), Meaty Plum Aphid Eggs, Olive Scale (Partatoria Scale), Plum Aphid Eggs, San Josa Scale	3 to 5 gals.	Apply PN All Season in 40 to 125 gals. of water pur acre. NOTE: For plums and prunes, spray only during the delayed dorman priod 201 sensitive varieties (such as, Lut and limited to Moyer or Queen Ann) may develop dead limbs or scaffolds, which will reduce yield.

DORMANT - DILUTE SPRAYS

Crop	Peste	PN All Season per 100 gallons water	Further Use Instructions
Ornamentals Shade Trees Apple, Birch, Conifers (see NOTE), Dogwood,	Black Scale, Brown Apricot Scale, Oyster Shell Scale, Red Spider Mites and Eggs, San Jose Scale, Soft	2 gals.	For dormant sprays on deciduous varieties.
Flowering Cherry, Flowering Peach, Hawthorne, Holly, Magnolia, Maple, Oak, Privet, Pyracantha, Sycamores	Brown Scale.	1 gal.	For dormant sprays on other varieties. NOTE: Conifer application to blue spruce may cause discoloration. Do not use on conifers or ferns not tolerant to oil.

SEED CROP USES

Crop	Pests	PN Ali Season per 100 gallons water	Further Use Instructions
Alfalfa (Seed Alfalfa Only)	Spider Mites	1 qt.	For use on seed alfalfa only. Apply in 20 to 30 gals. of water by ground application or 10 to 15 gals. of water by air with recommended rates of Kelthane7EC®.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Protect from water contamination during storage and transfer. Contamination with water may make this product unfit for its intended use. Always check emulsification (see Mixing Directions) before use. For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-892-0099.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW, AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN.

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without timitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Selter harmless from any claims relating to such factors.

Relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer. Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. The EXCLUSIVE REMEDY OF THE BUYER OF USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY CONTRACT, REGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

Copyright® 2006 by Wilbur-Ellis Company, WiLBUR-ELLIS Logo® is a registered trademark of WILBUR-ELLIS Company, Kelthane® is a registered trademark of DowAgroScience.