



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

September 10, 2025

Lynn Georges
Director of Regulatory Affairs
Wilbur-Ellis Company LLC
4582 S. Ulster Street, Suite 400
Denver, CO 80237

Subject: Label Amendment - Registration Review Mitigation for Permethrin
Product Name: Stiletto
EPA Registration Number: 2935-551
Case Number: 471540
Application Date: February 8, 2022

Dear Lynn Georges:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Permethrin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label

printing. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Christian Bongard by phone at (202) 566-2248, or via email at Bongard.Christian@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Jaclyn Pyne". The signature is written in a cursive, flowing style.

Jaclyn Pyne, Team Leader
Risk Management and Implementation Branch 3
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

ACCEPTED

09/10/2025

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 2935-551

PERMETHRIN

GROUP

3A

INSECTICIDE

**STILETTO®**

For the control of armyworm, cricket, cutworm and darkling beetle

ACTIVE INGREDIENT:

Permethrin * **0.5%

OTHER INGREDIENTS:99.5%

TOTAL:100.0%

*(3-Phenoxyphenyl)methyl(±)-cis, trans 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate

**Cis/trans isomers ratio: Min. 35% (±) cis; Max. 55% (±) cis

Min. 45% (±) trans; Max. 65% (±) trans

EPA Reg. No. 2935-551

EPA Est. No. 66876-OR-001

**KEEP OUT OF REACH OF CHILDREN
CAUTION****FIRST AID**

If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.

Have product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the following EMERGENCY TELEPHONE NUMBERS:
(800) 424-9300 CHEMTREC (transportation and spills)
(800) 222-1222 POISON CONTROL CENTER (human health)
(888) 426-4435 ASPCA (animal health)

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Wear long-sleeved shirt, long pants and socks and shoes.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate and butyl rubber \geq 14 mils.

All loaders, applicators, and other handlers must wear:

- Long-sleeve shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves for all handlers, except for applicators using motorized ground equipment, pilots, and flaggers.

See Engineering Controls for more requirements.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (6)].

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Under some conditions, it may also have a potential for transport into surface water in runoff (primarily adsorbed to suspended soil particles), for several months or more after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, and areas overlying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlying tile drainage systems that drain to surface waters.

NON-TARGET ORGANISM ADVISORY STATEMENT: This pesticide is highly toxic to bees and other pollinating insects exposed to direct treatment or to residues in/on blooming crops or weeds. **Protect pollinating insects by following label directions intended to minimize drift and reduce pesticide risk to these organisms.**

How to Report Bee Kills: It is recommended that users contact both the state lead agency and the U.S. Environmental Protection Agency to report bee kills due to pesticide application. Bee kills can be reported to EPA at beekill@epa.gov. To contact your state lead agency, see the current listing of state pesticide regulatory agencies at the National Pesticide Information Center's website: http://npic.orst.edu/reg/state_agencies.html

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, are:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material.

Unless otherwise directed by supplemental labeling, follow the Directions for Use as shown in each crop group section. STILETTO is a pelletized formulation for the control of armyworm, cricket, cutworm and darkling beetle. This product may be applied by hand equipment, ground equipment or aircraft. Apply when insects appear or feeding is noticed. The higher rate should be used as pest populations increase. With the exception of crops listed on this label rotational crops cannot be planted within 60 days of last application.

INSECT RESISTANCE MANAGEMENT

For resistance management, STILETTO contains a Group 3A insecticide. Any insect population may contain individuals naturally resistant to this product and other Group 3A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of STILETTO or Group 3A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area. For further information or to report suspected resistance contact Wilbur-Ellis Company, LLC at 720-306-6340.

VEGETATIVE FILTER STRIPS

Construct and maintain a vegetative filter strip, according to the width specified below, of grass or other permanent vegetation between the field edge and nearby down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; streams, marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing permethrin onto fields where a maintained vegetative filter strip of **at least 25 feet** exists between the field edge and where a down gradient aquatic habitat exists. This minimum required width of 25 feet may be reduced or removed under the following conditions:

- For Western irrigated agriculture, a maintained vegetative filter strip of at least 10 feet wide is required. Western irrigated agriculture is defined as irrigated farmland in the following states: WA, OR, CA, ID, NV, UT, AZ, MT, WY, CO, NM, and TX (west of I-35).
 - For Western irrigated agriculture, if a sediment control basin is present, a vegetative filter strip is not required.
- In all other areas, a vegetative filter strip with a minimum width of 25 feet is required, unless the following conditions are met. The vegetative filter strip requirement may be reduced from 25 feet to 15 feet if at least one of the following applies:
 - The area of application is considered prime farmland (as defined in 7 CFR Section 657.5).
 - Conservation tillage is being implemented on the area of application. Conservation tillage is defined as any system that leaves at least 30% of the soil surface covered by residue after planting. Conservation tillage practices can include mulch-till, no-till, or strip-till.
 - A functional terrace system is maintained on the area of application.
 - Water and sediment control basins for the area of application are functional and maintained.
 - The area of application is less than, or equal to, 10 acres.

For further guidance on vegetated filter strips, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses Natural Resources Conservation Services. <https://www.regulations.gov/document/EPA-HQ-OPP-2008-0331-0175>

BUFFER ZONES TO WATER BODIES

Ground Application

- Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

USE RATES AND RESTRICTIONS

Note: The preharvest interval is shown after each crop heading in parenthesis ().

ALFALFA, ALFALFA GROWN FOR SEED (0 day PHI*)		
Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-40 lb/acre (0.1-0.2 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
*Do not apply within 14 days of harvest for rates greater than 0.1 lbs. a.i. (20 lbs. product) per acre. • Do not apply to mixed stands with intentionally grown forage grasses and/or legumes. • Retreatment interval is 30 days. • For seed alfalfa, make applications early morning or late evening when bees are not actively foraging.		

ARTICHOKE (0 day PHI)		
Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-60 lb/acre (0.1-0.3 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
• Do not apply more than 1.5 lbs. a.i. (300 lbs. product) per acre per season. • Retreatment interval is 10 days. • Avoid application when honey bees are actively foraging by applying during early morning or during evening hours.		

ASPARAGUS (1 day PHI)		
Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	10-20 lb/acre (0.05-0.10 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 0.4 lbs. a.i. (80 lbs. product) per acre per season. Retreatment interval is 7 days. Avoid application when honey bees are actively foraging by applying during early morning or during evening hours. 		

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, CHINESE BROCCOLI (gai lon, white flowering broccoli), BROCCOLI RAAB (rapini) (1 day PHI)		
Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	10-20 lb/acre (0.05-0.10 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 0.8 lbs. a.i. (160 lbs. product) per acre per season. Retreatment interval is 5 days. Avoid application when honey bees are actively foraging by applying during early morning or during evening hours. 		

CABBAGE; CABBAGE, CHINESE (napa) (tight-heading varieties only) (1 day PHI)		
Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-40 lb/acre (0.1-0.2 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 0.4 lbs. a.i. (80 lbs. product) [0.8 lbs. a.i. (160 lbs. product) for Hawaii] per acre per season. Retreatment interval is 5 days. Avoid application when honey bees are actively foraging by applying during early morning or during evening hours. 		

COLLARDS and TURNIPS (1 day PHI)		
Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	10-20 lb/acre (0.05-0.1 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 0.8 lbs. a.i. (160 lbs. product) per acre per season. Retreatment interval is 3 days. Do not graze treated areas or feed crop to livestock. For use on collards in AR, AZ, GA, IL, IN, NC, OK, SC, and TX only. For use on turnips in FL, GA, IL, IN, OK, SC, TX and WA. Avoid application when honey bees are actively foraging by applying during early morning or during evening hours. 		

CORN (FIELD), FIELD CORN GROWN FOR SEED, POPCORN		
Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-40 lb/acre (0.1-0.2 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 0.45 lbs. a.i. (90 lbs. product) per acre per season. Retreatment interval is 7 days. Forage may be harvested on the day of application. Do not make treatments less than 6 days apart or apply less than 30 days prior to harvest of grain or fodder. 		

CORN, SWEET (1 day PHI)		
Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-40 lb/acre (0.1-0.2 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 1.2 lbs. a.i. (240 lbs. product) per acre per season. Retreatment interval is 3 days. 		

CUCURBIT VEGETABLES (1 day PHI)		
Chayote (fruit); Chinese waxgourd (Chinese preserving melon); Citron melon; Cucumber; Gherkin; Gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); <i>Momordica</i> spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); Muskmelon (includes true cantaloupe, cantaloupe, casaba, crenshaw, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); Pumpkin; Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); Squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); Watermelon (includes hybrids and/or varieties of <i>Citrullus</i> spp.).		
Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-40 lb/acre (0.1-0.2 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 1.6 lbs. a.i. (320 lbs. product) per acre per season. Retreatment interval is 7 days. Avoid application when honey bees are actively foraging by applying during early morning or during evening hours. 		

EGGPLANT (3 day PHI)		
Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-40 lb/acre (0.1-0.2 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 0.6 lbs. a.i. (120 lbs. product) [1.0 lbs. a.i. (200 lbs. product for Hawaii)] per acre per season. Retreatment interval is 7 days. Avoid application when honey bees are actively foraging by applying during early morning or during evening hours. 		

ONIONS, BULB and GARLIC (0 day PHI)		
Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-60 lb/acre (0.1-0.3 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 2.0 lbs. a.i. (400 lbs. product) per acre per season. 		

LEAFY VEGETABLE (EXCEPT BRASSICA) CROP GROUPING (1 day PHI)

Including all varieties and hybrids of Amaranth, Arugula, Cardoon, Celery, Celtuce, Chervil, Chinese celery, Chrysanthemum (edible-leaved and garland), Corn salad, Cress (garden and upland), Dandelion, Dock, Endive, Florence fennel, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio, Rhubarb, Swiss chard, Spinach

Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-40 lb/acre (0.1-0.2 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 0.8 lbs. a.i. (160 lbs. product) [1.2 lbs. a.i. (240 lbs. product) for Hawaii] per acre per season. Retreatment interval is 7 days. For spinach, do not apply more than 1.0 lbs. a.i. (200 lbs. product) per acre per season. Avoid application when honey bees are actively foraging by applying during early morning or during evening hours. 		

ORNAMENTAL NURSERY STOCK (field grown)

Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-40 lb/acre (0.1-0.2 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.

PEPPERS, BELL (3 day PHI)

Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-40 lb/acre (0.1-0.2 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 1.6 lbs. a.i. (320 lbs. product) per acre per season. Retreatment interval is 5 days. Avoid application when honey bees are actively foraging by applying during early morning or during evening hours. 		

POTATOES (14 day PHI)

Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-40 lb/acre (0.1-0.2 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 1.6 lbs. a.i. (320 lbs. product) per acre per season. Retreatment interval is 10 days. 		

TOMATOES, TOMATILLOS (0 day PHI)

Insects Controlled	Rate of Application	Comments
Armyworms Crickets Cutworms Darkling beetle	20-40 lb/acre (0.1-0.2 lb ai/acre)	Use higher recommended dosage for increased pest pressure or for increased residual pest control.
<ul style="list-style-type: none"> Do not apply more than 0.6 lbs. a.i. (120 lbs. product) [0.8 lbs. a.i. (160 lbs. product) for Hawaii] per acre per season. Retreatment interval is 7 days. Do not use on cherry tomatoes or other varieties in which the mature tomatoes will be less than 1 inch in diameter. 		

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container only and keep sealed. Store in a cool, dry and closed storage area. Keep container tightly closed when not in use. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Use caution when moving, opening, closing or pouring.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable Container: Do not reuse or refill this container. Completely empty bag into application equipment then offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW, AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN.

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS", AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR-ELLIS® logo, STILETTO® and INSECTICIDE® logo are registered trademarks of Wilbur-Ellis Company LLC.

NET CONTENTS: 50 POUNDS

Batch Code: _____

2025-0828

Manufactured for:
WILBUR-ELLIS COMPANY LLC
4582 S. Ulster Street, Suite 400, Denver, CO 80237



WILBUR-ELLIS®