

1/16/2015
UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

1/20

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

JAN 16 2015

Anne Downs
Senior Registration Specialist
Wilbur-Ellis Company
2903 S. Cedar Avenue
Fresno, CA 93725

Subject: Label Notification per PRN 98-10 – Addition of Logo and Chemical
Classification Language
Product Name: Base Camp LV 6
EPA Registration Number: 2935-553
Application Date: October 23, 2014
Decision Number: 497399

Dear Ms. Downs:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, you may contact Terri Stowe at (703) 305-6117 or via email at stowe.terri@epa.gov.

Sincerely,

Kathryn V. Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Redline Copy
Only added
No other changes

NOTIFICATION
JAN 16 2015

PULL HERE TO OPEN
PRESS TO RESEAL

E-1000109-014-00393

FIRST AID (cont.)

POTENTIAL NUMBER: Each of product containers of label with you will call to a person, control center or company for assistance. You may also call at 1-800-424-9300 for emergency medical treatment information.

NOTES: PHYSICIAN: Contains paraquat and other toxic and very toxic pesticides. Do not use if large eye treatment, suggest to contact a doctor or ophthalmologist.

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Open dumping is prohibited. Do not store this product near lawns, shrubs, vegetables, or impasses by not done near heat or open flame. Release all empty containers by thoroughly igniting, steam can. Also, any spill while in the dry absorbent and dispose of waste as indicated under "Pesticide Disposal." For safety and protection of unauthorized users, all pesticides should be stored in locked facilities to prevent access and areas of application should be stored in separate areas with enough distance between the pesticide to provide clear identification. Opened partially used pesticides should be stored in original labeled containers unless approved by the manufacturer. Containers necessary to take proper damage, carefully mark and family members of the new container.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or dilute is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the nearest state representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Non-refillable containers (2.5 gallons, 3 gallons and 55 gallons) do not use or pressure containers after they are empty. In all cases, high pressure containers (for example, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350, 1400, 1450, 1500, 1550, 1600, 1650, 1700, 1750, 1800, 1850, 1900, 1950, 2000, 2050, 2100, 2150, 2200, 2250, 2300, 2350, 2400, 2450, 2500, 2550, 2600, 2650, 2700, 2750, 2800, 2850, 2900, 2950, 3000, 3050, 3100, 3150, 3200, 3250, 3300, 3350, 3400, 3450, 3500, 3550, 3600, 3650, 3700, 3750, 3800, 3850, 3900, 3950, 4000, 4050, 4100, 4150, 4200, 4250, 4300, 4350, 4400, 4450, 4500, 4550, 4600, 4650, 4700, 4750, 4800, 4850, 4900, 4950, 5000, 5050, 5100, 5150, 5200, 5250, 5300, 5350, 5400, 5450, 5500, 5550, 5600, 5650, 5700, 5750, 5800, 5850, 5900, 5950, 6000, 6050, 6100, 6150, 6200, 6250, 6300, 6350, 6400, 6450, 6500, 6550, 6600, 6650, 6700, 6750, 6800, 6850, 6900, 6950, 7000, 7050, 7100, 7150, 7200, 7250, 7300, 7350, 7400, 7450, 7500, 7550, 7600, 7650, 7700, 7750, 7800, 7850, 7900, 7950, 8000, 8050, 8100, 8150, 8200, 8250, 8300, 8350, 8400, 8450, 8500, 8550, 8600, 8650, 8700, 8750, 8800, 8850, 8900, 8950, 9000, 9050, 9100, 9150, 9200, 9250, 9300, 9350, 9400, 9450, 9500, 9550, 9600, 9650, 9700, 9750, 9800, 9850, 9900, 9950, 10000, 10050, 10100, 10150, 10200, 10250, 10300, 10350, 10400, 10450, 10500, 10550, 10600, 10650, 10700, 10750, 10800, 10850, 10900, 10950, 11000, 11050, 11100, 11150, 11200, 11250, 11300, 11350, 11400, 11450, 11500, 11550, 11600, 11650, 11700, 11750, 11800, 11850, 11900, 11950, 12000, 12050, 12100, 12150, 12200, 12250, 12300, 12350, 12400, 12450, 12500, 12550, 12600, 12650, 12700, 12750, 12800, 12850, 12900, 12950, 13000, 13050, 13100, 13150, 13200, 13250, 13300, 13350, 13400, 13450, 13500, 13550, 13600, 13650, 13700, 13750, 13800, 13850, 13900, 13950, 14000, 14050, 14100, 14150, 14200, 14250, 14300, 14350, 14400, 14450, 14500, 14550, 14600, 14650, 14700, 14750, 14800, 14850, 14900, 14950, 15000, 15050, 15100, 15150, 15200, 15250, 15300, 15350, 15400, 15450, 15500, 15550, 15600, 15650, 15700, 15750, 15800, 15850, 15900, 15950, 16000, 16050, 16100, 16150, 16200, 16250, 16300, 16350, 16400, 16450, 16500, 16550, 16600, 16650, 16700, 16750, 16800, 16850, 16900, 16950, 17000, 17050, 17100, 17150, 17200, 17250, 17300, 17350, 17400, 17450, 17500, 17550, 17600, 17650, 17700, 17750, 17800, 17850, 17900, 17950, 18000, 18050, 18100, 18150, 18200, 18250, 18300, 18350, 18400, 18450, 18500, 18550, 18600, 18650, 18700, 18750, 18800, 18850, 18900, 18950, 19000, 19050, 19100, 19150, 19200, 19250, 19300, 19350, 19400, 19450, 19500, 19550, 19600, 19650, 19700, 19750, 19800, 19850, 19900, 19950, 20000, 20050, 20100, 20150, 20200, 20250, 20300, 20350, 20400, 20450, 20500, 20550, 20600, 20650, 20700, 20750, 20800, 20850, 20900, 20950, 21000, 21050, 21100, 21150, 21200, 21250, 21300, 21350, 21400, 21450, 21500, 21550, 21600, 21650, 21700, 21750, 21800, 21850, 21900, 21950, 22000, 22050, 22100, 22150, 22200, 22250, 22300, 22350, 22400, 22450, 22500, 22550, 22600, 22650, 22700, 22750, 22800, 22850, 22900, 22950, 23000, 23050, 23100, 23150, 23200, 23250, 23300, 23350, 23400, 23450, 23500, 23550, 23600, 23650, 23700, 23750, 23800, 23850, 23900, 23950, 24000, 24050, 24100, 24150, 24200, 24250, 24300, 24350, 24400, 24450, 24500, 24550, 24600, 24650, 24700, 24750, 24800, 24850, 24900, 24950, 25000, 25050, 25100, 25150, 25200, 25250, 25300, 25350, 25400, 25450, 25500, 25550, 25600, 25650, 25700, 25750, 25800, 25850, 25900, 25950, 26000, 26050, 26100, 26150, 26200, 26250, 26300, 26350, 26400, 26450, 26500, 26550, 26600, 26650, 26700, 26750, 26800, 26850, 26900, 26950, 27000, 27050, 27100, 27150, 27200, 27250, 27300, 27350, 27400, 27450, 27500, 27550, 27600, 27650, 27700, 27750, 27800, 27850, 27900, 27950, 28000, 28050, 28100, 28150, 28200, 28250, 28300, 28350, 28400, 28450, 28500, 28550, 28600, 28650, 28700, 28750, 28800, 28850, 28900, 28950, 29000, 29050, 29100, 29150, 29200, 29250, 29300, 29350, 29400, 29450, 29500, 29550, 29600, 29650, 29700, 29750, 29800, 29850, 29900, 29950, 30000, 30050, 30100, 30150, 30200, 30250, 30300, 30350, 30400, 30450, 30500, 30550, 30600, 30650, 30700, 30750, 30800, 30850, 30900, 30950, 31000, 31050, 31100, 31150, 31200, 31250, 31300, 31350, 31400, 31450, 31500, 31550, 31600, 31650, 31700, 31750, 31800, 31850, 31900, 31950, 32000, 32050, 32100, 32150, 32200, 32250, 32300, 32350, 32400, 32450, 32500, 32550, 32600, 32650, 32700, 32750, 32800, 32850, 32900, 32950, 33000, 33050, 33100, 33150, 33200, 33250, 33300, 33350, 33400, 33450, 33500, 33550, 33600, 33650, 33700, 33750, 33800, 33850, 33900, 33950, 34000, 34050, 34100, 34150, 34200, 34250, 34300, 34350, 34400, 34450, 34500, 34550, 34600, 34650, 34700, 34750, 34800, 34850, 34900, 34950, 35000, 35050, 35100, 35150, 35200, 35250, 35300, 35350, 35400, 35450, 35500, 35550, 35600, 35650, 35700, 35750, 35800, 35850, 35900, 35950, 36000, 36050, 36100, 36150, 36200, 36250, 36300, 36350, 36400, 36450, 36500, 36550, 36600, 36650, 36700, 36750, 36800, 36850, 36900, 36950, 37000, 37050, 37100, 37150, 37200, 37250, 37300, 37350, 37400, 37450, 37500, 37550, 37600, 37650, 37700, 37750, 37800, 37850, 37900, 37950, 38000, 38050, 38100, 38150, 38200, 38250, 38300, 38350, 38400, 38450, 38500, 38550, 38600, 38650, 38700, 38750, 38800, 38850, 38900, 38950, 39000, 39050, 39100, 39150, 39200, 39250, 39300, 39350, 39400, 39450, 39500, 39550, 39600, 39650, 39700, 39750, 39800, 39850, 39900, 39950, 40000, 40050, 40100, 40150, 40200, 40250, 40300, 40350, 40400, 40450, 40500, 40550, 40600, 40650, 40700, 40750, 40800, 40850, 40900, 40950, 41000, 41050, 41100, 41150, 41200, 41250, 41300, 41350, 41400, 41450, 41500, 41550, 41600, 41650, 41700, 41750, 41800, 41850, 41900, 41950, 42000, 42050, 42100, 42150, 42200, 42250, 42300, 42350, 42400, 42450, 42500, 42550, 42600, 42650, 42700, 42750, 42800, 42850, 42900, 42950, 43000, 43050, 43100, 43150, 43200, 43250, 43300, 43350, 43400, 43450, 43500, 43550, 43600, 43650, 43700, 43750, 43800, 43850, 43900, 43950, 44000, 44050, 44100, 44150, 44200, 44250, 44300, 44350, 44400, 44450, 44500, 44550, 44600, 44650, 44700, 44750, 44800, 44850, 44900, 44950, 45000, 45050, 45100, 45150, 45200, 45250, 45300, 45350, 45400, 45450, 45500, 45550, 45600, 45650, 45700, 45750, 45800, 45850, 45900, 45950, 46000, 46050, 46100, 46150, 46200, 46250, 46300, 46350, 46400, 46450, 46500, 46550, 46600, 46650, 46700, 46750, 46800, 46850, 46900, 46950, 47000, 47050, 47100, 47150, 47200, 47250, 47300, 47350, 47400, 47450, 47500, 47550, 47600, 47650, 47700, 47750, 47800, 47850, 47900, 47950, 48000, 48050, 48100, 48150, 48200, 48250, 48300, 48350, 48400, 48450, 48500, 48550, 48600, 48650, 48700, 48750, 48800, 48850, 48900, 48950, 49000, 49050, 49100, 49150, 49200, 49250, 49300, 49350, 49400, 49450, 49500, 49550, 49600, 49650, 49700, 49750, 49800, 49850, 49900, 49950, 50000, 50050, 50100, 50150, 50200, 50250, 50300, 50350, 50400, 50450, 50500, 50550, 50600, 50650, 50700, 50750, 50800, 50850, 50900, 50950, 51000, 51050, 51100, 51150, 51200, 51250, 51300, 51350, 51400, 51450, 51500, 51550, 51600, 51650, 51700, 51750, 51800, 51850, 51900, 51950, 52000, 52050, 52100, 52150, 52200, 52250, 52300, 52350, 52400, 52450, 52500, 52550, 52600, 52650, 52700, 52750, 52800, 52850, 52900, 52950, 53000, 53050, 53100, 53150, 53200, 53250, 53300, 53350, 53400, 53450, 53500, 53550, 53600, 53650, 53700, 53750, 53800, 53850, 53900, 53950, 54000, 54050, 54100, 54150, 54200, 54250, 54300, 54350, 54400, 54450, 54500, 54550, 54600, 54650, 54700, 54750, 54800, 54850, 54900, 54950, 55000, 55050, 55100, 55150, 55200, 55250, 55300, 55350, 55400, 55450, 55500, 55550, 55600, 55650, 55700, 55750, 55800, 55850, 55900, 55950, 56000, 56050, 56100, 56150, 56200, 56250, 56300, 56350, 56400, 56450, 56500, 56550, 56600, 56650, 56700, 56750, 56800, 56850, 56900, 56950, 57000, 57050, 57100, 57150, 57200, 57250, 57300, 57350, 57400, 57450, 57500, 57550, 57600, 57650, 57700, 57750, 57800, 57850, 57900, 57950, 58000, 58050, 58100, 58150, 58200, 58250, 58300, 58350, 58400, 58450, 58500, 58550, 58600, 58650, 58700, 58750, 58800, 58850, 58900, 58950, 59000, 59050, 59100, 59150, 59200, 59250, 59300, 59350, 59400, 59450, 59500, 59550, 59600, 59650, 59700, 59750, 59800, 59850, 59900, 59950, 60000, 60050, 60100, 60150, 60200, 60250, 60300, 60350, 60400, 60450, 60500, 60550, 60600, 60650, 60700, 60750, 60800, 60850, 60900, 60950, 61000, 61050, 61100, 61150, 61200, 61250, 61300, 61350, 61400, 61450, 61500, 61550, 61600, 61650, 61700, 61750, 61800, 61850, 61900, 61950, 62000, 62050, 62100, 62150, 62200, 62250, 62300, 62350, 62400, 62450, 62500, 62550, 62600, 62650, 62700, 62750, 62800, 62850, 62900, 62950, 63000, 63050, 63100, 63150, 63200, 63250, 63300, 63350, 63400, 63450, 63500, 63550, 63600, 63650, 63700, 63750, 63800, 63850, 63900, 63950, 64000, 64050, 64100, 64150, 64200, 64250, 64300, 64350, 64400, 64450, 64500, 64550, 64600, 64650, 64700, 64750, 64800, 64850, 64900, 64950, 65000, 65050, 65100, 65150, 65200, 65250, 65300, 65350, 65400, 65450, 65500, 65550, 65600, 65650, 65700, 65750, 65800, 65850, 65900, 65950, 66000, 66050, 66100, 66150, 66200, 66250, 66300, 66350, 66400, 66450, 66500, 66550, 66600, 66650, 66700, 66750, 66800, 66850, 66900, 66950, 67000, 67050, 67100, 67150, 67200, 67250, 67300, 67350, 67400, 67450, 67500, 67550, 67600, 67650, 67700, 67750, 67800, 67850, 67900, 67950, 68000, 68050, 68100, 68150, 68200, 68250, 68300, 68350, 68400, 68450, 68500, 68550, 68600, 68650, 68700, 68750, 68800, 68850, 68900, 68950, 69000, 69050, 69100, 69150, 69200, 69250, 69300, 69350, 69400, 69450, 69500, 69550, 69600, 69650, 69700, 69750, 69800, 69850, 69900, 69950, 70000, 70050, 70100, 70150, 70200, 70250, 70300, 70350, 70400, 70450, 70500, 70550, 70600, 70650, 70700, 70750, 70800, 70850, 70900, 70950, 71000, 71050, 71100, 71150, 71200, 71250, 71300, 71350, 71400, 71450, 71500, 71550, 71600, 71650, 71700, 71750, 71800, 71850, 71900, 71950, 72000, 72050, 72100, 72150, 72200, 72250, 72300, 72350, 72400, 72450, 72500, 72550, 72600, 72650, 72700, 72750, 72800, 72850, 72900, 72950, 73000, 73050, 73100, 73150, 73200, 73250, 73300, 73350, 73400, 73450, 73500, 73550, 73600, 73650, 73700, 73750, 73800, 73850, 73900, 73950, 74000, 74050, 74100, 74150, 74200, 74250, 74300, 74350, 74400, 74450, 74500, 74550, 74600, 74650, 74700, 74750, 74800, 74850, 74900, 74950, 75000, 75050, 75100, 75150, 75200, 75250, 75300, 75350, 75400, 75450, 75500, 75550, 75600, 75650, 75700, 75750, 75800, 75850, 75900, 75950, 76000, 76050, 76100, 76150, 76200, 76250, 76300, 76350, 76400, 76450, 76500, 76550, 76600, 76650, 76700, 76750, 76800, 76850, 76900, 76950, 77000, 77050, 77100, 77150, 77200, 77250, 77300, 77350, 77400, 77450, 77500, 77550, 77600, 77650, 77700, 77750, 77800, 77850, 77900, 77950, 78000, 78050, 78100, 78150, 78200, 78250, 78300, 78350, 78400, 78450, 78500, 78550, 78600, 78650, 78700, 78750, 78800, 78850, 78900, 78950, 79000, 79050, 79100, 79150, 79200, 79250, 79300, 79350, 79400, 79450, 79500, 79550, 79600, 79650, 79700, 79750, 79800, 79850, 79900, 79950, 80000, 80050, 80100, 80150, 80200, 80250, 80300, 80350, 80400, 80450, 80500, 80550, 80600, 80650, 80700, 80750, 80800, 80850, 80900, 80950, 81000, 81050, 81100, 81150, 81200, 81250, 81300, 81350, 81400, 81450, 81500, 81550, 81600, 81650, 81700, 81750, 81800, 81850, 81900, 81950, 82000, 82050, 82100, 82150, 82200, 82250, 82300, 82350, 82400, 82450, 82500, 82550, 82600, 82650, 82700, 82750, 82800, 82850, 82900, 82950, 83000, 83050, 83100, 83150, 83200, 83250, 83300, 83350, 83400, 83450, 83500, 83550, 83600, 83650, 83700, 83750, 83800, 83850, 83900, 83950, 84000, 84050, 84100, 84150, 84200, 84250, 84300, 84350, 84400, 8445

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

NOTE TO PHYSICIAN: Contains petroleum distillate. Vomiting may cause aspiration pneumonia. If lavage is performed, suggest endotracheal and/or esophagoscopy control.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are neoprene or nitrile rubber. If you want more options, follow the instructions for Category E on an EPA chemical-resistance category selection chart.

All mixers; loaders, applicators, flaggers and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves
- Chemical-resistant apron when applying with any hand-held nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

ENGINEERING CONTROLS


Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.



ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate. Apply this product only as directed on label.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et. Al. v. EP, C01-132C, (W.D. WA). For further information, please refer to <http://www.epa.gov/espp/wtc>.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.


PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (or pets) to enter the treated area until sprays have dried.



PRODUCT INFORMATION

The performance of Base Camp® LV 6 may be affected by local conditions, crop varieties, and application method. User should consult local extension service, agricultural experiment station or university weed specialists, and State regulatory agencies for recommendations in your area.

Best results are obtained when Base Camp LV 6 is applied to young succulent weeds that are actively growing. The lower specified rates will be satisfactory on susceptible annual weeds. For perennial weeds and conditions found in very dry areas of the western states, where control is difficult, the higher specified rates should be used.

When Base Camp LV 6 is used for weed control in crops, the growth stage of the crop must be considered.

Some plants and weeds, especially woody varieties, are difficult to control and may require repeat applications.

Apply 1 to 5 gallons of total spray by air or 5 to 25 gallons by ground equipment unless otherwise directed. In either case, use the same amount of 2,4-D per acre, as specified in the ensuing use rates. For crop uses, do not mix with oil, surfactants, or other adjuvants unless specifically recommended. Such combinations may reduce the herbicide's selectivity and could result in crop damage.

Many states have regulations concerning aerial application of 2,4-D formulations. Consult local regulatory authorities before making applications. Although this product is a low volatile formulation, temperatures above 90°F may release vapors that can result in damage to susceptible crops growing nearby.

Base Camp LV 6 should not be allowed to come into contact with desirable, susceptible plants such as beans, cotton, fruit trees, grapes, legumes, ornamentals, peas, tomatoes, and other vegetables. Base Camp LV 6 should not be used in greenhouses. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

If stored below freezing, efficacy is not affected if product is warmed to 40°F and agitated before using.

Spray equipment that is used to apply any 2,4-D products should not be used for any other purpose until thoroughly cleaned.

Spray Preparation:

Add the specified amount of Base Camp LV 6 to approximately one-half the volume of water to be used for spraying. Agitate well, then add the remainder of the water. Continue agitation during application until spray tank is empty.

TANK MIXING PRECAUTIONS

- Read carefully and follow all applicable use directions, precautions and limitations on the respective product labels.
- Do not exceed the specified label use rates. Do not tank mix with another pesticide product that contains the same active ingredient as this product unless the label of either tank mix partner specifies the maximum dosage that may be used.

Use in Liquid Nitrogen Fertilizer:

Base Camp LV 6 may be combined with liquid nitrogen fertilizer suitable for foliar application on corn, grass, pastures, or small grains in one operation. Use Base Camp LV 6 according to directions on this label for those crops. Use liquid nitrogen fertilizer at rates recommended by supplier or Extension Service Specialist. Mix the Base Camp LV 6 and fertilizer according to the following instructions:

Fill the spray tank approximately one-half full with the liquid nitrogen fertilizer. Add the Base Camp LV 6 while agitating the tank. Add the remainder of the fertilizer while continuing to agitate. Apply immediately, maintaining agitation during application until tank is empty. DO NOT APPLY DURING COLD (NEAR FREEZING) WEATHER. Spray mixture must be used immediately and must not be stored.

Note: If adequate and continuous agitation is not maintained, separation of the spray mixture and/or clogging of the nozzles is likely to occur.

SPRAYER CLEAN OUT

To avoid injury to desirable plants, equipment used to apply Base Camp LV 6 should be thoroughly cleaned before re-use or applying other chemicals.


- Rinse and flush application equipment thoroughly after use at least three times with water. Dispose of all rinse water by application to treatment area or apply to non-cropland area away from water supplies.
- During the second rinse, add 1 quart of household ammonia for every 25 gallons of water. Circulate the solution through entire system so that all internal surfaces are contacted (15-20 minutes). Let the solutions stand for several hours, preferably overnight.
- Flush the solution out of the spray tank through the boom.
- Rinse the system twice with clean water, recirculating and draining each time.
- Remove the nozzles and screens and clean separately.
- If equipment is to be used to apply another pesticide or agricultural chemical to a 2,4-D susceptible crop, additional steps may be required to remove all traces of 2,4-D including cleaning of disassembled parts and replacement of hoses or other fittings that may contain absorbed 2,4-D.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE standard 572) or a volume-mean diameter of 385 microns or greater for spinning atomizer nozzles.



When applying sprays that contain 2,4-D mixed with other active ingredients that require a medium or more fine spray, apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for non-target species, non-target crops) within 250 feet downwind. If applying a medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist or, b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial equipment and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

For aerial equipment, the boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or right-of-way applications.

When applications are made in a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

For ground boom application, do not apply with a nozzle height greater than 4 feet above the crop canopy.

Use low-pressure sprays to minimize drift. Where states have regulations that specify minimum spray volumes, they should be observed.

APPLICATION INSTRUCTIONS

Spray volume: Apply with calibrated air or ground equipment using sufficient spray volume to provide adequate coverage of target weeds or as otherwise directed in specific use directions. For broadcast application, apply the specified rate of Base Camp LV 6 in a spray volume of 2 or more gallons per acre by air and 10 or more gallons per acre for ground equipment. In general, spray volume should be increased as crop canopy height and weed density increase in order to obtain adequate spray coverage. Do not apply less than 2 gallons total spray volume per acre.

Use Rates: Generally, lower rates in specified rate ranges will be satisfactory for more sensitive weeds species, when weeds are small, and when environmental conditions are favorable for rapid growth. Use higher rates in the specified rate range for less sensitive species and under less favorable growing conditions. For crop uses, do not mix with emulsifiable oil or other adjuvants unless specifically recommended on this label. Deep-rooted perennial weeds such as Canada thistle, field bindweed and many woody plants usually require repeated applications for effective control.


Spot Treatments: To prevent misapplication, spot treatments should be applied with a calibrated boom or with hand sprayers using a fixed spray volume per 1,000 sq. ft. as indicated below.

Hand-Held Sprayers: Hand-held sprayers may be used for spot applications of Base Camp LV 6. Care should be taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on the application rate for an area of 1,000 sq. ft. Mix the amount of Base Camp LV 6 (fl. oz. or ml) corresponding to the desired broadcast rate in 1 to 3 gallons of spray. To calculate the amount of Base Camp LV 6 required for larger areas, multiply the table value (fl. oz. or ml) by the thousands of sq. ft. to be treated. An area of 1,000 sq. ft. is approximately 10.5 x 10.5 yards (strides) in size.

Rate Conversion Table for Spot Treatment:

Label Broadcast Rate (pt/acre)							
1/3	1/2	2/3	1	1-1/3	2	2-2/3	5-1/3
Equivalent Amount of Base Camp LV 6 per 1,000 sq. ft.							
1/8 fl. oz. ¹ (3.7 ml)	1/5 fl. oz. (5.9 ml)	1/4 fl. oz. (7.4 ml)	3/8 fl. oz. (11 ml)	1/2 fl. oz. (15 ml)	3/4 fl. oz. (22 ml)	1 fl. oz. (30 ml)	2 fl. oz. (60 ml)

¹Conversion factors:
1 pt. = 16 fl. oz.;
1 fl. oz. = 29.6 (30) ml



USE SITES

Base Camp LV 6 is used to control broadleaf weeds in cereal crops, corn, and sorghum; weeds and brush in rangeland, pastures, rights-of-way, and other listed non-crop uses.

WEEDS CONTROLLED

Annual or Biennial Weeds:

- beggartick⁽¹⁾
- bittercress, small-flowered⁽²⁾
- bitterweed
- broomweed, common⁽¹⁾
- burdock, common
- buttercup, small-flowered⁽¹⁾⁽²⁾
- carpetweed
- cinquefoil, common⁽²⁾
- cinquefoil, rough⁽²⁾
- cocklebur, common
- coffeeweed
- copperleaf, Virginia⁽²⁾
- croton, Texas
- croton, woolly
- fixweed
- galinsoga
- geranium, Carolina⁽²⁾
- hemp, wild
- horseweed (marestail)⁽²⁾
- jewelweed
- jimsonweed
- knotweed⁽¹⁾
- kochia
- lambquarter, common
- lettuce, prickly⁽¹⁾⁽²⁾
- lettuce, wild
- lupine
- mallow, little⁽¹⁾
- mallow, Venice⁽¹⁾
- marshelder
- morningglory, annual
- morningglory, ivy
- morningglory, woolly
- mousetail⁽²⁾
- mustards (except blue mustard)
- parsnip, wild
- pennycress (fanweed)
- pepperweed (*Lepidium spp.*)⁽¹⁾⁽²⁾
- pigweed (*Amaranthus spp.*)⁽¹⁾
- poorjoe
- primrose, common
- purslane, common⁽²⁾
- pursley, Florida
- radish, wild
- ragweed, common
- ragweed, giant
- rape, wild
- rocket, yellow
- salsify, common⁽¹⁾
- salsify, western⁽¹⁾
- shepherdspurse
- sicklepod
- smartweed (annual species)⁽¹⁾⁽²⁾
- sneezeweed, bitter
- sowthistle, annual
- sowthistle, spiny
- spanishneedle
- sunflower
- sweetclover
- tansymustard
- thistle, bull
- thistle, musk⁽¹⁾
- thistle, Russian (tumbleweed)⁽¹⁾
- velvetleaf
- vetch


WEEDS CONTROLLED (cont.)

Perennial Weeds

- alfalfa⁽¹⁾⁽²⁾
- artichoke, Jerusalem⁽¹⁾
- aster, many-flower⁽¹⁾
- Austrian fieldcress⁽¹⁾
- bindweed (hedge, field and European)⁽¹⁾⁽²⁾
- blue lettuce
- blueweed, Texas
- broomweed
- bullnettle⁽¹⁾⁽²⁾
- carrot, wild⁽¹⁾
- catnip
- chicory
- clover, red⁽¹⁾⁽²⁾
- coffeeweed
- cress, hoary⁽¹⁾
- dandelion
- dock⁽¹⁾
- dogbane⁽¹⁾
- eveningprimrose, cutleaf⁽²⁾
- garlic, wild
- goldenrod
- hawkweed, orange⁽¹⁾
- heal-all
- ironweed, western⁽²⁾
- ivy, ground⁽¹⁾
- nettles (including stinging)⁽¹⁾
- onion, wild⁽¹⁾
- pennywort
- plantain
- ragwort, tansy⁽¹⁾
- sowthistle, perennial
- thistle, Canada⁽¹⁾⁽²⁾
- vervain⁽¹⁾
- wormwood

^{(1)Difficult-to-Control Weeds:} These weeds are only partially controlled and may require repeat applications and/or use of the higher specified rate of Base Camp LV 6 even under ideal conditions of application.

^{(2)Base Camp LV 6 may not be used to control this weed species in the state of California.}



CROPS

CEREAL GRAINS (Wheat, Barley, Millet, Oats, Rye)

(Not Underseeded with Legumes)

CROP / APPLICATION TIMING	Base Camp LV.6 (PT / ACRE)	SPECIFIC USE DIRECTIONS
Wheat, Barley, Millet, Rye Annual and biennial Broadleaf weeds Perennial broadleaf weeds	1/3 to 1-1/3 ¹ 2/3 to 1-1/3 ¹	Apply after crop is fully tillered and before boot stage of growth (usually 4 to 8 inches tall) but not forming joints in the stem. Do not apply before tillering or from early boot through the milk stage of growth.
Oats (Spring Seeded) (Fall Seeded Southern)	1/3 1/2 to 1 ¹	Apply after crop is fully tillered and before boot stage of growth (usually 4 to 8 inches tall) and weeds are small. Do not apply before tillering or from early boot through the milk stage of growth. Do not apply during or immediately following cold weather.
Preharvest application (all cereals)	2/3	Apply using air or ground equipment to control weeds that could interfere with harvest, or to suppress perennial weeds. Apply when grain is in dough stage. Do not apply from early boot through the milk stage of growth.

¹Use the lower rate in the rate range if small annual or biennial weeds are the major problem. Use the higher rate if perennial weeds or annual or biennial weeds are present which are considered to be hard-to-kill as determined by local experience. Higher rates increase the risk of crop injury and should be used only where weed control justifies such risk. Do not apply Base Camp LV 6 at the crop seedling stage of growth. Consult state agricultural experiment station or extension service weed specialists for recommendations or suggestions to fit local conditions.

CEREAL GRAIN RESTRICTIONS:

Postemergence:


- Do not make more than one application per crop cycle.
- Do not apply more than 29.0 fluid ounces per acre per application.

Preharvest:

- Do not make more than one application per crop cycle.
- Do not apply more than 11.6 fluid ounces per acre per application.

Preharvest interval is 14 days.

Base Camp LV 6 contains 0.7 pounds a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 1.75 pounds of a.e. per acre per year.



12/20

ORNAMENTAL TURF

(Excluding Grasses Grown for Seed of Sod Farms)

(Includes lawns, golf courses, cemeteries and parks, airfields, roadsides, and vacant lots)

Use Requirements for Ornamental Turf Areas: When this product is applied to ornamental turf areas, follow PPE and reentry instructions in the "Non-agricultural Use Requirements" section of this label.


TREATMENT SITE (APPLICATION/TIMING)	Base Camp LV 6 (PT / ACRE)	SPECIFIC USE DIRECTIONS
Ornamental turf (postemergence)		Apply when weeds are small and actively growing. For best results, apply when soil moisture is adequate for active weed growth. Deep-rooted perennial weeds such as bindweed and Canada thistle may require repeat applications. Do not apply to newly seeded grasses until well established (five-leaf stage or later) and then use a maximum of 2/3 pints per acre. Cool season grasses are tolerant of higher rates.
Seeding grass (five-leaf stage or later)	1/2 to 2/3	
Well-established grasses	1-1/3 to 2	
Biennial and perennial broadleaf weeds	2	

PRECAUTIONS:

- Do not use on creeping grasses such as bentgrass except for spot treatment.
- Do not use on susceptible southern grasses such as St. Augustine.
- Do not use on dichondra or other herbaceous ground covers; legumes may be damaged or killed.
- Reseeding: Delay reseeding at least 30 days following application. Preferably, for spring applications reseed in the fall and for fall applications, reseed in the spring.

ORNAMENTAL TURFGRASS RESTRICTIONS:

- Do not apply more than 34.8 fluid ounces per acre per application.
- Do not make more than 2 applications per year.
- Minimum spray interval between broadcast applications is 30 days.
- If grazing or haying is anticipated, do not apply more than 2 pints per acre per application. Do not harvest grass for hay from treated areas for 7 days after application.



13/20

GRASSES GROWN FOR SEED OR SOD FARMS

Agricultural Use Requirements: When used in grass grown for seed or sod farms, follow PPE and reentry instructions in the "Agricultural Use Requirement" section of this label.

TREATMENT SITE (APPLICATION TIMING)	Base Camp LV 6 (PT / ACRE)	SPECIFIC USE DIRECTIONS
Grasses grown for seed (Postemergence Use)		Apply when weeds are small and actively growing. For best results, apply when soil moisture is adequate for active weed growth. Do not apply to newly seeded grasses until well established (five-leaf stage or later) and then use a maximum of 2/3 pints per acre. Cool season grasses are tolerant of higher rates. Do not apply to grass in the early boot through milk stage if seed production is desired. When grass is well established, higher rates of up to 2-2/3 pints per acre may be applied for control of hard-to-kill annual or perennial weeds. Deep-rooted perennials such as bindweed and Canada thistle may require repeat applications. Avoid mowing sod farms for 1 to 2 days before or after application. Delay irrigation until the day following application.
Seedling grass (five-leaf stage or later)	1/2 to 2/3	
Well-established grasses	2/3 to 2-2/3	
Sod Farms (Postemergence)	1/3 to 2-2/3	


PRECAUTIONS:

- Do not use on creeping grasses such as bentgrass except for spot treatment;
- Do not use on susceptible southern grasses such as St. Augustine;
- Do not use on dichondra or other herbaceous ground covers; legumes may be damaged or killed;
- Reseeding: Delay reseeding at least 30 days following application. Preferably, with spring application reseed in the fall and with fall applications, reseed in the spring.

GRASSES GROWN FOR SEED OR SOD RESTRICTIONS:

- Do not make more than 2 applications per year.
- Do not apply more than 46.4 fluid ounces per acre per application.
- Minimum of 21 days between applications.

Base Camp LV 6 contains 0.7 pounds a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of a.e. per acre per year (excluding spot treatments).



FALLOW LAND AND CROP STUBBLE

Fallow land is idle land, postharvest to crops or between crops.


TYPE OF WEEDS	Base Camp LV 6 (PT / ACRE)	SPECIFIC USE DIRECTIONS
Annual broadleaf weeds	2/3 to 1-1/3	Use a lower rate in the rate range when weeds are small (2 to 3 inches tall) and actively growing. Use a higher rate in the rate range when weeds are larger and under less favorable growth conditions.
Biennial broadleaf weeds	1-1/3 to 2-2/3	Apply when musk thistles or other biennial species are in the seedling to rosette stage and before development of flower stalks. The lower rate can be used in the spring during the rosette stage. Use the highest rate in the fall or after flower stalks have developed.
Perennial broadleaf weeds	1-1/3 to 2-2/3	Apply when perennial weeds are in bud to early bloom stage or while in good vegetative growth.
Wild garlic and onion in crop stubble	2-2/3	Apply to new regrowth of wild garlic or onion that occurs in the fall after harvest of other crops.

PRECAUTIONS: For best weed control results, do not cultivate for at least 2 weeks after application or until top growth is dead.

FALLOW-LAND RESTRICTIONS:

- Do not make more than two applications per year.
- Do not apply more than 2-2/3 pints per acre per application.
- Minimum spray interval between applications is 30 days.
- Plant only labeled crops within 30 days following last application.
- If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.

Base Camp LV 6 contains 0.7 pounds a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of a.e. per acre per year.



16/20

RANGELAND, ESTABLISHED GRASS PASTURES (cont.)

PRECAUTIONS:

- Do not use on bentgrass, alfalfa, clover, or other legumes.
- Do not use on newly seeded areas until grass is well established.
- Do not use from early boot to milk stage where grass seed production is desired.

RANGELAND & PASTURE RESTRICTIONS:

Livestock Feeding Restrictions:

- Do not graze dairy animals on treated areas within 7 days after application.
- Do not graze meat animals on treated areas within 3 days before slaughter.
- Do not cut forage for hay within 7 days of application.
- For government program grasslands, follow program grazing restrictions if more restrictive than those given above.

When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of a.e. per acre per year.

For susceptible annual and biennial broadleaf weeds do not exceed 23.2 fluid ounces per acre per application. For moderately susceptible biennial and perennial broadleaf weeds and woody plants, do not exceed 46.4 fluid ounces per acre per application. For spot treatment do not exceed 46.4 fluid ounces per acre. Do not make more than 2 applications per year. Minimum spray interval between applications is 30 days. If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard apply.

Base Camp LV 6 contains 0.7 pounds a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of a.e. per acre per year.


POTATOES

Fresh Market Only

APPLICATION TIMING / STAGE OF GROWTH	Base Camp LV 6 (PT./ACRE)	SPECIFIC USE DIRECTIONS
Postemergence	1/10 pt (1.6 fl. oz.) / acre	Make first application when potatoes are in the pre-bud stage (about 7 to 10 inches high) and make a second application about 10 to 14 days later.

POTATO RESTRICTIONS:


- Preharvest Interval: Do not harvest within 45 days of application.
- Do not apply more than 0.07 lbs a.e. (1/10 pint per acre) of Base Camp LV 6 per application.
- Do not exceed two applications per crop.
- A minimum of 10 days between applications is required.
- Do not apply more than 0.14 lbs a.e. (1/5 pint per acre) of Base Camp LV 6 per growing season.



FORESTRY USES

Forest site preparation, forest roadsides, brush control, established conifer release (including Christmas trees and reforestation areas)

TREATMENT SITE METHOD OF APPLICATION	Base Camp LV 6 (PT / ACRE)	SPECIFIC USE DIRECTIONS
Annual weeds Biennial and perennial broadleaf weeds and susceptible woody plants	1-1/3 to 2-2/3 2-2/3 to 5-2/3	Apply when weeds are small and growing actively before the bud stage. Apply when biennial and perennial species are in the seedling to rosette stage and before flower stalks appear. For difficult to control perennial broadleaf weeds and woody species, use up to 2-2/3 quarts of Base Camp LV6 and 1 to 4 quarts of Garlon 3A herbicide per acre. For conifer release, make application in early spring before budbreak of conifers when weeds are small and actively growing.
Spot Treatment to control broadleaf weeds	See instructions for "Spot Treatment"	Note: To control broadleaf weeds in small areas with a hand sprayer, apply a use rate equivalent to the specified broadcast rate and spray to thoroughly wet all foliage. See rate conversion table and instructions for "Spot Treatment" and use of hand-held sprayers under "Application".
Conifer Release: Species such as white pine, ponderosa pine, jack pine, red pine, black spruce, white spruce, red spruce, and balsam fir	2 to 4	To control competing hardwood species such as alder, aspen, birch, hazel, and willow, apply from mid to late summer when growth of conifer trees has hardened off and woody plants are still actively growing. Apply with ground or air equipment, using sufficient spray volume to ensure complete coverage. Because this treatment may cause occasional conifer injury, do not apply if such injury cannot be tolerated.
Directed Spray: Conifer plantations including pine	5-2/3 pints per 100 gal.	Apply when brush or weeds are actively growing by directing the spray so as to avoid contact with conifer foliage and injurious amounts of spray. Apply in oil, oil-water, or water carrier in a spray volume of 10 to 100 gallons per acre.
Basal Spray (May also be used in rangeland, pastures, and non-cropland)		Thoroughly wet the base and root collar of all stems until the spray begins to accumulate around the root collar at the ground line. Wetting stems with the mixture may also aid in control.
Surface of Cut Stumps (May also be used in rangeland, pastures, and non-cropland)	11-1/3 pints per 100 gal. 1.75 fl. oz./gal. of water.	Apply as soon as possible after cutting trees. Thoroughly soak the entire stump with the 2,4-D mixture including cut surface, bark and exposed roots.
Frill and Girdle (May also be used in rangeland, pastures, and non-cropland)		Cut frills (overlapping V-shaped notch cut downward through the bark in a continuous ring around the base of the tree) using an axe or other suitable tool. Saturate the freshly cut frills with the 2,4-D mixture.
Tree Injection Application (May also be used in rangeland, pastures, and non-cropland)	(1 to 1.4 ml per injection site)	To control and prevent resprouting of unwanted hardwood trees such as elm, hickory, oak, and sweetgum in forests and other non-crop areas, apply by injecting at a rate of 1 ml of undiluted Base Camp LV 6 per inch of trunk diameter as measured at breast height (DBH), approximately 4.5 ft. above the ground. Injection sites, however, should be as close to the root collar as possible, and the injection bit must penetrate the inner bark. Applications may be made throughout the year, but for best results apply between May 15 and October 15. Maples should not be treated during the spring sap flow. For hard to control species such as ash, maple, and dogwood, use 1.4 ml of undiluted Base-Camp LV 6 per injection site or double the number of 0.7 ml injections. Note: No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.



Conditions of Sale and Limitation of Warranty and Liability

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS," AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES, OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

WILBUR ELLIS Logo, Ideas to Grow With and Base Camp are registered trademarks, and Active's Tri-Droplet Logo is a trademark of Wilbur-Ellis Company. Garlon is a registered trademark of Dow AgroSciences.

F-100214 • K-100314

In Case of Emergency,
Call CHEMTREC: (800) 424-9300

Manufactured for
WILBUR ELLIS COMPANY
PO BOX 16458 • FRESNO, CA 93755

