FEOR THE CONTROL OF BOLL WEEVIL, THRIPS, CUTWORMS, COTTON FLEAHOPPER, RAPID AND TARNISH EPLANT BUGS, COTTON LEAFWORM, FALL ARMYWORM AND GARDEN WEBWORM ON COTTON AND CUT-WORMS: ARMYWORMS, GRASSHOPPERS AND FALL ARMYWORM IN SMALL GRAINS.

lon t

Ke

Hi

frd

lal

D

MARNING KEEP OUT OF REACH OF

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Campher containing 67% to 69% Chlorine)

Petroleum Distillates

INERT INGREDIENTS

TOTAL

Contains 6 lbs. Technical Toxaphene per gallo

DIRECTIONS FOR USE

COTTON INSECT CONTROL — EARLY SEASON: thrips apply 1 gallon RED PANTHER TOXAPHEN water to cover 7 acres. For control of cutworm TOXAPHENE in sufficient water to cover 1^{1}_{2} - 3

MID AND LATE SEASON — For control of boll fleahopper, rapid and tarnish plant bug, cotton armyworm and garden webworm apply 1 gallon To sufficient water to cover 112-3 acres. Repeat at 4until infestation is brought under control.

For best control with ground equipment apply amount of insecticide 3-6 gallons of water per acre.

For airplane application apply recommended amount cide in 3 to 10 gallons of water per acre.

Do not feed gin waste from cotton treated with Toxaphene to stock.

Do not graze livestock in cotton fields treated late in the season with Toxaphene.

Do not apply Toxaphene in excess of 4 lbs. per acre per applica-

E BELOW FOR ADDITIONAL WARNINGS.

AIN INSECT CONTROL — Wheat, Barley, Oais, and trol of cutworms apply 1 gallon Toxaphene in sufto cover 3 acres. For armyworms and grasshoppers to cover 4 acres. For fall armyworms apply 1 gal-4 acres. Do not apply more than once after heads form. Do not apply within one week of harvest on have oats. Do not apply within two weeks of harvest on barley. Do not feed or graze forage from grains treated with Toxaphene to dairy animals or animals being finished for slaugh-

> May be fatal if swallowed. Do not breathe spray mist. n skin, in eyes, or on clothing. Wash thoroughly. FROM CHILDREN AND DOMESTIC ANIMALS. nation of feed and foodstuffs. Do not use on dairy airy barns.

body of water.

fish and wildlife. Birds feeding on treated areas Do not apply when weather conditions favor drift ted. Apply this product only as specified on this

E EMPTY CONTAINERS. Destroy by perforating burying in a safe place.

s toxic to bees and should not be applied when bees fisiting the area.

SE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN

NON-WARRANTY: Coahoma Chemical Company, Inc., makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

FOR THE CONTROL OF BOLL WEEVIL, THRIPS, CUTWORMS, COTTON FLEAHOPPER, RAPID AND TARNISH PLANT BUGS, COTTON LEAFWORM, FALL ARMYWORM AND GARDEN WEBWORM ON COTTON AND CUTWORMS, ARMYWORMS, GRASSHOPPERS AND FALL ARMYWORM IN SMALL GRAINS.

KEEP OUT OF REACH OF CHILDREN. SEE BELOW FOR ADDITIONAL WARNINGS.

ACTIVE INGREDIENTS:	SMALL GRAIN TO THE RIVER OF THE
INERT INGREDIENTS	ficient water to a apply 1 gallon to to cover 3-4 are begun to form the second particular to the
DIRECTIONS FOR USE	at 70° F. WARNING. M. Do not get on the contract of the contr
CONTRACTOR CONTRACTOR EN VISEASON: I talporation de la REPORTATION DE LA ALHENE water tour accompany de la Bourse de la Contractor ap TONARE ENFORCEMENT DE LA CONTRACTOR DE LA CONTRACTOR DE	Avoid contaminate animals or in dame to the sufficient of any second contaminate animals or in dame to the sufficient of any second contaminate animals or in dame to the sufficient of any second contaminate animals or in dame to the sufficient of any second contaminate animals or in dame to the sufficient of any second contaminate animals or in dame to the sufficient of any second contaminate animals or in dame to the sufficient of any second contaminate animals or in dame to the sufficient of animals or in dame to the s
le de la companya de	reevil, cotton by be killed. In the safety of the safety o
	of insecti-
	DO NOT US: FLAME.

.

• •