FOR POST-EMERGENCE WEED CONTROL IN COTTON.

For control of weeds and grasses, including Johnson grass, nut grass, water free control of weeds and cocklebur.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT!

ACTIVE INGREDIENT:

MIXING DIRECTIONS

For proper mixing of RED PANTHER MMA fill the spray tank half full of water, then add the required amount of MMA. Start the pump circulating the spray mixture and finish filling the tank with water. MMA is completely soluble in water.

DIRECTIONS FOR USE

COTTON: Broadcast Application: Mix 3 pints RED PANTHER MMA plus 2-4 pints of RED PANTHER SURFACTANT IN 50 gallons of water. Apply 40 gallons of this mixture per acre. Thorough coverage of weeds and grass is absolutely necessary for maximum control. Adjust nozzles to direct the spray in a manner to allow maximum coverage of weeds. Keep spray off cotton foliage. Apply only when cotton is 3 inches high to first bloom.

Band Treatment: Mix 3 pints RED PANTHER MMA plus 2-4 pints of RED PANTHER SURFACTANT in 50 gallons of water. Apply at the rate of 1 gallon of spray solution per each 1 inch of band width to be treated on cotton grown in 40 inch rows. (Example: For a 12-inch band apply 12 gallons per acre.) A second application may be required during humid growing conditions. Time the second application, if necessary, about 1 to 3 weeks after the first application. Adjust nozzles to direct the spray in a manner to allow maximum coverage of weeds. Keep spray off cotton foliage. Apply only when cotton is 3 inches high to first bloom.

A Tys use RED PANTHER SURFACTANT with RED PANTHER MMA as it is not nearly as effective when used alone.

GENERAL GRASS AND WEED CONTROL: RED PANTHER MMA is useful for controlling Johnson grass, nut grass, water grass, and goose grass on ditch banks, rights-of-way, storage yards, and other non-crop areas. Mix 3 pints RED PANTHER MMA plus 2 to 4 PINTS of RED PANTHER SURFACTANT in 50 gallons of water. Spray the unwanted vegetation thoroughly, to the point of run-off. Make certain that complete coverage is obtained. Power sprayers are preferred, but any spray equipment which will give good coverage may be used. If regrowth occurs in spots, retreat as required.

APPLICATION METHODS

RED PANTHER MMA should be applied with a low volume sprayer, having satisfactory pumping and bypass action. Nozzles should be placed so a; to avoid spraying the cotton, but also to give thorough coverage of the weeds and grasses. Adjust nozzles to direct the spray in a manner to allow miximum coverage of weeds. Keep spray off cotton foliage.

APPLY ONLY WHEN COTTON IS 3 INCHES HIGH TO FIRST BLOOM. DO NOT APPLY AFTER FIRST BLOOM.

DO NOT FEED TREATED FOLIAGE TO LIVESTOCK OR GRAZE TREATED AREAS.

Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally.

WARN ING

Manuful if available and injury may result. Do not store spray solution in tank for a prolonged period, and clean application equipment thoroughly after every use by flushing with water. Do not flush spray equipment near or close by desirable plants. DO NOT CONTAMINATE WATERS USED FOR DOMESTIC OR ANIHAL CONSUMPTION, WATER USED BY WILDLIFE OR AQUATIC LIVE, WATER USED FOR IRRIGATION PURPOSES, OR WHERE FISH ARE FOUND. DO NOT RE-USE CONTAINER. DESTROY EMPTY CONTAINER.

NON-WARRANTY: Coahoma Chemical Company, Inc., makes no warranty of any kind, express or implied, on the use of this material and assumes no responsibility ownatsoever for any personal injury, property damage, or other typeof loss resulting from the use or handling of this material. The user assumes all risks for use and handling whether in accordance with directions or not.

USDA REG. NO. 2974-94

Manufactured By COAHOMA CHEMICAL COMPANY, INC. Clarksdale, Mississippi

ACCEPTED

JUN 28 1965

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT POR FCONOMIC POISON REGISTERED UNDER NO. 2974-94