

KEEP OUT OF REACH OF CHILDREN. SEE ANTIDOTE STATEMENT AND OTHER  
WARNING STATEMENTS BELOW

<p>Aromatic Petroleum Distillate</p> <p><b>INERT INGREDIENTS</b></p> <p>*EPN - A DuPont P Methyl Parathion contains 3 lbs. Methyl</p> <p><b>IN RED PA</b> e spray pu arance of se until a unitron</p> <p>or control of thrip P.E.C. per acre, 4-5 day in</p> <p>or control of me species plication w eded.</p> <p>Mississippi, Louisiana idworm, apply 1-1.3 p eded.</p> <p>ground equipment is llons of water per act or airplane application</p>	<p>Distillate</p> <p>uct</p> <p>athion and 2 lbs. EPN per allon.</p> <p><b>MIXING DIRECTIONS</b> MP with water or by cucuk into the tank. A proper m sion with no free oil on th ion is formed.</p> <p><b>DIRECTIONS FOR USE</b> s bugs, and cotton fleas rst application eded.</p> <p>cotton aph ites, apply on warrant</p> <p>nd Arkan- s of EPN-A</p> <p>sed apply</p> <p>only recom</p>	<p>50% 50% 50% 00%</p> <p><b>TOTAL</b></p> <p>ng the mixture th ough ture will have the ap- surface. Agitate b fore</p> <p>apply 2-3 pint of EPN- on warrants and repeat</p> <p>, garden webworm and per acre. Make first 4-5 day intervals as</p> <p>pollworm and tobacco t at 5 day intervals as</p> <p>int of EPN-A P in 3-6</p> <p>EPN-A P gallons</p>
--	---	--

**TREATMENT:** Atropine is a specific therapeutic antagonist of choice for cholinergic nervous system poisoning. If there are signs of parasympathetic overactivity, atropine sulfate should be injected in 1-2 mg. doses at 10-minute intervals, in a total dose of 10-20 mg. until complete atropinization has occurred. Morphine should be administered if necessary for postural drainage. Observe patient continuously for 24 hours. Repeat exposure to cholinesterase inhibitors may occur without warning. Allow no further exposure until cholinesterase regeneration has been allowed as determined by blood test.

**PRECAUTIONS IN USING**

**KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.** Do not contaminate food or feed products. Keep away from heat and open flame.

Do not allow or allow to drift to areas occupied by unprotected persons or animals. Keep all unprotected persons and animals out of treated areas until after drifting insecticide and volatile residues have dissipated. Do not allow drift to adjacent crops or livestock. Do not allow meat animals to feed on plants or crops during blood-feeding period. Do not apply to areas treated with insecticides active against fish and wildlife. Keep out of eyes, steam, and water.

DO NOT USE IN AREAS WHERE  
WINDS BLOW FROM TREATED AREAS  
TOWARD UNPROTECTED AREAS  
OR WHERE WINDS BLOW FROM  
TREATED AREAS TOWARD UNPROTECTED AREAS

# EPN-MP EMULSIFIABLE CONCENTRATE

For control of boll weevil, bollworm, aphids, cotton fleahopper, cotton leafworm, thrips, some species of spider mites, garden webworm, lygus bugs on cotton.

**DANGER - KEEP OUT OF REACH OF CHILDREN. SEE ANTIDOTE STATEMENT AND OTHER WARNING STATEMENTS BELOW.**

**ACTIVE INGREDIENTS:**

- \*O-Ethyl O-p-nitrophenyl phenylphosphonothioate 31.50%
- \*\*O,O-dimethyl O-p-nitrophenyl Thiophosphate 31.50%

**EPN-MP**

**DIRECTIONS:**

For control of boll weevil, bollworm, aphids, cotton fleahopper, cotton leafworm, thrips, some species of spider mites, garden webworm, lygus bugs on cotton.

Apply 1 pint of EPN-MP per acre. Repeat applications as needed.

**SYMPTOMS:** Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

**TREATMENT:** Administer 1-2 mg of atropine subcutaneously for every 100 mg of symptoms. Repeat every 30-60 minutes until the patient is breathing and has a pulse. Do not give if patient is comatose.

**PRECAUTIONS:** Do not apply to areas where humans or animals are present. Do not apply to areas where bees are present. Do not apply to areas where birds are present. Do not apply to areas where fish are present.

**POISON**

**ANTIDOTE AND FIRST AID TREATMENT—ATROPINE IS THE EMERGENCY ANTIDOTE FOR EPN OR METHYL PARATHION POISONING.** Consult your physician about obtaining an adequate supply of 1:1000 grain atropine tablets for emergency use. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING. **INTERNAL:** If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 table-spoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. **EXTERNAL:** If the material has been spilled on the skin, immediately remove patient from vicinity of the EPN-MP, remove all contaminated clothing and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least 10 minutes. If WARNING SYMPTOMS APPEAR Administer two 1:1000 grain atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (See WARNING SYMPTOMS below) Keep patient warm and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician. **NOTE TO PHYSICIAN: WARNING**

**DANGER**

**POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO EPN OR METHYL PARATHION POISONING.**

Do not swallow or get in eyes, on skin or on clothing. Do not breathe vapors or spray mist. Wear clean natural rubber gloves, clean, waterproof clothing, and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for EPN or Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If material gets into the eyes, immediately flush with water for at least ten minutes and get medical attention. In case of contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before reuse. In case of spillage, wash down with large amounts of water. **DO NOT RE-USE CONTAINER. DESTROY EMPTY CONTAINER.** Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

**NON-WARRANTY:** Cohnoma Chemical Company, Inc. make no warranty, express or implied, concerning this product or its use which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.