

CAUTION: KEEP OUT OF REACH OF CHILDREN. Protect pets and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for warfarin.

Baits should be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes.

CAROLINA'S RAT BAIT

SUGGESTED USE. Place 2 to 4 tablespoonsful of contents of this package in locations frequented by rats and mice and protect from children, dogs, cats and livestock by means of bait boxes or cages where necessary. CAROLINA'S RAT BAIT should be replenished as consumed. Use no decomposed, mouldy or sour bait.

Where a continuous source of infestation prevails from nearby dumps or fields, permanent bait stations should be used and CAROLINA'S RAT BAIT replenished as needed.

This package may be used as an individual bait station. Punch out the perforated space on back of box and nail box to the floor or on a board.

DIRECTIONS FOR USE

RATS — Place 2 to 6 ounces of bait in dry containers near to burrows and runways, or where rats are known to feed. As soon as infestation subsides, bait for a period of not less than 2 days and continue until all signs of feeding have ceased. This is a cumulative poison.

MICE — Place 1/2 to 1 tablespoonful amounts of bait in shallow containers where mice or rats are known to have been seen. Bait stations should be spaced 8 to 12 feet apart. Bait for an uninterrupted supply of bait for a period of not less than 15 days.

IMPORTANT NOTICE

BAITING SHOULD CONTINUE UNTIL COMPLETE LACK OF FEEDING IS NOTED.

CAUTION: KEEP OUT OF REACH OF CHILDREN. Protect pets and domestic animals from bait. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for warfarin.


Baits should be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes.

CAROLINA'S RAT BAIT

ONE POUND NET

CAROLINA'S RAT BAIT

ACTIVE INGREDIENTS
 Warfarin (3-(4-aminophenyl)-4-hydroxycoumarin) 0.025%
 N-(2-quinexalyl)phenanthridine (sulfarquinolone) 0.025%
NET WEIGHT 99.95%
NET CONTENTS 99.95%
 USDA Registration No. 27644



CAUTION: KEEP OUT OF REACH OF CHILDREN.
 See side panel for additional cautions.

CAROLINA GARDEN AND FARM SERVICE, INC., 279 E. MAIN AVE., GASTONIA, N.C.

CAROLINA'S RAT BAIT