

**BRAYTON**

**75-25**

# GRAIN FUMIGANT

## AN EFFECTIVE GRAIN FUMIGANT FOR CONTROLLING INSECT PESTS IN STORED GRAIN

### INSTRUCTIONS FOR TREATING GRAIN IN FARM GRANARIES

1. Weigh the grain to be treated and determine the amount of fumigant to use. The amount of fumigant to use is based on the weight of the grain to be treated. The amount of fumigant to use is based on the weight of the grain to be treated.

2. The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary.

3. The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary.

4. The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary.

5. The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary.

### ACTIVE INGREDIENTS

Phosphine	75.2
Diethyl phosphine	24.8
<b>TOTAL</b>	<b>100.0</b>

EPA REGISTRATION NO. 2593

**DANGER!**

KEEP OUT OF REACH OF CHILDREN

KEEP AWAY FROM HEAT OR OPEN FLAMES

**POISON**

VOLATILE LIQUID - VAPOR HARMFUL

Phosphine is a colorless, odorless, non-flammable gas. It is highly toxic to humans and animals. It is also highly flammable and can form explosive mixtures with air.

Diethyl phosphine is a colorless, odorless, non-flammable gas. It is highly toxic to humans and animals. It is also highly flammable and can form explosive mixtures with air.

The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary.

The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary. The fumigant should be applied to the grain in a granary.

APPLICATION: 1) Any method of application should have as its purpose even distribution of the liquid grain fumigant over the surface of the grain. 2) Avoid breathing fumigant vapors; apply fumigant from outside. 3) A power sprayer, or pump or strup pump may be used to apply liquid fumigant. Nozzle should be removed from spray rod. 4) The fumigant should be applied as stream and not atomized. This can be done by prying through a length of 1/2 inch pipe with end fitted and mashed together to form a fan shape. 5) After fumigant has been applied the bin should be closed as tightly as possible and the grain left undisturbed for at least 72 hours.

### INSTRUCTIONS FOR TREATING GRAIN IN FARM GRAIN ELEVATORS

1. The fumigant should be applied to the grain in a grain elevator. The fumigant should be applied to the grain in a grain elevator. The fumigant should be applied to the grain in a grain elevator.

2. The fumigant should be applied to the grain in a grain elevator. The fumigant should be applied to the grain in a grain elevator. The fumigant should be applied to the grain in a grain elevator.

3. The fumigant should be applied to the grain in a grain elevator. The fumigant should be applied to the grain in a grain elevator. The fumigant should be applied to the grain in a grain elevator.

*Brayton*

EB-5

# GRAIN FUMIGANT

**An Effective Fumigant For  
Controlling Insect Pests In Stored Grain**

(Grain and Rice Weevils, Flour Beetles, and Grain Moths)

**ACTIVE INGREDIENTS**

Ethylene Dibromide	7.2%
Ethylene Dichloride	29.2%
Carbon Tetrachloride	63.6%
TOTAL	100.0%

\*U.S. Pat. 2,391,890

USDA REGISTRATION No. 2993-23

**POISON** ☹ **DANGER**

**KEEP OUT OF REACH OF CHILDREN**

Hazardous Vapor and Liquid — May Be Fatal If Inhaled or Swallowed



**Brayton Chemicals, Inc., Burlington, Iowa**

DIRECTIONS

1. Shake well before use.

2. Use as directed.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.  
STORE IN TIGHTLY CLOSED CONTAINER, IN A COOL  
LOCATION AWAY FROM DWELLINGS