

**Brayton**

# MILL and WAREHOUSE SPRAY

## For Insect Control in Mills, Warehouses, and Food Processing Plants.

**ACTIVE INGREDIENTS.**

N-Octyl bicycloheptene dicarboximide .....	666%
Technical piperonyl butoxide* .....	.400%
Pyrethrins .....	.200%
Petroleum Solvent .....	98.734%
	<hr/>
	100.000%

\*Equivalent to .32% (Butylcarbityl) (6-Propylpiperonyl) ether and .08% related compounds.

USDA REG. No. 2993-27

### - WARNING -

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not spray towards or near an open flame. Do not pour, spill or store near fire or flame. Direct application on food products can cause contamination.

NOTE: The seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

FLUID CONTENTS \_\_\_\_\_ GALLONS

# BRAYTON CHEMICALS, Inc.

BURLINGTON, IOWA

### DIRECTIONS FOR USE

For control of accessible stages of weevils, flour beetles, saw toothed grain beetles, cadelle, Indian meal moth, adult and larvae, crickets, silverfish, ants, spiders, cheese and flour mites, flies, and flying moths.

As a contact spray - apply with a good sprayer using a large droplet flat spray nozzle. Use approximately one gallon of the insecticide to each 500 sq. ft. of surface.

**For space application** use in microsol generators or electric, compressed air, or steam atomizing devices. When fogging for control flies, flying moths, mosquitoes, gnats, wasps, and skippers, close all doors, windows, and openings during fogging operation and keep closed for 10 to 20 minutes after spraying. Repeat treatment as necessary.

Do not remain in treated areas and ventilate the area after treatment is completed.

### DIRECTIONS FOR USE

For control of crawling insects listed elsewhere, spray face of disjunctures, cracks, crevices, niches, under skirts, equipment, scales, water mounts, in conveyors, prooters, sifters, around the base of machinery and equipment, and all places where insects might harbor and multiply. Repeat treatment as necessary.

For control of ants, spray nests, along trailways, area around doorways, door sills, window frames, cracks in foundation walls, and other places where pests enter the premises.

**CAUTION** - When used in Food Processing Plants and Flour Mills spray only when plant is not in operation. Remove or cover food products. Do not spray directly on food processing equipment.

6/28/50  
2193-27