P-B FOG CONCENTRATE

FOR INSECT CONTROL IN FOOD PLANTS

ACTIVE INGREDIENTS

Pyrethrins		0.50%
Technical piperonyl butoxide*		0.40%
Petroleum solvent		99.10%
	TOTAL	100,00%

*Equivalent to .3200% of (butylcarbityl) (6-propyl piperonyl) ether and .0800% of related compounds.

USDA REG. No. 2993-43

-- WARNING -KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not spray near or toward open flame. Do not pour, spill or store near tire or flame. Avoid contamination of food and food processing equipment. Do not use inside Food Plants while Plants are in operation.

"IOTE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

FLUID CONTENTS

GALLONS



Visibile Areas Apply with compressed as soray as an ingging device and have a larger of the control of the cont

Saw H windows door and opening draining togging operation. Keep deviation of the philadeon Region.
 10/20 minutes after application. Region freatment when necessary.

Follow directions and recommendations of equipment manufacturer for proper setting and position of fogging machine. "Do not leuse empty container Wash thoroughly with water and détergent. Disca d'in a safe place."

9/2/70

2993-43

Directions For Use

For the control of accessible stages of weevils, flour beetles, meat moth larvae, cadelle, saw-toothed grain beetle, roaches, silverfish, ants, spiders, carpet beetles, cheese and flour mites, flies and flying moths

As A Surface Spray: for the control of crawling insects listed above, apply directly onto walls, partitions, floors, in cracks, crevices, under skids, equipment, scales, motor mounts, on ledges, in conveyors, proofers behind switch boxes, sinks, storage cabinets, floor drains, locker and lunch rooms and all places where insects may harbor and multiply

Clean areas thoroughly before spraying. Dust and dirt absorb insecticides which limits effectiveness. Repeat treatment when necessary