

7 1 6
445-6

Net Contents

BINGMAN'S

CIODRIN

LIVESTOCK SPRAY

Emulsifiable Concentrate

for Control of
FACE FLIES . STABLE FLIES
HORN FLIES . LICE
LONE STAR TICKS

on
Dairy Cattle, Beef Cattle, Sheep
Goats, or Hogs

ACTIVE INGREDIENTS:

Dimethyl phosphate of alpha-methylbenzyl 3-hydroxy-cis-crotonate--- 25.5%

Petroleum hydrocarbons----- 59.7%

INERT INGREDIENTS----- 14.8%

Total by weight----- 100.0%

W A R N I N G (12 pt. type)

KEEP OUT OF REACH OF CHILDREN (8 pt. type)

See back panel for additional cautions

BINGMAN LABORATORIES, INC.
CALDWELL, OHIO

* * * * *

DIRECTIONS FOR LIVESTOCK PEST CONTROL

For control of stable flies, horn flies, face flies, house flies, and lone star ticks on dairy cattle, beef cattle, sheep, goats, and hogs; and for control of house flies, face flies, and stable flies in and around livestock buildings. Do not use this spray directly on horses; horse barns must be vacated before spraying.

Mix 1 quart Livestock Spray Concentrate per 32 gallons water. For smaller quantities mix 1 ounce (2 tablespoons) per gallon of water. Spray thoroughly, covering all parts of animals. Use approximately 2 quarts spray for large animals, and proportionately less for smaller animals. Spraying premises such as inside and outside barn walls, eaves, fences, and other brooding or resting places will aid in reducing fly population in area. Repeat applications as necessary to maintain control, but not more than once weekly.

LOW VOLUME APPLICATION for controlling face flies, house flies, stable flies, and horn flies on dairy and beef cattle mix 2 oz. Livestock Spray Concentrate per 1 1/2 pnts water, or use 1 quart per 3 gallons water. Spray thoroughly covering all parts of the animal, using 1 to 2 fluid ounces of spray per animal. Apply daily with conventional hand or power sprayer.

Improved house fly and face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 8 gallons spray.

BACK RUBBER APPLICATION

BINGMAN LABORATORIES, INC.
CALDWELL, OHIO

* * * * *

DIRECTIONS FOR LIVESTOCK PEST CONTROL

For control of stable flies, horn flies, face flies, house flies, and lone star ticks on dairy cattle, beef cattle, sheep, goats, and hogs; and for control of house flies, face flies, and stable flies in and around livestock buildings. Do not use this spray directly on horses; horse barns must be vacated before spraying.

Mix 1 quart Livestock Spray Concentrate per 32 gallons water. For smaller quantities mix 1 ounce (2 tablespoons) per gallon of water. Spray thoroughly, covering all parts of animals. Use approximately 2 quarts spray for large animals, and proportionately less for smaller animals. Spraying premises such as inside and outside barn walls, eaves, fences, and other brooding or resting places will aid in reducing fly population in area. Repeat applications as necessary to maintain control, but not more than once weekly.

LOW VOLUME APPLICATION for controlling face flies, house flies, stable flies, and horn flies on dairy and beef cattle mix 2 oz. Livestock Spray Concentrate per $1\frac{1}{2}$ pints water, or use 1 quart per 3 gallons water. Spray thoroughly covering all parts of the animal, using 1 to 2 fluid ounces of spray per animal. Apply daily with conventional hand or power sprayer.

Improved house fly and face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 8 gallons spray.

BACK RUBBER APPLICATION

For use in back rubbers or face rubbers for control of horn flies and face flies mix 1 quart Livestock Spray Concentrate in 7 gallons back rubber base oil. This may be diesel fuel, or combination of diesel fuel and motor oil in accordance with local recommendations.

CAUTION: Do not contaminate feed or foodstuffs. To protect fish and wildlife do not contaminate streams, lakes, or ponds with this material. During commercial spraying or prolonged exposure in spraying and mixing operations, wear clean synthetic rubber gloves and a mask or respirator of a type found suitable by the U. S. Department of Agriculture for protection against Ciodrin insecticide. Use Livestock Spray Concentrate only for recommended purposes and at recommended dosages.

WARNING: HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. Wash thoroughly with soap and water after handling and before eating or smoking. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water. For eyes get medical attention. Keep away from heat and open flame.

U.S.D.A. Reg. No.