USE DIRECTIONS

FOR CLEANING

Lightly soiled areas—surfaces to be cleaned should be mopped or brushed with a solution containing 1 ounce of Edco—Syn per gallon of water. Rinse cleaned area with fresh water. Heavily soiled areas mop or brush surface with a solution containing 2 ounces of Edco—Syn per gallon of water.

GENERAL PURPOSE DISINFECTION

To disinfect walls, floors, and other hard surfaces in homes, schools, restaurants, rest rooms and other public areas, mop or brush surface to be treated with a solution containing I ounce of Edco—Syn per gallon of water. Surfaces that are heavily soiled should be cleaned with 2 ounces of Edco—syn per gallon of water and then disinfected. All treated food contact surfaces and or equipment must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse. To disinfect locker rooms, shower stalls, walkways around pools and other areas against pathogenic fungi, mop area with a solution containing I sunce of Edco—Syn in I gallon of water. This solution will destroy Trichophyton mentagrophytes, a causative agent of athletes foot.

HOSPITAL DISINFECTION

To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 1 ounce of Edco—Syn per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.



DETERGENT-DISINFECTANT-DEODORANT-GERMICIDAL-FUNGICIDE-SANITIZER

ACTIVE INGREDIENTS

Sodium 0-benzyl-p-chlorophenate	7%
tetrasodium ethylene-diamine tetraacetate	0%
Sodium p-tert-butylphenate	4%
✓ Isopropyl "Alcohol"	8%
sodium dodecylbenzene sulfonate	
INERT INGREDIENTS82.8	1%

EPA Reg. No. 3040-30


DANGER KEEP OUT OF REACH OF CHILDREN

See First Aid and additional precautions on side panel. Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying in a safe place.

MANUFACTURED BY

Edco Chemical Co. Inc. 1048 Key Road Columbia, S. C. 29205

NET CONTENTS



SANITIZATION

To sanitize walls, floors, and equipment in food processing plants, first clean thoroughly with a solution containing I ounce of Edca-Syn per gallon of water. If surface is heavily soiled, use Edco-Syn at a rate of 2 ounces per gallon to insure thorough cleaning. Sanitize cleaned surface with a rinse solution containing I ounce of Edco-Syn per gallon of water. All treated food contact surfaces and/or equipment mest be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse.

DEODORIZATION

To deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 2 sunces of Edco-Syn per gallon of water. Edco-Syn is an effect tive antibacterial cleaner designed for use in household, industrial and institutional cleaning. When used as directed, Edco-Syn will disinfect hard surfaces, such as floors, walls, counter tops and furniture. 14 may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where obnoxious odors may develop. This product may be used as a general purpose antibacterial-antifungal detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other areas where efficient cleaning and antimicrobial action is desired. See left panel for use directions.

DANGER

KEEP OUT OF REACH OF CHILDREN.

Corrosive. May cause eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucesal demage may contraindicate the use of asstric levage. Measure and convulsion may be needed.