

USE DIRECTIONS

SANITIZATION OF EATING UTENSILS AND FOOD HANDLING EQUIPMENT

Clean dishes, silverware, cooking utensils and food handling equipment with a suitable detergent and rinse to remove food soils. Then immerse in a solution containing 1 ounce of Edco Germolene in 4 gallons of water and let air dry. Do not towel dry. It is not necessary to rinse sanitized articles with potable water before reuse.

SANITIZATION OF FOOD PROCESSING EQUIPMENT

To sanitize beverage and food processing or handling equipment, first clean thoroughly and rinse to remove soils. Then brush or sponge equipment with a solution containing 1 ounce of Edco Germolene in 4 gallons of water. This solution should be circulated through transfer lines, blending vessels and filling equipment to thoroughly sanitize all areas that contact food. Allow equipment to drain dry before reuse.

SANITIZATION OF DAIRY EQUIPMENT

After milking, rinse equipment with potable warm water. Then wash equipment thoroughly with a suitable detergent and rinse with potable water to remove all soil and milk solids. Sanitize by sponging or brushing equipment with a solution containing 1 ounce of Edco Germolene in 4 gallons of water. Freshly prepared sanitizing solution should be circulated through milking machines, transfer lines and bulk holding equipment to assure thorough sanitization of those surfaces that contact milk. Allow equipment to drain dry after sanitization. It is not necessary to rinse sanitized equipment with potable water before reuse.

For udder and flank washing, prepare a solution containing 1 ounce of Edco Germolene in 4 gallons of warm water. Wash udder and flanks, using a fresh towel or sponge for each cow. Avoid contamination of sanitizing solutions with dirt or soil. Be certain the area to be treated is thoroughly wetted with the sanitizing solution.



Edco

GERMOLENE

DISINFECTANT - DEODORIZER - SANITIZER

ACCEPTED
MAY 6 1974
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 3040-37

ACTIVE INGREDIENTS

n-Alkyl (50% C₁₄, 40% C₁₂, 10% C₁₆) dimethyl benzyl ammonium chloride10.0%

INERT INGREDIENTS90.0%
100.0%

EPA Reg. No 3040-37

Edco Germolene is a concentrate and should be diluted before using.

DANGER

KEEP OUT OF REACH OF CHILDREN

See side panel for additional precautions and first aid.

Rinse empty container thoroughly with fresh water and discard it.

MANUFACTURED BY

Edco Chemical Co. Inc.
1048 Key Road
Columbia, S. C. 29205

NET CONTENTS

POULTRY EQUIPMENT DISINFECTION

Poultry brooders, watering founts and feeding equipment may be disinfected after thorough cleaning by sponging or brushing with a solution containing 1 ounce of Edco Germolene in 2 gallons of water. Small utensils should be immersed in this solution. NOTE: All equipment that will contact food or drinking water must be rinsed with water of drinking quality after disinfection and before reuse.

Hatching eggs may be disinfected by dipping in a solution containing 1 ounce of Edco Germolene per 2 gallons of water. Do not use this product on eggs intended for human consumption.

KENNELS AND PET ANIMAL QUARTERS DISINFECTION

To disinfect and deodorize animal quarters mop or sponge a solution containing 1 ounce of Edco Germolene per 2 gallons of water to cleaned floors and walls. Feeding equipment disinfected with this solution should be rinsed with water of drinking quality before reuse.

GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces that will not contact food products in homes, schools, rest rooms and other public places, mop or sponge surface to be treated with a solution containing 1 ounce of Edco Germolene in 2 gallons of water. Surfaces that are soiled must be thoroughly cleaned and rinsed free of soil before disinfection. This disinfectant solution will effectively destroy the gram positive micro-organism, Staphylococcus aureus and the gram negative micro-organism, Salmonella choleraesuis.

HOSPITAL DISINFECTION

To disinfect walls, floors, furnishings and other hard surfaces in hospitals, nursing homes, laboratories and other medical environments, mop or sponge surface with a solution containing 1 ounce of Edco Germolene per gallon of water. Surfaces that are soiled must be thoroughly cleaned and rinsed free of soil before disinfection.

Surgical thorough disinfecting of water sodium gallon corrosia

This c destroy Staphyl micro-o Pseudo

This p F, of t nance, Public (32 gra as CaC by the trizers

DANI

KEEP Corros ritator clothin ling. ination

FIRST

In cas skin minute and wi

If swa of mil these of wat

NOTE

Probal the us circule convul



Edco

GERMOLENE

DISINFECTANT - DEODORIZER - SANITIZER

ACCEPTED
MAY 6 1974
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 3040-37

ACTIVE INGREDIENTS

n-Alkyl (50% C₁₄, 40% C₁₂, 10% C₁₆) dimethyl benzyl ammonium chloride10.0%

INERT INGREDIENTS90.0%

100.0%

EPA Reg. No 3040-37

Edco Germolene is a concentrate and should be diluted before using.

DANGER

KEEP OUT OF REACH OF CHILDREN

See side panel for additional precautions and first aid.

Rinse empty container thoroughly with fresh water and discard it.

MANUFACTURED BY

Edco Chemical Co. Inc.
1048 Key Road
Columbia, S. C. 29205

NET CONTENTS

POULTRY EQUIPMENT DISINFECTION

Poultry brooders, watering founts and feeding equipment may be disinfected after thorough cleaning by sponging or brushing with a solution containing 1 ounce of Edco Germolene in 2 gallons of water. Small utensils should be immersed in this solution. NOTE: All equipment that will contact food or drinking water must be rinsed with water of drinking quality after disinfection and before reuse.

Hatching eggs may be disinfected by dipping in a solution containing 1 ounce of Edco Germolene per 2 gallons of water. Do not use this product on eggs intended for human consumption.

KENNELS AND PET ANIMAL QUARTERS DISINFECTION

To disinfect and deodorize animal quarters mop or sponge a solution containing 1 ounce of Edco Germolene per 2 gallons of water to cleaned floors and walls. Feeding equipment disinfected with this solution should be rinsed with water of drinking quality before reuse.

GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces that will not contact food products in homes, schools, rest rooms and other public places, mop or sponge surface to be treated with a solution containing 1 ounce of Edco Germolene in 2 gallons of water. Surfaces that are soiled must be thoroughly cleaned and rinsed free of soil before disinfection. This disinfectant solution will effectively destroy the gram positive micro-organism, Staphylococcus aureus and the gram negative micro-organism, Salmonella choleraesuis.

HOSPITAL DISINFECTION

To disinfect walls, floors, furnishings and other hard surfaces in hospitals, nursing homes, laboratories and other medical environments, mop or sponge surface with a solution containing 1 ounce of Edco Germolene per gallon of water. Surfaces that are soiled must be thoroughly cleaned and rinsed free of soil before disinfection.

Surgical or medical instruments must be thoroughly scrubbed to remove soil and then disinfected by immersion in a solution containing 1 ounce of Edco Germolene per gallon of water for 10 minutes. One tablespoon of sodium nitrite should be added to each gallon of disinfectant solution to inhibit corrosion of metal instruments.

This disinfectant solution will effectively destroy the gram positive micro-organism Staphylococcus aureus, and the gram negative micro-organisms Salmonella choleraesuis, and Pseudomonas aeruginosa.

This product fulfils the criteria of appendix F, of the grade "A" Pastuerized Milk Ordinance, 1965 recommendations of the U. S. Public Health Service in waters up to 550 ppm (32 grains per gallon) of hardness calculated as CaCO₃ when tested by the method outlined by the AOAC Germicidal and Detergent Sanitizers Official Method.

DANGER

KEEP OUT OF REACH OF CHILDREN
Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.