

ACCEPTED
SEP 11 1991

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg No. **3050-77**

PRECAUTIONARY STATEMENTS
Hazards To Humans and Domestic Animals

Do not get in eyes, or on skin or clothing. Wear goggles or safety glasses while mixing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Do not apply directly to food, water or feed supplements. Do not apply directly to animals.

When using this product in animal housing areas, do not apply as a coarse surface spray directly on food, water or feed supplements.

Do not apply as a coarse surface spray directly on animals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DO NOT USE THIS PRODUCT IN OR ON ELECTRICAL EQUIPMENT DUE TO THE POSSIBILITY OF SHOCK HAZARD.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place away from heat or open flame. Store on a solid, level surface in an upright position. Keep container closed when not in use. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, filler's earth, etc. Dispose of waste at an approved waste disposal facility.

Pesticide Disposal: Pesticide, spray mixture or rinsate may be disposed of on site according to label instructions, or at an approved waste disposal facility.

Container Disposal: Triple Rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

2 of 5

CONTAINER LABEL - FRONT PANEL

PROTECTICIDE_™
COYNE FORMULA 211

**Effective, Fast Acting
PROTECTICIDE_™ Insecticide Provides
Flushing Action and Quick
Knockdown. Specifically Designed
for Use In and Around Poultry
Houses as a Space, Area
or Contact Spray.**

Authorized by USDA for use in edible
product areas of official establishments
operating under the Meat, Poultry, Shell
Egg Grading and Egg Products
Inspection Program.

ACTIVE INGREDIENTS:	
Pyrethrins	0.5%
*Piperonyl Butoxide, Technical	5.0%
INERT INGREDIENTS:	94.5%
TOTAL	100.0%

*Equivalent to 4.00% (butylcarbityl)(6-propylpiperonyl) ether and 1.00% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or poison control center immediately.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

**SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

AVISO

PRECAUCION AL USUARIO

Si usted no inglé's, no use este producto hasta que la etiqueta le haya sido explicada ampliamente

PROTECTICIDE is a trademark of Cardinal Chemical Company Inc. All rights reserved

EPA Reg No. 3050-79
EPA Est. No 3050-CA-1

Coyne Chemical Company
3171 Fitzgerald Road
Rancho Cordova, CA 95742

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

General: This Product has been specifically designed for indoor and outdoor use in, around the perimeter of, and outside of poultry houses and poultry processing plants. For use undiluted or diluted with water and applied as a space spray with cold aerosol generators (ULV), automatic spraying systems, and hand held mechanical foggers or misters. Also for use undiluted or diluted with water and applied as a wet surface spray with mechanical misting equipment or compressed air sprayers.

Official establishments operating under the Federal Meat, Poultry, Shell Egg Grading, and Egg Products Inspection Program: Before using this product, exposed food products and packaging materials must be removed or protected. After using this product, facilities, equipment, and utensils must be washed with an effective cleaning compound and thoroughly rinsed with potable water after spraying. Before food products may be placed back into the area where this product has been used, the area should be sufficiently free of odor to prevent product contamination. Do not apply while food processing is underway.

This product should be used in, around the perimeter of, and outside of poultry houses and poultry processing plants as a space or surface spray for the control of exposed stages of the following insects:


Ants, Hornets, Brown Dog Ticks, Horn Flies, Cockroaches, Horse Flies, Crickets, House Flies, Darkling Beetles, Lice, Deer Flies, Mites, Earwigs, Mosquitoes, Face Flies, Silverfish, Fannia Flies, Scorpions, Firebrats, Small Flying Moths, Fleas, Spiders, Flies, Stored Product Moths, Fruit Flies, Wasps, Fungus Gnats, Yellow Jackets, and other similar insects.

Dilution Instructions:

Use Coyne Formula 211 insecticide to control pests indicated in the areas listed below by application as a 0.10% - 0.05% to control light infestations, or full strength to quickly reduce heavy infestations. To use this product diluted, dilute just prior to use as follows:

0.10% Pyrethrins - Add 1 part Concentrate to 4 parts water.

0.05% Pyrethrins - Add 1 part Concentrate to 9 parts water.



INDOOR USE

Space Spray Application

In poultry houses, poultry processing plants, and poultry ranch equipment and tool sheds as a space spray, undiluted or diluted and applied with conventional ULV cold aerosol generators, automatic spraying systems, and hand held mechanical foggers or misters, for the control of exposed stages of the following insects:

Ants, Hornets, Brown Dog Ticks, Horn Flies, Cockroaches, Horse Flies, Crickets, House Flies, Darkling Beetles, Lice, Deer Flies, Mites, Earwigs, Mosquitoes, Face Flies, Silverfish, Fannia Flies, Scorpions, Firebrats, Small Flying Moths, Fleas, Spiders, Flies, Stored Product Moths, Fruit Flies, Wasps, Fungus Gnats, Yellow Jackets, and other similar insects.

Whenever possible, space spray application should be made in the late evening when insects are at rest. Use a conventional ULV cold aerosol generator, thermal fogger, or fogger adjusted to deliver aerosol size droplets. Close buildings and enclosures as tightly as possible. Apply at a rate of 1-2 ounces per 1000 cubic feet of space. In poultry houses, direct spray toward the upper portions of the enclosure above the birds, filling the barn with mist or fog. For best results, keep treated areas closed for at least 25-30 minutes after treatment. Vacate treated area and ventilate before reentering. Repeat treatment if infestation reoccurs.

OUTDOOR USE

Space Spray Application

For use outside of and around the perimeter of poultry houses, poultry processing plants, and poultry ranch equipment and tool sheds as a space spray, undiluted or diluted and applied with conventional ULV cold aerosol generators, automatic spraying systems, and hand held mechanical foggers or misters, for the control of exposed stages of the following insects:

Ants, Hornets, Brown Dog Ticks, Horn Flies, Cockroaches, Horse Flies, Crickets, House Flies, Darkling Beetles, Lice, Deer Flies, Mites, Earwigs, Mosquitoes, Face Flies, Silverfish, Fannia Flies, Scorpions, Firebrats, Small Flying Moths, Fleas, Spiders, Flies, Stored Product Moths, Fruit Flies, Wasps, Fungus Gnats, Yellow Jackets, and other similar insects.

Whenever possible, space spray application should be made in the late evening when insects are at rest. Apply when there is a light breeze approximately 5 miles per hour. Spray from the windward side so the mist or fog will be carried across the area to be treated. Outside treatment includes, but is not limited to, such areas as undersides of eaves, around farm equipment, above and around exposed manure piles, in and around shrubbery and vegetation, stagnant pools, marshy areas and other similar places where insects may rest or hide. Apply at a rate of 1-2 ounces per 1000 cubic feet. Repeat treatment if infestation reoccurs.

INDOOR/OUTDOOR USE

Surface Spray Application for Crawling Insects

For use as a surface spray, undiluted or diluted and applied with mechanical misting equipment or compressed air sprayers in, around the perimeter of, and outside of poultry houses, poultry processing plants, and poultry ranch equipment and tool sheds for the control of exposed stages of the following crawling insects:

Ants, Earwigs, Dog Ticks, Cockroaches, Crickets, Darkling Beetles, Lice, Mites, Earwigs, Silverfish, Spiders, Firebrats, Fleas, Spiders, and other similar insects.

Apply as a coarse, wet surface spray including, but not limited to such areas as trails, harborage areas, interior walls, exterior walls, nests, points of entry, dumpsters, tool sheds, gate houses, cracks and crevices, ranch equipment, beams, trash cans, ceilings, behind equipment, moldings, doors, around window frames, undersides of eaves, rafters, foundations, tractors, electric carts, manure piles, poultry hauling equipment, building foundations and similar insect hiding places. Spray liberally contacting as many insects as possible. Repeat treatment if infestation reoccurs.

Surface Spray Application for Resting Flying Insects

For use as a surface spray, undiluted or diluted and applied with mechanical misting equipment or compressed air sprayers in, around the perimeter of, and outside of poultry houses, poultry processing plants, and poultry ranch equipment and tool sheds for the control of exposed stages of the following resting flying insects:

Hornets, Horn Flies, Horse Flies, House Flies, Deer Flies, Mosquitoes, Face Flies, Fannia Flies, Small Flying Moths, Flies, Stored Product Moths, Fruit Flies, Wasps, Fungus Gnats, Yellow Jackets, and other similar insects.

Whenever possible, surface application should be made in late evening when insects are at rest. Apply as a coarse, wet surface spray including, but not limited to such areas as trails, harborage areas, interior walls, exterior walls, nests, points of entry, dumpsters, tool sheds, gate houses, cracks and crevices, ranch equipment, beams, trash cans, ceilings, behind equipment, moldings, doors, around window frames, undersides of eaves, rafters, foundations, tractors, electric carts, manure piles, poultry hauling equipment, building foundations, and similar insect hiding places. Spray liberally contacting as many insects as possible. Repeat treatment if infestation reoccurs.

WARRANTY STATEMENT: Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product is beyond the control of the manufacturer. No guarantee, expressed or implied, is made as to the effect of such or the results to be obtained if not used in accordance with the directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such or in combination with other materials.