

**DANGER—POISON—PRECAUTIONS** ☠

**CAN KILL YOU IF SWALLOWED**

This product can kill you if swallowed even in small amounts: spray mist or dust may be fatal if swallowed.


**CAN KILL YOU BY SKIN CONTACT**

This product can kill you if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes)



**CAN KILL YOU IF BREATHED**

This product can kill you if vapors, spray mist or dust are breathed.



**POISONOUS TO FISH AND WILDLIFE**

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where birds are likely to roost.

Do not use on or near any food crops, food, feed, or equipment. Do not use on or near any animal. Apply this product only as specified on this label.


**AGRICULTURAL PRODUCTS COMPANY, INC.**  
ARTESIA, NEW MEXICO — MESQUITE, NEW MEXICO

See panels below  
for antidote  
and precautions

**STOP! READ THE LABEL**



Keep out of reach  
of children



**BIOTROL BTB PARATHION 2-2**

**ACTIVE INGREDIENTS:**

Parathion\* 2.00%

Bacillus thuringiensis Berliner (Type 200) (Entomopathogenic) 0.06%

Inert Ingredients 97.94%

**INERT INGREDIENTS:**

Equivalent to 100% active ingredient

BIOTROL XK Reg. T.M. Nutrilite Products, Inc.

50 lbs. Net Weight

EPA Reg. No. 3051-64

**NOT FOR HOME USE**

EPA Est. No. 3051-NM-1

### USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

(1) Wear water proof pants, coat, hat, rubber boots or rubber overshoes. (2) Wear safety goggles. (3) Wear mask or respirator approved by the U.S. Bureau of Mines for parathion protection. (4) Wear heavy duty, natural rubber gloves.

### WORK SAFETY RULES


Keep all unprotected persons and children away from treated area or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. Call a doctor (physician) clinic or hospital—immediately. Explain that the victim has been exposed to parathion and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital.

**IMPORTANT:** Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

**AFTER WORK,** take off all work clothes and shoes. Shower using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

EPA Reg. No. 3051-64



### SYMPTOMS OF POISONING

Parathion is a very dangerous poison. It rapidly enters the body on contact with all skin surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment or they may die.

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, cramps, weakness, blurred vision, pin point pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooling or frothing of mouth and nose, muscle spasms and coma.

### FIRST AID TREATMENT

Call a doctor (physician) clinic or hospital immediately. Explain that the victim has been exposed to parathion and describe his condition.

**If breathing has stopped,** start artificial respiration immediately and maintain until doctor sees victim. **If swallowed** and victim is awake (conscious) make him vomit quickly. Induce vomiting by sticking finger down throat or by giving soapy or strong salty water to drink. Repeat until vomit is clear. **Never** give anything by mouth to an unconscious person. Have victim lie down and keep quiet. See doctor immediately.

**In case of contact,** immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. See doctor immediately.

### NOTE TO PHYSICIAN

Antidote—administer atropine sulfate in large doses, TWO to FOUR mg. intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 10 minute intervals until signs of atropinization appear. 2-PAM (pralidoxime) is an antidotal and may be administered in conjunction with atropine. **DO NOT GIVE MORPHINE OR TRANQUILIZERS.** Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement. **VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.**

### POST TREATED AREA

Consult your State Agricultural Extension Service or Experiment Station regarding posting treated areas.

### CAUTION

Do not eat, drink, or apply cosmetics. Destroy by burning in the water or by incineration. Stay away from smoke or fumes.

### DIRECTIONS FOR USE

**LETTUCE-CABBAGE:** For control of cabbage looper, use 25 pounds per acre. Apply dust so that all foliage is thoroughly covered. Do not apply to head lettuce within 7 days of harvest. Do not apply to cabbage within 10 days of harvest.

### NOTICE TO BUYER

Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label use directions or under abnormal use conditions, or under conditions not reasonable foreseeable to seller. Buyer assumes all risk of any such use. Seller makes no other warranties, express or implied.

### CAUTION

This product is very toxic to bees exposed to it during flowering. Do not apply this material to blooming crops or weeds where bees are active. For more information, contact your State Agricultural Extension Service.

