

4-10-53

PARATHION 8
Insecticide

100
100
100
100

DANGER!

KEEP OUT OF REACH OF CHILDREN

★ POISON ★

See Side Panel For Antidote And Other Required Warning Statements

USDA Reg. No. 3051-

Active Ingredients:

Parathion (O,O-diethyl O-p-nitrophenyl phosphorothioate)	81.7%
Pyridine	6.2%
Other Ingredients	12.1%

(1 gallon contains 8 lbs. parathion)

Net Contents: Gallon

Net Contents: Gallon



WARNING

KEEP AWAY FROM CHILDREN

POISON

CAUTION DO NOT EAT

DONT

PRUITT CO.

MEXICO

9/23/68
3051-65

DANGER

AVOID SKIN CONTACT, INHALATION OR SWALLOWING
POISONOUS AND IRRITANT THROUGH SKIN
REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS,
PROGRESSIVELY INCREASE SUSCEPTIBILITY TO PARATHION POISON.

(a) Not to be Used In Eyes, On Skin, Or Clothing

Wear clean heavy rubber gloves and clean waterproof or freshly laundered protective clothing (overalls, cap, etc.) Clothing must not get wet through. Destroy and replace gloves frequently. Wash thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing and protective clothing thoroughly with soap and hot water before re-use. In case of spill, immediately remove contaminated clothing and wash skin thoroughly with soap and water.

(b) Not to be Used As Vapor or Spray Mist

Wear respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Airplane pilots should wear full-face, canister-type masks. If unmasked, provide mechanical exhaust ventilation. Remove unprotected persons and domestic animals from operating and adjacent areas where drift may drift; prohibit re-entry until drift and vapor dissipate.

(c) Not to be Used As Fertilizer Feed Products

Do not use as a fertilizer or feed product such as a soda ash, lime, clay or sawdust. Sweep up and remove wash area thoroughly with a full-strength liquid household chlorine bleach.

Wipe up spillage with a cloth. Drain pail completely. Add 1/2 gallon water, 1/2 cup of household bleach and 1/2 cup of lye. Tighten closure. Rotate pail to wet all surfaces and let stand for 15 minutes. Drain completely and rinse several times with water. Tighten closure. Puncture and crush pail to prevent re-use.

COMBUSTIBLE

Do not use near open flame or store near heat or open flame. Keep container tightly closed with adequate ventilation. Boiling point 100°C (212°F).

POISON

ANTIDOTE: Atropine is an antidote. Consult your physician about obtaining a supply of atropine tablets for emergency use. Never take atropine unless symptoms of poisoning have occurred.

FIRST AID: In symptoms or signs of poisoning (See Note to Physician) include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor. Give two atropine tablets (each 1/100 grain) at once. Call a doctor at once in all cases of suspected poisoning.

Remove contaminated clothing and

unprotected persons and domestic animals from operating and adjacent areas where drift may drift; prohibit re-entry until drift and vapor dissipate.

Do Not Use on Plants, Foliage or Feed Products

Do not mix with an absorbent such as a soda ash, lime, clay or sawdust.

Wash, up to 1000, wash area thoroughly with a full strength liquid household

disinfectant.

Wash hands and face completely. Add 1/2 gallon water to cup of
the paint and pour it over. Tighten closure. Rotate pail to wet all surfaces
and let it stand for at least 15 minutes. Drain completely and rinse several times
with water. Tighten closure. Puncture and crush pail to prevent reuse.

COMBUSTIBLE

Do not use near a fire or store near heat or open flame. Keep container
closed with adequate ventilation.

Keep away from temperatures below 0° F.

✕ POISON ✕

ANTIDOTE: Atropine is an antidote. Consult your physician about obtaining a
supply of 1/100 grain tablets for emergency use. Never take atropine unless
symptoms of poisoning have occurred.

FIRST AID: If symptoms or signs of poisoning (See Note to Physician) include
blurred vision, abdominal cramps and tightness in the chest, don't wait for a
doctor but give two atropine tablets (each 1/100 grain) at once.

Call a physician at once in all cases of suspected poisoning.

Remove patient immediately from the area. Remove contaminated clothing and
wash the skin clean with plenty of soap and water to remove all traces of the
pesticide. If swallowed, induce vomiting by giving warm salty or soapy water.
Have patient lie down and keep quiet.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness in
chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea,
vomiting, diarrhea and abdominal cramps.

TREATMENT: Give atropine intramuscularly or intravenously 2 to 4 milligrams
(1 to 2 tablets 1/100 grain each) at once and every hour as required until
patient is awake. Twenty to 30 milligrams or more may be required during the
first 24 hours. Never give morphine or phenothiazine tranquilizers. Clear
chest by natural drainage. Artificial respiration or oxygen administration

may be necessary to keep patient continuously 48 hours. Repeated exposure to
this pesticide may without warning, cause increasing susceptibility
to very small doses of any cholinesterase inhibitor. Allow no further exposure
until complete recovery has taken place as determined by blood tests.

This product is highly toxic to fish and wildlife. Do not contaminate any body
of water, especially when conditions favor drift of spray from areas treated. To
protect bees, do not spray when crops are in bloom. Remove or cover hives if
spraying is necessary during bloom.

9/23/68
3051-65

DISCLAIMER

Agricultural Products Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonable fit for the use therein described when used in accordance with the directions for use. Any damage resulting from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc. Agricultural Products Company makes no other express or implied warranty including any implied warranty of FITNESS or of MERCHANTABILITY. BUYER assumes the risk of any use contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable by Agricultural Products Company.

DIRECTIONS FOR USE

Do not use this product for any uses other than those specified on this label. Application should be made only by experienced personnel using approved safety equipment. Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do Not allow mixture to stand. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible. Unless otherwise noted, apply recommended rates by ground equipment in not less than 10 gals. of water per acre; by aircraft equipment, in 2 to 10 gals. of water per acre consistent with crop growth and good coverage. Consult your state agricultural experiment station or agricultural extension service for proper timing of applications.

VEGETABLE, VINE AND FIELD CROPS

BEANS - LAYER WETCH: Aphids and Alfalfa Weevil Larvae-Use $\frac{1}{8}$ pt. per acre. Apply within 15 days of harvest or forage use.
BEANS - MOTTLED: Armyworm, Leafhopper, Leaf Miner, Leaf Roller and Mexican Bean Beetle-Use $\frac{1}{8}$ pt. per acre. Repeat applications at 7-10 day intervals until control is obtained. Do not apply within 15 days of harvest.
CABBAGE: Aphids, Armyworm, Diamond-back Moth, Imported Cabbageworm, Thrips-Use $\frac{1}{8}$ pt. per acre. Do not apply to cabbage, kale mustard and turnips within 10 days of harvest; 7 days on broccoli, Brussels sprouts and cauliflower.
CORN: Army worm, Corn Rootworm Adults-Use $\frac{1}{8}$ pt. per acre. For Corn Rootworm, make full coverage applications to foliage when adult beetles become abundant and repeat as necessary. Do not apply within 12 days of picking, cutting or forage.

combined has proved them to be physically compatible.
Unless otherwise noted, apply recommended rates by ground equipment in not less than 10 gals. of water per acre; by aircraft equipment, in 2 to 10 gals. of water per acre consistent with crop growth and good coverage.
Consult your state agricultural experiment station or agricultural extension agent for proper timing of applications.

VEGETABLE, VINE AND FIELD CROPS

ALFALFA (CLOVER WETCH): Aphids and Alfalfa Weevil Larvae—Use $\frac{1}{8}$ pt. per acre. Do not apply within 15 days of harvest or forage use.

BEANS: Aphids, Armyworm, Leafhopper, Leaf Miner, Leaf Roller and Mexican Bean Beetle—Use $\frac{1}{2}$ pt. per acre. Repeat applications at 7-10 day intervals until controlled. Do not apply within 15 days of harvest.

COLE CROPS (Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kale, Mustard, Turnip): Aphids, Armyworm, Diamond-back Moth, Imported Cabbageworm, Thrips—Use $\frac{1}{2}$ pt. per acre. Do not apply to cabbage, kale mustard and turnips within 10 days of harvest; 7 days on broccoli, Brussels sprouts and cauliflower.

CORN: Army worm, Corn Rootworm Adults—Use $\frac{1}{2}$ pt. per acre. For Corn Rootworm, make full-coverage applications to foliage when adult beetles become abundant and repeat as necessary. Do not apply within 12 days of picking, cutting or forage.

COTTON: Cotton Aphid, Cotton Leafworm, Spider Mites, Stinkbugs—Apply $\frac{1}{2}$ pt. per acre. Cotton Leaf Perforator, Fleahoppers, Lygus Bugs—Apply 1 pt. per acre. Do not apply within 5 days of hand picking.

CUCURBITS (Cucumber, Squash and Melon): Aphids, Cucumber Beetle, Melonworm, Blightworm, Serpentine Leafminer, Squash Bug and Rindworm on watermelons—Use $\frac{1}{2}$ pt. per acre. Do not apply unless plants are dry not until runners begin to wilt. Do not apply within 15 days of harvest on cucumbers and squash. 7 days on melon.

EGGPLANT: Aphids, Leaf Miners, Red Spider Mites, Thrips—Apply $\frac{1}{2}$ pt. per acre. Do not apply within 10 days of harvest.

LETTUCE: Aphids, Leafhoppers, Lygus Bug—Use $\frac{1}{2}$ pt. per acre. Do not apply within 10 days of harvest on head lettuce; 21 days on leaf and bibb lettuce.

OKRA: Aphids, Leaf Miners, Stinkbugs—Apply $\frac{1}{2}$ pt. per acre. Do not apply within 10 days of harvest.

PEAS: Aphids, Pea Weevil—Use $\frac{1}{2}$ pt. per acre. Do not apply within 10 days of harvest.

PEPPERS: Flea Beetle, Serpentine Leaf Miner, Pepper Maggot, Thrips—Use $\frac{1}{2}$ pt. per acre. Do not apply within 15 days of harvest.

POTATO (TUBER): Aphids, Flea Beetle, Leafhopper, Looper, Serpentine Leaf Miner, Blightworm—Use $\frac{1}{2}$ pt. per acre. Do not apply within 10 days of harvest on tomatoes and 10 days of harvest on potatoes.

SMALL GRAINS (Barley, Oats, Wheat): Aphids, Armyworms, Greenbug—Use $\frac{3}{8}$ pt. per acre. Do not apply within 15 days of harvest.

WATERMELON: Aphids, Blister Beetles, Flea Beetles, Leafhoppers, Leaf Miners, Rindworm—Use $\frac{1}{2}$ pt. per acre. Do not apply within 15 days of harvest.