

PARATHION 4
Insecticide

DANGER!

KEEP OUT OF REACH OF CHILDREN

☠ POISON ☠

See Side Panel For Antidote and Other Warnings

USDA Reg. No. 3051

Active Ingredients:

Parathion (4 lbs. parathion)	45.7%
Xylene (methyl aromatic hydrocarbon solvent)	42.5%
Inert Ingredients	11.8%
*Contains 4 lbs. parathion	

Net Contents: Gallon

ACCEPTED

SEP 24 1968

FEDERAL INSECTICIDE
AND FUNGICIDE ACT
POISON REGISTER
SUBJECT
TO ALL REGULATIONS

[Faint, illegible text, possibly a signature or stamp]

PARATHION 4
DONN

SEP 24 1966
UNDER THE FEDERAL
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 3251-66
TO ATTACHED COMMISSION



WARNING
KEEP AWAY FROM FEED or FOOD PRODUCTS
POISON

CAUTION—DO NOT DROP
IF LEAKING **DON'T** BREATHE FUMES
TOUCH CONTENTS
SWALLOW

This is to certify that the contents of this package
are properly described by name and are packed
and marked and are in proper condition for
transportation according to the Regula-
tions prescribed by the Interstate
Commerce Commission.

Shipper's name required
hereon for ship-
ments by
EXPRESS

AGRICULTURAL PRODUCTS CO.

MESQUITE, NEW MEXICO

AGCO

ARTESIA, NEW MEXICO

19/20

DANGER!

9/24/66

3051-66

POISONOUS BY SKIN CONTACT, INHALATION OR SWALLOWING

RAPIDLY ABSORBED THROUGH SKIN

REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO PARATHION POISONING.

Do Not Get In Eyes, On Skin, On Clothing

Wear clean, heavy rubber gloves and clean waterproof or freshly laundered protective clothing (coveralls, cap, etc.). Clothing must not get wet through. Destroy and replace gloves frequently. Wash thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water.

Do Not Breathe Vapor Or Spray Mist

Wear respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Airplane pilots should wear full face, canister-type mask. If handled indoors, provide mechanical exhaust ventilation. Remove unprotected persons and domestic animals from operating and adjacent areas where insecticide may drift; prohibit re-entry until drift and vapor dissipate.

Do Not Contaminate Food Or Feed Products

Cover spillage with an absorbent such as a soda ash, lime, clay or sand. Sweep up and bury. Wash area thoroughly with a full-strength, liquid household chlorine bleach. This product is toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label. Disposal Of Pails: Drain pail completely. Add $\frac{1}{2}$ gallon water, $\frac{1}{4}$ cup of detergent and $\frac{1}{4}$ pound of lye. Tighten closure. Rotate pail to wet all surfaces and let stand for at least 15 minutes. Drain completely and rinse several times with water. Tighten closure. Puncture and crush pail to prevent re-use.

COMBUSTIBLE

Do not use, pour, spill or store near heat or open flame. Keep container closed. Use with adequate ventilation.

POISON *

ANTIDOTE: Atropine is an antidote. Consult your physician about obtaining a supply of 1/100 grain tablets for emergency use. Never take atropine unless symptoms of poisoning have occurred.

FIRST AID: If symptoms or signs of poisoning (See Note to Physicians) include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets (each 1/100 grain) at once.

Call a physician at once in all cases of suspected poisoning.

Remove patient immediately from the area. Remove contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces of the pesticide.

parathion protection. Airplane pilots should wear full face, canister-type mask. If handled indoors, provide mechanical exhaust ventilation. Remove unprotected persons and domestic animals from operating and adjacent areas where insecticide may drift; prohibit re-entry until drift and vapor dissipate.

Do Not Contaminate Food Or Feed Products

Cover spillage with an absorbent such as a soda ash, lime, clay or sand. Sweep up and bury. Wash area thoroughly with a full-strength, liquid household chlorine bleach. This product is toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label.

Disposal Of Pails: Drain pail completely. Add $\frac{1}{2}$ gallon water, $\frac{1}{4}$ cup of detergent and $\frac{1}{4}$ pound of lye. Tighten closure. Rotate pail to wet all surfaces and let stand for at least 15 minutes. Drain completely and rinse several times with water. Tighten closure. Puncture and crush pail to prevent re-use.

COMBUSTIBLE

Do not use, pour, spill or store near heat or open flame. Keep container closed. Use with adequate ventilation.

POISON

ANTIDOTE: Atropine is an antidote. Consult your physician about obtaining a supply of 1/100 grain tablets for emergency use. Never take atropine unless symptoms of poisoning have occurred.

FIRST AID: If symptoms or signs of poisoning (See Note to Physicians) include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets (each 1/100 grain) at once.

Call a physician at once in all cases of suspected poisoning.

Remove patient immediately from the area. Remove contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces of the pesticide. If swallowed, induce vomiting by giving warm salty or soapy water. Have patient lie down and keep quiet.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Give atropine intramuscularly or intravenously, 2 to 4 milligrams (3 to 6 tablets, 1/100 grains each) at once and every hour as required until pupils dilate. Twenty to 30 milligrams or more may be required during the first 24 hours. Never give morphine or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause increasing susceptibility inhibitor. Allow no further exposure until cholinesterase regeneration has taken place as determined by blood tests.

Do not use this product for any uses other than those specified on this label.

DIRECTIONS FOR USE
DISCLAIMER

9/24/68

3051-66

Agricultural Products Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

Agricultural Products Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

BUYER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by Agricultural Products Company.

Application should be made only by experienced personnel using approved safety equipment.

Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do not allow mixture to stand.

If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible.

When applying by ground equipment, use specified dosage in not less than 10 gallons of water per acre; by aircraft equipment, in 2-5 gallons of water per acre.

BEANS: Aphids, Armyworms, Leafhoppers, Leaf Miner, Leaf Roller and Mexican Bean Beetle—Use 1 pt. per acre in sufficient water for thorough coverage. Repeat applications at 7 to 10-day intervals until controlled. Do not apply within 15 days of harvest.

COLE CROPS (Cabbage, Broccoli, Brussel Sprouts, Cauliflower, Kale, Mustard, Turnip): Aphids, Armyworm, Diamond-back Moth, Imported Cabbageworm, Thrips—Use $\frac{1}{2}$ pt. per acre in sufficient water for thorough coverage. Do not apply within 7 days of harvest.

COTTON: Aphids, Red Spider Mites, Stinkbugs, Loopers—Use $\frac{1}{2}$ to $\frac{3}{4}$ pt per acre in sufficient water for thorough coverage. Do not apply within 5 days of hand-picking.

CUCURBITS (Cucumbers, Squash and Melons): Aphids, Cucumber Beetle, Melonworm, Pickleworm, Serpentine Leaf Miner—Use 1 pt. per acre in sufficient water for thorough coverage. Do not apply unless plants are dry nor until runners begin to form. Do not apply within 15 days of harvest on cucumbers and squash. 7 days on melons.

SPRAYER CROPS (Alfalfa, Beans, Clover, Corn, Fava Beans, Soybeans, Sunflower)

normal conditions, or under conditions not reasonably foreseeable by Agricultural Products Company.

Application should be made only by experienced personnel using approved safety equipment.

Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do not allow mixture to stand.

If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible.

When applying by ground equipment, use specified dosage in not less than 10 gallons of water per acre; by aircraft equipment, in 2-5 gallons of water per acre.

BEANS: Aphids, Armyworms, Leafhoppers, Leaf Miner, Leaf Roller and Mexican Bean Beetle—Use 1 pt. per acre in sufficient water for thorough coverage. Repeat applications at 7 to 10-day intervals until controlled. Do not apply within 15 days of harvest.

COLE CROPS (Cabbage, Broccoli, Brussel Sprouts, Cauliflower, Kale, Mustard, Turnip): Aphids, Armyworm, Diamond-back Moth, Imported Cabbageworm, Thrips—Use $\frac{1}{2}$ pt. per acre in sufficient water for thorough coverage. Do not apply within 7 days of harvest.

COTTON: Aphids, Red Spider Mites, Stinkbugs, Loopers—Use $\frac{1}{2}$ to $\frac{3}{4}$ pt per acre in sufficient water for thorough coverage. Do not apply within 5 days of hand-picking.

CUCURBITS (Cucumbers, Squash and Melons); Aphids, Cucumber Beetle, Melonworm, Pickleworm, Serpentine Leaf Miner—Use 1 pt. per acre in sufficient water for thorough coverage. Do not apply unless plants are dry nor until runners begin to form. Do not apply within 15 days of harvest on cucumbers and squash. 7 days on melons.

FORAGE CROPS (Alfalfa, Barley, Clover, Oats, Vetch, Wheat): Aphids, Armyworms, Blister Beetles, Grasshoppers, Green Bugs—Use $\frac{1}{2}$ to $\frac{3}{4}$ pt. per acre in sufficient water for thorough coverage at first sign of infestation. Do not apply within 15 days of harvest or forage use.

LETTUCE: Aphids, Leafhoppers—Use $\frac{1}{2}$ pt. per acre in sufficient water for through coverage. Do not apply within 7 days of harvest on head lettuce; or within 14 days of harvest on leaf and bibb lettuce.

PEPPER: Aphids, Flea Beetle, Serpentine Leaf Miner, Pepper Maggot—Use $\frac{1}{2}$ pt. per acre in sufficient water for thorough coverage. Do not apply within 15 days of harvest.

POTATO, TOMATO: Aphids, Flea Beetle, Leafhopper, Serpentine Leaf Miner—Use $\frac{1}{2}$ pt. per acre in sufficient water for thorough coverage. Do not apply within 10 days of harvest on tomatoes and 5 days of harvest on potatoes.

Protect from temperatures below 0° F.