



Not Content: ~~Label~~

**ACCEPTED**  
**NOV 5 1968**  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROENTIGIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. **3051-67** SUBJECT  
TO ATTACHED COMMENTS.

AGRICULTURAL PRODUCTS CO.

MESQUITE, NEW MEXICO

AGCO

ARTESIA, NEW MEXICO

25/26

**DANGER!**

**POISONOUS BY SKIN CONTACT, INHALATION OR SWALLOWING**

**RAPIDLY ABSORBED THROUGH SKIN**

**REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO METHYL PARATHION POISONING.**

**Do Not Get In Eyes, On Skin, On Clothing**

Wear clean heavy rubber gloves, goggles and clean waterproof or freshly laundered protective clothing (coveralls, cap, etc.). Clothing must not get wet through. Destroy and replace gloves frequently. Wash thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water.

**Do Not Breathe Vapor Or Spray Mist**

Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for methyl parathion protection. Airplane pilots should wear full-face, canister-type mask. If handled indoors, provide mechanical exhaust ventilation. Remove unprotected persons and domestic animals from operating and adjacent areas where insecticide may drift; prohibit re-entry until drift and vapor dissipate.

**Do Not Contaminate Food Or Feed Products**

Cover spillage with an absorbent such as a soda ash, lime, clay or sand. Sweep up and bury. Wash area thoroughly with a full-strength, liquid household chlorine bleach.

**DISPOSAL OF PAIL:** Drain pail completely. Add 2 gallon of water, 1/4 cup of detergent and 1/2 pound of lye. Tighten closure. Rotate pail to wet all surfaces and let stand for at least 15 minutes. Drain completely and rinse several times with water. Tighten closure. Puncture and crush pail to prevent re-use.

**COMBUSTIBLE**

Do not use, pour, spill or store near heat or open flame. Keep container closed. Use with adequate ventilation.

✕ POISON ✕

**ANTIDOTE:** Atropine is an antidote for methyl parathion. Consult your physician about obtaining a supply of 1/100 grain tablets for emergency use. Never take atropine unless symptoms of poisoning have occurred.

**FIRST AID:** If symptoms or signs of poisoning (See Note to Physician) include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets (each 1/100 grain) at once. Call a physician at once in all cases of suspected poisoning.

Remove patient immediately from the area. Remove contaminated clothing and wash

Insecticide may drift, prohibit re-entry until drift and vapor dissipate.  
**Do Not Contaminate Food Or Feed Products**  
Cover spillage with an absorbent such as a soda ash, lime, clay or sandust.  
Sweep up and bury. Wash area thoroughly with a full-strength, liquid household chlorine bleach.

**DISPOSAL OF PAILS:** Drain pail completely. Add  $\frac{1}{2}$  gallon of water,  $\frac{1}{4}$  cup of detergent and  $\frac{1}{4}$  pound of lye. Tighten closure. Rotate pail to wet all surfaces and let stand for at least 15 minutes. Drain completely and rinse several times with water. Tighten closure. Puncture and crush pail to prevent re-use.

#### COMBUSTIBLE

Do not use, pour, spill or store near heat or open flame. Keep container closed. Use with adequate ventilation.

✕ POISON ✕

**ANTIDOTE:** Atropine is an antidote for methyl parathion. Consult your physician about obtaining a supply of 1/100 grain tablets for emergency use. Never take atropine unless symptoms of poisoning have occurred.

**FIRST AID:** If symptoms or signs of poisoning (See Note to Physician) include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets (each 1/100 grain) at once.

Call a physician at once in all cases of suspected poisoning.

Remove patient immediately from the area. Remove contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces of the pesticide. If swallowed, induce vomiting by giving warm salty or soapy water. Have patient lie down and keep quiet.

**NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

**NOTE TO PHYSICIANS:** Warning symptoms of methyl parathion poisoning include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

**TREATMENT:** Give atropine intramuscularly or intravenously, 2 to 4 milligrams (3 to 6 tablets, 1/100 grain each) at once and every hour as required until pupils dilate. Twenty to 30 milligrams or more may be required during the first 24 hours. Never give morphine or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause increasing susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has taken place as determined by blood tests.

**NOTE:** Inspect area temperatures below 45° F.

Do not use this product for any uses other than those specified on this label.

This product is toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label. Birds and other wildlife in treated areas may be killed.

INSTRUCTIONS FOR USE  
DISCLAIMER

The Agricultural Products Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

Agricultural Products Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

BUYER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by Agricultural Products Company.

Application should be made only by experienced personnel using approved safety equipment.

Four recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do not allow mixture to stand.

If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible.

Make full coverage application and repeat as necessary.

**PEACH AND APRICOT:** European red mites and green peach aphid—Apply  $\frac{1}{2}$  pt. per 100 gallons of water. Apply twice at 5-day intervals for control of European red mites. Do not apply within 14 days of harvest.

**VEGETABLE AND FIELD CROPS:** Rates are given in terms of pints of METHYL PARATHION 4 E. G. per acre. When applying by ground equipment, use specified dosage in not less than 10 gallons of water per acre; by aircraft equipment in 2-5 gallons of water per acre. The number shown in parentheses() indicates the number of days which should be allowed between last application of harvest or grazing. Unless otherwise specified, this product will control aphids, armyworms(up to 3rd instar) loopers, flea beetles, leafhoppers and mites on the following crops:

**ALFALFA, CLOVER, FETTER:** Spotted alfalfa aphid— $\frac{1}{2}$  pt. Apply in early spring. Follow state recommendations. For the control of aphids, armyworms(up to 3rd instar), leafhopper and mites— $2\frac{1}{2}$  pts.(20).

**ALFALFA HAY:** Alfalfa weevil larvae—Apply  $\frac{3}{4}$  pt. per acre by ground or air. Apply when buds and shoots first appear ragged or when larvae count first reaches 25 per sweep (15).

**ARTICHOKES, BEETS, CARROTS, CELERY, PEAS, PEPPERS:** 1 pt.(7) artichokes, (15) other crops listed except (21) days on beets if tops are to be used for food.

**BEAN:** (green) 2 pts. (21)

**BEAN (except green), TOMATOES:** 2 pts. (15).

**COLE CROPS:** Cabbage, Cauliflower, Broccoli, Brussels sprouts, Collards, Kale and Kohlrabi—Use 2 pts. (21).

**COTTON:** Aphids, boll weevil, cotton leafworm, leafhopper, lygus, red spider mites, thrips, cotton leaf perforator— $\frac{1}{2}$ -1 pt. Application should be made at 4-5 day intervals until control is obtained.

For the control of armyworms(up to 3rd instar) on cotton:  $2\frac{1}{2}$  pts. Workers entering treated fields within 24 hours of application should wear protective clothing (5).

**LETTUCE, SPINACH:** 2 pts. (21).

**POTATOES:** 2 pts. (5).

**SMALL GRAINS:** Barley, Oats, Rye and wheat; Aphids(green bug), black grass bug (tribe), brown wheat mite(Petrolia) and stink bugs—Apply  $\frac{1}{2}$ -1 pt. For armyworms(up to 3rd instar) and leafhoppers, apply  $2\frac{1}{2}$  pts. Use limitations:  $\frac{1}{2}$  pt. once, over  $\frac{1}{2}$  and up to  $1\frac{1}{2}$  pts.—15 days; over  $1\frac{1}{2}$  pts.—Do not apply after heads begin to form.