

SANITIZED BRAND MILDEW PREVENTATIVE  
For Commercial & Industrial Use

Active Ingredient	Dodecyl Dimethyl Ammonium Chloride	50%
Inert Ingredient		50%
		100%

Use: A mildew preventive for commercial & institutional laundries.

**DANGER** - Keep Out Of Reach Of Children.

Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

**FIRST AID:**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:**

Possible mucosal damage may contraindicate the use of gastric lavage. Measures may be necessary such as respiratory depression, and convulsion may be required.

Directions for use for mildew prevention, use Sanitized Brand Preventative as follows:

For 2 week protection use 2 ounces per 100 pounds of fabric (dry weight).

For 4 week protection use 4 ounces per 100 pounds of fabric (dry weight).

For 6 week protection use 6 ounces per 100 pounds of fabric (dry weight). For 8 week protection use 8 ounces per 100 pounds of fabric (dry weight).

For 10 week protection use 10 ounces per 100 pounds of fabric (dry weight). For 12 week protection use 12 ounces per 100 pounds of fabric (dry weight).

For 14 week protection use 14 ounces per 100 pounds of fabric (dry weight). For 16 week protection use 16 ounces per 100 pounds of fabric (dry weight).

For 18 week protection use 18 ounces per 100 pounds of fabric (dry weight). For 20 week protection use 20 ounces per 100 pounds of fabric (dry weight).

For 22 week protection use 22 ounces per 100 pounds of fabric (dry weight). For 24 week protection use 24 ounces per 100 pounds of fabric (dry weight).

For 26 week protection use 26 ounces per 100 pounds of fabric (dry weight). For 28 week protection use 28 ounces per 100 pounds of fabric (dry weight).

For 30 week protection use 30 ounces per 100 pounds of fabric (dry weight). For 32 week protection use 32 ounces per 100 pounds of fabric (dry weight).

For 34 week protection use 34 ounces per 100 pounds of fabric (dry weight). For 36 week protection use 36 ounces per 100 pounds of fabric (dry weight).

ACCEPTED  
UNDER THE MILDEW PREVENTATIVE ACT  
FOR ELON UNIVERSITY  
MAY 1964