

SUPERIOR

50 LB. NET

WETTABLE SULPHUR

ACTIVE INGREDIENTS:

Sulphur Not Less Than 90.0%

INERT INGREDIENTS:

Not More Than 10.0%

Sulphur Fineness: 93% Passing No. 325 U.S.S. Sieve 100.0%

OBSERVE THE FOLLOWING USE PRECAUTIONS

Apply in early morning or evening when there is little or no wind.

Do not apply to plants (tomatoes, cucumbers, squash, melon, or young plants) which may be injured by the result.

Do not apply to foliage of apples, tung trees, citrus, roses, and certain other sensitive crops when temperatures are high. Do not apply sulphur to such crops when the temperature is expected to rise likely to exceed 90 degrees F.

CAUTION KEEP OUT OF REACH OF CHILDREN

RECOMMENDATIONS

APPLES, PEARS: Powdery Mildew: Use 6 to 8 lbs. per 100 gals. Redenvelope to 2 lbs. if frost weather threatens.

CITRUS: Rust Mite: Use 10 to 16 lbs. per 100 gals. Begin application at first sign of damage and repeat as necessary. Rust mites should be controlled 3 or 4 times a year to prevent russetting of fruit and greasy spots of leaves.

PEACH: Brown Spots: Scab: Use 6 to 8 lbs. per 100 gals. in pink cover and on harvest sprays.

BEANS, PEAS: Powdery Mildew: Rust: Use 6 to 8 lbs. in sufficient water to spray. Repeat every 10 to 14 days as necessary.

IMPORTANT: Use higher dosage rates during periods of heavy rainfall or rapid growth. Keep new growth covered with protective coating.

DECONTAMINATION AND DISPOSAL INSTRUCTIONS: Do not re-use the container. Rinse with water, then contents and bury in sand or other waste at least eight inches deep in an isolated location away from water supplies. Burn the residue in a furnace completely. Exercise extreme caution and stay well away from the smoke.

NOTICE

Seller warrants that the product conforms to its chemical description and is suitable for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions of use making it unreasonable to expect the buyer assumes the risk of any such use.

Manufactured by

Superior Fertilizer & Chemical Co.

P. O. BOX 1021

TAMPA, FLORIDA