

DIRECTIONS FOR USE

IMPORTANT Read these entire Directions and Conditions of Sale before using DYRENE 50% Wettable Powder.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. FAILURE TO FOLLOW THESE DIRECTIONS OR THE OCCURRENCE OF WEATHER CONDITIONS THAT ARE EXTRAORDINARY OR UNUSUAL FOR THE USE AREA MAY RESULT IN UNSATISFACTORY PERFORMANCE OF THE PRODUCT. CHEMAGRO AND THE SELLER OFFER, AND THE BUYER ACCEPTS AND USES THIS PRODUCT SUBJECT TO THE AFOREMENTIONED CONDITIONS WHICH ARE BEYOND THE CONTROL OF CHEMAGRO AND THEREFORE THE RESPONSIBILITY OF THE BUYER.

DYRENE 50% Wettable Powder effectively protects crops and turf grasses from various destructive plant diseases. Complete and thorough coverage of foliage or turf areas is necessary for satisfactory control.

CROP APPLICATIONS - Begin applications when disease appears and repeat if needed at intervals of 7 to 10 days and during extended periods of wet weather every 4 or 5 days. Observe time limitations for each use for interval between last application and harvest.

TURF APPLICATIONS - For a preventive program begin application when diseases first appear and if needed, repeat at intervals of 10 to 14 days, during extended periods of moist conditions every 7 days.

Schedule applications in accordance with local conditions. Consult your State Agricultural Experiment Station or Extension Service for specific use information and for State regulations which may contain additional restrictions or requirements.

RESTRICTIONS

Do not exceed the above maximum dosage. Do not use on other crops used for food or feed. Use only according to label directions. Do not treat food crops grown in the greenhouse. Do not graze treated areas. Do not feed grass clippings to livestock.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

CONTAINER AND WASTE DISPOSAL

Rinse equipment and dispose of waste by burying in non-crop lands away from water supplies. Dispose of container by burying with wastes or burning (keep out of smoke). Never reuse container.

CAUTION

Harmful if swallowed or inhaled. Avoid contact with eyes, skin, and clothing. Avoid breathing of dust or spray mist. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

NOTE: DYRENE may cause a temporary allergic skin reaction in a few susceptible persons. This condition should be treated as an allergic dermatitis. There is no evidence of after effects or permanent injury. If sensitivity occurs, protective clothing should be worn during periods of exposure, such as, during application or when contacting treated foliage.

Do not contaminate feed or food. Keep out of reach of children.

EPA Est. 3125 MO 1 909032 579280

DYRENE is a Reg. TM of Mobay Chemical Corporation.



DYRENE

50% WETTABLE POWDER FOLIAGE FUNGICIDE

FOR CONTROL OF CERTAIN PLANT DISEASES ON CERTAIN CROPS AND TURF GRASSES

NET WEIGHT 50 POUNDS

STORE IN A COOL PLACE

CAUTION

KEEP OUT OF REACH OF CHILDREN

(See additional caution statement on sale panel)

ACTIVE INGREDIENT:	
4,6 Dichloro-N (2-chlorophenyl)-1,3,5 triazin-2-amine	50%
INERT INGREDIENTS:	
	50%
	100%

EPA Reg. No. 3125 50 AA

STOP — READ THE LABEL BEFORE USE

RECOMMENDED APPLICATIONS

CROP	DISEASE	Pounds DYRENE 50%	REMARKS
FIELD CROPS			
Tobacco	Brown spot (<i>Alternaria</i>)	2	Apply specified dosage weekly 100 gallons of water per acre. Apply at 10 day intervals after each harvest.
FRUIT			
Blackberries Blueberries Elderberries Raspberries	Cankers of bud and stem Leaf spot Gummy stem blight (<i>Alternaria</i>)	2 to 3	Apply specified dosage 100 gallons of water per acre. Do not apply within 14 days of harvest for the above crops.
Strawberries	Leaf spot Leaf blight	2 to 3	
Blueberries Huckleberries	Dieback spot Leaf blight	3	Apply specified dosage 100 gallons of water per acre for use. Do not apply within 14 days of harvest for use only in the above crops.
Cranberries	Berry spot Black rot End rot Phomopsis rot	5	Apply specified dosage 100 gallons of water per acre for use. Do not apply within 14 days of harvest for use only in the above crops.
VEGETABLES			
Celery	Early blight Late blight (<i>Botrytis</i>)	2 to 3	Apply specified dosage 100 gallons of water per acre. Do not apply within 14 days of harvest.
Cucumbers	Anthrax Downy mildew Gummy stem blight Leaf spots (<i>Alternaria</i>) (<i>Cercospora</i>)	4 to 6	Apply specified dosage 100 gallons of water per acre.
Onions (bulb or dry) Garlic	<i>Botrytis</i> Purple blotch (<i>Alternaria</i>)	2 to 3	Apply specified dosage 100 gallons of water per acre. Not recommended for use.
Onions (Green) Shallots	<i>Botrytis</i> Purple blotch (<i>Alternaria</i>)	2 to 3	Apply specified dosage 100 gallons of water per acre. Not recommended for use.
Potatoes	Early blight Late blight	2 to 5	Apply specified dosage 100 gallons of water per acre.
Pumpkins Squash	Anthrax Downy mildew Gummy stem blight Leaf spots (<i>Alternaria</i>) (<i>Cercospora</i>)	4 to 6	Apply specified dosage 100 gallons of water per acre.
Tomatoes	Anthrax <i>Botrytis</i> Early blight Grey leaf spot Late blight Septoria leaf spot	2 to 5	Apply specified dosage 100 gallons of water per acre.
MELONS			
Cantaloupes Honeydew Muskmelons Watermelons Other Varieties	Anthrax Downy mildew Gummy stem blight Leaf spots (<i>Alternaria</i>) (<i>Cercospora</i>)	4 to 6	Apply specified dosage 100 gallons of water per acre.
ORNAMENTALS			
Gladioli	Leaf spot (<i>Botrytis</i>)	2	Apply specified dosage 100 gallons of water per acre as a full coverage. Repeat weekly during period recommended for use.
Lawns & Turf (Bluegrasses, Bent grasses, Ryegrasses & Fescues)	<i>Helminthosporium</i> Leaf spots <i>Sclerotinia</i> Dollar spot	1/4 to 1/2	Apply specified dosage 100 gallons of water per 1,000 square feet at 1 to 2 week intervals during the spring and summer months.

*This dosage is calculated for conventional hydraulic type sprayers. When spray are applied per acre increase the concentration of DYRENE in the order to apply amount of DYRENE per acre, equivalent to a full coverage.

120 017

Chemagro
Agricultural Division
Mobay Chemical Corporation
P.O. Box 1000, St. Louis, MO 63103

120 017

DIRECTIONS FOR USE

Read these entire Directions and Conditions of Sale before using 50% Wettable Powder.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED BY SEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. FOLLOW THESE DIRECTIONS OR THE OCCURRENCE OF WEATHER THAT ARE EXTRAORDINARY OR UNUSUAL FOR THE USE AREA IN UNSATISFACTORY PERFORMANCE OF THE PRODUCT AND THE BUYER ACCEPTS AND USES IT SUBJECT TO THE FOREMENTIONED CONDITIONS WHICH ARE THE CONTROL OF CHEMAGRO AND THEREFORE THE RESPONSIBILITY OF THE BUYER.

50% Wettable Powder effectively protects crops and turf grasses from various plant diseases. Complete and thorough coverage of foliage or turf areas is necessary for satisfactory control.

APPLICATIONS: Begin applications when disease appears and repeat if needed at 10 to 14 days and during extended periods of wet weather every 4 or 5 days. Limitations for each use for interval between last application and harvest.

PREVENTIVE APPLICATIONS: For a preventive program begin application when diseases first appear, repeat at intervals of 10 to 14 days, during extended periods of wet weather every 7 days.

Applications in accordance with local conditions. Consult your State Experiment Station or Extension Service for specific use information and restrictions which may contain additional restrictions or requirements.

RESTRICTIONS

Do not exceed the above maximum dosage. Do not use on other crops used for food or forage according to label directions. Do not treat food crops grown in the same area as not graze treated areas. Do not feed grass clippings to livestock.

It is toxic to fish. Keep out of lakes, streams or ponds. Do not apply in weather conditions favor drift from areas treated. Apply this product as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

CONTAINER AND WASTE DISPOSAL

Remove and dispose of waste by burying in non-crop lands away from water supplies. Dispose of container by burying with wastes or burning (not in smoke). Never reuse container.

CAUTION

Do not swallow or inhale. Avoid contact with eyes, skin, and clothing. Avoid breathing of dust or spray mist. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

DYRENE may cause a temporary allergic skin reaction in a few susceptible persons. This condition should be treated as an allergic reaction. There is no evidence of after effects or permanent injury. If sensitivity occurs, protective clothing should be worn during periods of exposure, such as, during application or when contacting treated areas.

Do not contaminate feed or food. Keep out of reach of children.

EPA Est. 3125-MO-1 909032 579280

Reg. TM of Mobay Chemical Corporation.



DYRENE

50% WETTABLE POWDER FOLIAGE FUNGICIDE

FOR CONTROL OF CERTAIN PLANT DISEASES
ON CERTAIN CROPS AND TURF GRASSES

NET WEIGHT 50 POUNDS

STORE IN A COOL PLACE

CAUTION

KEEP OUT OF REACH
OF CHILDREN

(See additional caution statement on side panel)

ACTIVE INGREDIENT:	
4,6-Dichloro-N (2-chlorophenyl)-1,3,5-triazin-2-amine	50%
INERT INGREDIENTS:	
	50%
	100%

EPA Reg. No. 3125-50 AA

STOP — READ THE LABEL BEFORE USE

RECOMMENDED APPLICATIONS

CROP	DISEASE	Pounds DYRENE 50%	REMARKS
FIELD CROPS			
Tobacco	Brown rot (<i>Alternaria</i>)	2	Apply specified dosage using approxi- mately 100 gallons of water per acre for good coverage.* Apply at weekly intervals during harvest, after each picking.
FRUIT			
Blackberries Dwarfberries Loganberries Raspberries	Canine and leaf spot (<i>Ascochyta blight</i>) Leaf spot (<i>Ascochyta blight</i>)	2 to 4	Apply specified dosage using 100 to 200 gallons of water per acre for good coverage.* Do not apply within 5 days of harvest for strawberries and within 14 days of harvest for the other berries.
Strawberries	Berry mildew (<i>Ascochyta blight</i>) Leaf spot Leaf blight	2 to 4	
Blackberries Huckleberries	Double spot (<i>Septoria leaf spot</i>)	3	Apply specified dosage using 150 gallons of water per acre for good coverage.* Do not apply within 14 days of harvest. Restricted for use only in North Carolina.
Cranberries	Berry spot Black rot Foot rot Phytophthora rot	6	Apply specified dosage using 100 gallons of water per acre for good coverage.* Do not apply within 14 days of harvest. Restricted for use only in Wisconsin.
VEGETABLES			
Celery	Early blight Late blight (<i>Phytophthora</i>)	2 to 6	Apply specified dosage using 75 to 200 gallons of water per acre for good coverage.* Treated celery should be washed and bunched.
Cucumbers	Anthraxnose Downy mildew Gummy stem blight Leaf spots (<i>Alternaria</i>) (<i>Cercospora</i>)	4 to 6	Apply specified dosage using 100 to 200 gallons of water per acre for good coverage.*
Onions (bulb or dry) Garlic	<i>Botrytis</i> Purple blotch (<i>Alternaria</i>)	2 to 6	Apply specified dosage using 75 to 200 gallons of water per acre for good coverage.* Not recommended for use in Florida.
Onions (Green) Shallots	<i>Botrytis</i> Purple blotch (<i>Alternaria</i>)	2 to 3	Apply specified dosage using 75 to 100 gallons of water per acre for good coverage.* Not recommended for use in Florida.
Potatoes	Early blight Late blight	2 to 5	Apply specified dosage using 75 to 175 gallons of water per acre for good coverage.*
Pumpkins Squash	Anthraxnose Downy mildew Gummy stem blight Leaf spots (<i>Alternaria</i>) (<i>Cercospora</i>)	4 to 6	Apply specified dosage using 100 to 200 gallons of water per acre for good coverage.*
Tomatoes	Anthraxnose <i>Botrytis</i> Early blight Grey leaf spot Late blight <i>Septoria leaf spot</i>	2 to 5	Apply specified dosage using 75 to 175 gallons of water per acre for good coverage.*
MELONS			
Cantaloupes Honeydew Muskmelons Watermelons Other Varieties	Anthraxnose Downy mildew Gummy stem blight Leaf spots (<i>Alternaria</i>) (<i>Cercospora</i>)	4 to 6	Apply specified dosage using 100 to 200 gallons of water per acre for good coverage.*
ORNAMENTALS			
Glabadii	Leaf spot (<i>Ascochyta blight</i>)	2	Apply specified dosage in 100 gallons of water as a full coverage spray.* Repeat twice weekly during periods of infection. Recommended for use in Florida only.
Lawns & Turf (Bluegrasses, Bent grasses, Rye grasses, & Fescues)	<i>Helminthosporium</i> Leaf spots <i>Sclerotinia</i> Dollar spot	1/4 to 1/2	Apply specified dosage in 5 gallons of water per 1,000 square feet. Repeat at 1 to 2 week intervals throughout the spring and summer months.

*This dosage is calculated for conventional hydraulic type sprayers. When larger volumes of spray are applied per acre increase the concentration of DYRENE in the spray mixture in order to apply amount of DYRENE per acre equivalent to a full coverage spray.

120 017

Chemagro

Agricultural Division
Mobay Chemical Corporation
P.O. Box 1000, St. Louis, Mo. 63103

120 017