

U.S. LABEL

Reason to Issue: To change PHI for for strawberries, celery, onion and potatoes and to show maximum recommended rates per season.


Date of Draft: 9/23/86 (Pre-Reg) (T)

EPA Reg. No. 3125-50-AA

® DYRENE

50Z WETTABLE POWDER

FOLIAGE FUNGICIDE



ACTIVE INGREDIENT:
2,4-Dichloro-6-(o-chloroanilino)-s-triazine 50Z

INERT INGREDIENTS: 50Z
100Z

AMENDMENT
To Previously Registered Labeling

Revise as follows:

RECOMMENDED APPLICATIONS

DYRENE 50Z Wettable Powder effectively protects crops from various destructive plant diseases. Complete and thorough coverage of foliage is necessary for satisfactory control.

CROP APPLICATIONS - Apply DYRENE in a protective spray schedule before disease symptoms appear. Repeat if needed at intervals of 7 to 10 days, or every 4 to 5 days during extended periods of wet weather. Observe time limitations for each crop use for interval between last application and harvest.

Schedule applications in accordance with local conditions. Consult your State Agricultural Experiment Station or Extension Service for specific use information and for State regulations which may contain additional restrictions or requirements.

DIRECTIONS FOR USE

CROP	DISEASE	Pounds DYRENE 50Z WP	REMARKS
FRUIT			
Strawberries	Gray mold (Botrytis) Common leaf spot Leaf scorch	2 to 5	Apply specified dosage using 100 to 200 gallons of water per acre for good coverage.* Additional applications may be made as needed, but do not exceed 30 pounds of DYRENE 50Z WP per acre per season. Do not apply within 3 days of harvest.

DIRECTIONS FOR USE (Cont'd)

CROP	DISEASE	Pounds DYRENE 50Z WP	REMARKS
<u>VEGETABLES</u>			
Celery	Early blight Late blight Rhizoctonia crater rot	2 to 6	Apply specified dosage using 75 to 200 gallons of water per acre for good coverage by ground equipment* or a minimum of 5 gallons of water per acre by air. Additional applications may be made as needed, but do not exceed 60 pounds of DYRENE 50Z WP per acre per season. Do not apply within 7 days of harvest. Treated celery should be trimmed.
Onions (Green)	Botrytis leaf blight Purple blotch (<u>Alternaria</u>)	2 to 3	Apply specified dosage using 75 to 100 gallons of water per acre for good coverage.* Additional applications may be made as needed but do not exceed 18 pounds of DYRENE 50Z WP per acre per season. Do not apply within 3 days of harvest. Not recommended for use in Florida.
Potatoes	Early blight Late blight	2 to 5	Apply specified dosage using 75 to 175 gallons of water per acre for good coverage by ground equipment* or a minimum of 5 gallons of water per acre by air. Additional applications may be made as needed, but do not exceed 60 pounds of DYRENE 50Z WP per acre per season. Do not apply within 3 days of harvest.
Tomatoes	Anthrachnose Botrytis gray mold and ghost spot Early blight Gray leaf spot Late blight Septoria leaf spot	2 to 5	Apply specified dosage using 75 to 175 gallons of water per acre for good coverage by ground equipment* or a minimum of 5 gallons of water per acre by air. Additional applications may be made as needed, but do not exceed 60 pounds of DYRENE 50Z WP per acre per season. Do not apply within 3 days of harvest.

*This dosage is calculated for conventional hydraulic-type sprayers. When lower volumes of spray are applied per acre, increase the concentration of DYRENE in the spray mixture in order to apply the amount of DYRENE per acre, equivalent to a full coverage spray.