

3125-50

Directions for use.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not use on seed crops intended to be used for feed or forage
Do not graze treated areas

Do not reenter treated fields within 24 hours following application of this product

IMPORTANT: Read these entire Directions and Conditions of Sale before using DYRENE 50% Wettable Powder.

CONDITIONS OF SALE: The directions on this label were determined through research to be the directions for correct use of this product. This product has been tested for a range of weather conditions similar to those weather conditions that are ordinary and customary in the geographic area where the product is used. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather, or from failure to follow label directions. In addition, failure to follow label directions may cause injury to other crops, animals, man, or the environment. Mobay offers, and the buyer accepts and uses, this product subject to the conditions that extraordinary or unusual weather, or failure to follow label directions are beyond the control of Mobay and are, therefore, the responsibility of the buyer.

DYRENE 50% Wettable Powder effectively protects crops from various destructive plant diseases. Complete and thorough coverage of foliage is necessary for satisfactory control.

CROP APPLICATIONS: Apply DYRENE in a protective spray schedule before disease symptoms appear. Repeat if needed at intervals of 7 to 10 days, and during extended periods of wet weather every 4 or 5 days. Observe time limitations for each use for interval between last application and harvest.

Schedule applications in accordance with local conditions. Consult your State Agricultural Experiment Station or Extension Service for specific use information and for State regulations which may contain additional restrictions or requirements.

Do not apply this product through any type of irrigation system

Restrictions

Do not exceed the label maximum dosage. Do not use on seed crop intended to be used for feed or forage. Do not use on other crops used for food or feed. Use only according to label direction. Do not treat food crops grown in the greenhouse. Do not graze treated areas

Re-entry statements

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not reenter treated fields within 24 hours following application of this product. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings should include those statements found under the label heading "Precautionary Statements" and "Re-entry Statements" included on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: **WARNING,** area treated with DYRENE 50% Wettable Powder on (date of application) Do not reenter treated fields within 24 hours following application of this product. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If swallowed, drink large quantities of water.

DYRENE

Dyrene® 50% Wettable Powder Foliage Fungicide for control of certain plant diseases on certain crops.

ACTIVE INGREDIENT:

4, 6-Dichloro-N(2-chlorophenyl)

- 1, 3, 5-triazin-2-amine

50%

INERT INGREDIENTS:

50%

100%

EPA Reg. No. 3125-50

EPA Est. 3125-MO-1

Store in a cool place.

STOP—Read the label before use.

Keep out of reach of children.

DANGER

(See side panel for Statements of Practical Treatment and Other Precautionary Statements)

PELIGRO

PRECAUCION AL USUARIO: Si usted no puede leer o entender Inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

NET WEIGHT 4 POUNDS

Mobay Corporation

A Bayer U.S.A. INC. COMPANY



Agricultural Chemicals Division
Box 4913, Kansas City, Mo. 64120

Storage and disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Do not use container in connection with food or drinking water. Completely empty bag into applicator or other container. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, out of smoke.

Store in a cool, dry place and in such a manner as to prevent contamination with other pesticides, fertilizers, food, and feeds. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason, do not use. Carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed above. In spill or leak incidents, keep unauthorized people away. For more information, may contact the Mobay Emergency Response Team for decontamination procedures or any other assistance that may be needed. The Mobay Kansas City Emergency Response Telephone Number is 242-2582, or contact Chemtrec at 800-424-9300.

Precautionary statements

Hazards to humans and domestic animals

DANGER

Causes irreversible eye damage. Harmful if swallowed. Do not get into eyes or on clothing. Avoid contact with skin. Wear goggles, shield or safety glasses. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing with hot water before reuse.

NOTE: Protective clothing should be worn during periods of use, such as during application, or when contacting foliage. DYRENE anilazine may cause an allergic skin reaction in some susceptible persons. This condition should be treated as an allergic dermatitis.

Keep out of reach of children.

Statements of practical treatment

If in eyes: Hold eyelids open and flush with steady, gentle stream of water for 15 minutes. Get medical attention. **If on skin:** Wash immediately with soap and warm water. If irritation occurs, get medical attention. **If swallowed,** drink promptly a large quantity of milk. If milk is not available, large quantities of water. Ask doctor for medical attention. Probable mucosal damage may contraindicate the use of gastric lavage.

Environmental hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or wetlands. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Cover or incorporate spills.

905400

DYRENE is a Reg. TM of Mobay Corporation



Recommended applications.

CROP	DISEASE	Pounds DYRENE 50% WP	REMARKS
FRUIT Strawberries	Gray mold (Botrytis) Common leaf spot Leaf scorch	2 to 6	Apply specified dosage using 100 to 200 gallons of water per acre for good coverage. Do not apply within 5 days of harvest.
VEGETABLES Celery	Early blight Late blight Rhizoctonia crater rot	2 to 6	Apply specified dosage using 75 to 200 gallons of water per acre for good coverage. Treated celery should be washed and trimmed.
Onions (Green)	Botrytis leaf blight Purple blotch (Alternaria)	2 to 3	Apply specified dosage using 75 to 100 gallons of water per acre for good coverage. Not recommended for use in Florida.
Potatoes	Early blight Late blight	2 to 5	Apply specified dosage using 75 to 175 gallons of water per acre for good coverage.
Tomatoes	Anthraxnose Botrytis gray mold and ghost spot Early blight Gray leaf spot Late blight Septoria leaf spot	2 to 5	Apply specified dosage using 75 to 175 gallons of water per acre for good coverage.

*This dosage is calculated for conventional hydraulic-type sprayers. When lower volumes of spray are applied per acre increase the concentration of DYRENE in the spray mixture in order to apply amount of DYRENE per acre, equivalent to a full coverage spray.

ACCEPTED
MAR 7 1988
Under the Federal Insecticide,
Fungicide and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg No. 3125-50