

U.S. LABEL

Reason for Issue: To add use on cucumber and summer squash.

Date of Draft: 03/01/91 (Pre-Reg)(T)

Supersedes Pre-Reg Draft Date: 01/23/91(T)

EPA Reg. No. 3125-50

**DYRENE**

50% WETTABLE POWDER

FOLIAGE FUNGICIDE

**ACCEPTED**

**JAN 16 1992**

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 3125-50

**ACTIVE INGREDIENT:**

4,6-Dichloro-N-(2-chlorophenyl)-1,3,5-triazin-2-amine ..... 50%

INERT INGREDIENTS: ..... 50%  
100%

**AMENDMENT TO PREVIOUSLY REGISTERED LABELING**

REVISE AS FOLLOWS:

**RECOMMENDED APPLICATIONS**

CROP	DISEASE	POUNDS DYRENE 50% WP	REMARKS
Cucumber Squash (Summer)	Anthracnose Downy mildew Gummy stem blight Leaf spots ( <u>Alternaria</u> ) ( <u>Cercospora</u> )	2 to 4	Apply specified dosage using 100 to 200 gallons of water per acre for good coverage.*  Apply DYRENE in a protective spray schedule before disease symptoms appear. Make additional applications at 7- to 10-day intervals, or every 4 to 5 days during extended periods of wet weather. Use the higher label rate under heavy disease pressure and/or on highly disease susceptible varieties.  Do not exceed 12 pounds of DYRENE 50% WP per acre per season. Do not apply within 7 days of harvest.

\*This dosage is calculated for conventional hydraulic-type sprayers. When lower volumes of spray are applied per acre, increase the concentration of DYRENE in the spray mixture in order to apply amount of DYRENE per acre equivalent to a full coverage spray.