U.S. LABEL

Reason to Issue: To add use on potatoes as a pre-plant broadcast application and allow for a post-emergence sidedress following broadcast.

USDA Reg. No. 3125-58

Date of Draft: 8/8/68 (Pre-Rek.)

® DI-SYSTON

5Z GRANULAR

Systemic Insecticide

AMENDMENT
To Previously Registered Label

ACCEPTED

UNDER THE PERSONAL BENEFICIDE FUNGICIDE AND ROBERTICIDE ACT FOR ECCAO-AC POROS BENEFICE DE UNDER NO.3 / C ATTACHED COMMENTS.

Amend as follows:

RECOMMENDED DI-SYSTON APPLICATIONS

CROP	INSECT	DOSAGE DI-SYSTON 5Z	remarks
Potatoes	Aphids Flea Beetles (except tuber flea beetles) Leafhoppers Potato Psyllid Southern Potato Wireworm Colorado Potato Beetle (early season reduction)	46 to 70 ounces per 1000 feet of row (for any row spacing*) or 40 to 60 pounds per acre (38 inch row spacing)	Place the granules in the seed furrow or in a band on each side of the seed furrow at planting. An application may also be made as a side-dressing after plants become established. Do not apply within 75 days of harvest. Higher rate to be used in organic soils.
		60 to 80 pounds per acre	BROADCAST APPLICATION Apply specified dosage per acre as a pre-plant broadcast application. York granules into the soil to a depth of 2 to 3 inches using a disc or other suitable equipment. Plant crop in usual manner. A post-emergence sidedress may also be made at rates of 46 to 70 ounces per 1000 feet of row (for any row spacing*), after plants become established. Do not make more than two applications per season nor apply within 75 days of harvest.

ALL LABEL TEXT ABOVE, EXCEPT THAT WHICH IS UNDERLINED, HAS BEEN PREVIOUSLY ACCEPTED FOR REGISTRATION.

Above amendment is the original submitted with application.

BEST AVAILABLE COPY