

U.S. LABEL

**Reason to Issue:** To add the "application at planting" use for control of aphids in sorghum.

Date of Draft: 2/23/70 (Pre-Reg.) (S)

USDA Reg. No. 3125-61

®  
DI-SYSTON

10% Granular

SYSTEMIC INSECTICIDE

AMENDMENT

To Previously Registered Labeling

REVISE AS FOLLOWS:

RECOMMENDED APPLICATIONS

CROP	PEST	Dosage DI-SYSTON 10% Granular	REMARKS
Sorghum	Aphids (Greenbugs)	9 to 12 ozs. per 1000 ft. of row (for any row spacing*) or 7 1/2 to 10 pounds per acre (40-inch row spacing)	<u>APPLICATION AT PLANTING: IN-FURROW</u> Place the granules in the seed furrow approximately 1/2 inch above the seed at planting. Do not apply directly on the seed. See "NOTE" below.
		12 ozs. per 1,000 ft. of row (for any new spacing*) or 10 pounds per acre (40-inch row spacing)	<u>APPLICATION AT PLANTING: BAND</u> Place the granules in a 4 to 6 inch band directly behind the planter shoe in front of the press wheel. Do not apply directly on the seed. See "NOTE" below.  <u>POST-PLANTING APPLICATION: WHORL</u> Apply specified dosage by broadcasting into the whorl. Do not apply within 30 days of harvest of grain or 14 days of harvest for forage on fodder. See "NOTE" below.
NOTE: Do not apply DI-SYSTON 10% Granular to sorghum more than once per crop season regardless of method of application.			

\*ON ROW CROPS WHERE ROW SPACING IS EXTREMELY NARROW, DO NOT PLACE TREATED ZONES (BAND OR FURROW) CLOSER TOGETHER THAN 6 INCHES.

**ACCEPTED**  
  
**JUL 2 1970**  
  
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 3125-61