

U.S. LABEL

Reason to Issue: To recommend 3 post-planting applications into the sorghum whorl following an at-planting application.

Date of Draft: 6/7/72 (Pre-Reg.) (S)

EPA Reg. No. 3125-61

ACCEPTED
SEP 1 1972
FEDERAL INSECTICIDE AND FUNGICIDE ACT
EPA REGISTRATION NO. 3125-61

DI-SYSTON
10% Granular

SYSTEMIC INSECTICIDE

AMENDMENT

To Previously Registered Labeling

RECOMMENDED APPLICATIONS

CROP	INSECT	DOSAGE DI-SYSTON 10% Granular	REMARKS
Sorghum	Aphids (Greenbugs)	9 to 12 ozs. per 1000 ft. of row (for any row spacing*) -or- 7-1/2 to 10 pounds per acre (40-inch row spacing)	<u>APPLICATION AT PLANTING: IN-FURROW</u> Place the granules in the seed furrow approximately 1/2 inch above the seed at planting. Do not apply directly on the seed. See "NOTE" below.
		12 ozs. per 1000 ft. of row (for any row spacing*) or 10 pounds per acre (40-inch row spacing)	<u>APPLICATION AT PLANTING: BAND</u> Place the granules in a 4 to 6 inch band directly behind the planter shoe in front of the press wheel. Do not apply directly on the seed. See "NOTE" below. <u>POST-PLANTING APPLICATION: WHORL</u> Apply specified dosage by broadcasting into the whorl. Do not apply more than 3 times per season nor within 30 days of harvest of grain or 14 days of harvest for forage or fodder. See "NOTE" below.
		<u>NOTE:</u> Post-planting broadcast applications into the whorl may be made following an at-planting application.	

*ON ROW CROPS WHERE ROW SPACING IS EXTREMELY NARROW, DO NOT PLACE TREATED ZONES (BAND OR FURROW) CLOSER TOGETHER THAN 6 INCHES.