



# DYRENE

## 5% DUST

FOLIAGE FUNGICIDE

FOR CONTROL OF PLANT DISEASES

GUARANTEE

ACTIVE INGREDIENT	
2,4-Dichloro-6-(o-chloroanilino)-s-triazine	5%
INERT INGREDIENTS	95%
	100%

U.S. Patent No. 2,720,480  
 Canadian Patent No. 546,027  
 EPA Reg. No. 3125-62

STOP READ THE LABEL BEFORE USE

### CAUTION

KEEP OUT OF REACH OF CHILDREN

(See side panel for caution statements)

NET CONTENTS                      POUNDS

## CHEMAGRO

Division Of Bayer Corporation  
 Box 4913, Kansas City, Mo. 64120

DIRECTIONS FOR USE

**IMPORTANT:** Read these entire Directions and Conditions of Sale including the Warranty and Limitation of Damages provision before using DYRENE 5 Dust.

**CONDITIONS OF SALE:** These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller, Chemagro and the Seller offer, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

**DYRENE 5** Dust foliage fungicide effectively protects crops from various destructive plant diseases including early and late blights. Complete and thorough coverage of foliage is necessary for satisfactory control. Begin applications before disease appears and repeat treatments as needed, usually at intervals of 7 to 10 days, except during extended periods of cool, wet weather when applications every 4 or 5 days may be advisable. The DYRENE dust program should be continued throughout the growing season. Schedule applications in accordance with local conditions. Consult your State Agricultural Experiment Station or Extension Service for specific use information.

RESTRICTIONS

Do not exceed the recommended maximum dosage. Under severe conditions of late blight on tomatoes and potatoes, consult State Agricultural Experiment Station or Extension Service for specific use information. Do not use on other crops used for food or forage.

This product is toxic to fish. Keep out of any body of water. Do not apply under conditions that favor drift.

CONTAINER AND WASTE DISPOSAL

Rinse equipment and dispose of waste by burying in non-crop lands away from water supplies. Dispose of container by burying with wastes or burning (keep out of smoke). Never reuse container.

CAUTION

Harmful if swallowed or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing of dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

Do not contaminate feed or food. Keep out of reach of children.

WARRANTY AND LIMITATIONS OF DAMAGES

Chemagro warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Direction for Use and Conditions of Sale, subject to the risks referred to therein. CHEMAGRO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, and no agent of Chemagro is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential or commercial damages such as loss of profits or value of crop.

RECOMMENDED APPLICATIONS

CROP	DISEASE	Period of DYRENE 5	REMARKS
<b>FRUIT</b>	Cane and leaf rust		
Blackberries Dewberries Raspberries Loganberries	<i>(Kuehneella) rot</i> <i>Botrytis</i> fruit rot Cane and leaf spot <i>(Septoria rubi)</i>	30 to 60	Apply specified dosage per acre. Do not apply within 14 days of harvest.
Strawberries	Gray mold <i>(Botrytis)</i> Leaf spot Leaf scorch	20 to 40	Apply specified dosage per acre. Do not apply within 14 days of harvest.
<b>VEGETABLES</b>			
Celery	Early blight Late blight <i>Rhizoctonia</i>	20 to 40	Apply specified dosage per acre. Treated celery should be washed and trimmed.
Cucumbers	Anthracnose Downy mildew Gummy stem blight Leaf spots <i>(Alternaria)</i> <i>(Cercospora)</i>	30 to 60	Apply specified dosage per acre.
Onions (Bulb or dry) Garlic	<i>Botrytis</i> Purple blotch <i>(Alternaria)</i>	20 to 40	Apply specified dosage per acre. Not recommended for use in Florida.
Onions (green) Shallots	<i>Botrytis</i> Purple blotch <i>(Alternaria)</i>	20 to 40	Apply specified dosage per acre. Not recommended for use in Florida.
Potatoes	Early blight Late blight	20 to 40	Apply specified dosage per acre.
Pumpkins Squash	Anthracnose Downy mildew Gummy stem blight Leaf spots <i>(Alternaria)</i> <i>(Cercospora)</i>	30 to 60	Apply specified dosage per acre.
Tomatoes	Anthracnose <i>Botrytis</i> Early blight Gray leaf spot Late blight <i>Septoria</i> leaf spot	20 to 30	Apply specified dosage per acre.
<b>MELONS</b>			
Honeydew Muskmelons Watermelons Other Varieties	Anthracnose Downy mildew Gummy stem blight Leaf spots <i>(Alternaria)</i> <i>(Cercospora)</i>	30 to 60	Apply specified dosage per acre.