

U.S. LABEL-(S)

Reason to Issue: To propose tolerances for soybeans, walnuts, dry beans, blackeyed peas, and potatoes currently registered on a "no-residue" basis.

Date of Draft: 7/31/69 (L)

USDA Reg. No. 3125-63

®
GUTHION

3% Dust
Crop Insecticide

Currently Registered Text

ACCEPTED

JUN 16 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 3125-63

RECOMMENDED GUTHION APPLICATIONS

CROP	INSECT	Pounds GUTHION 3%	REMARKS
<u>FIELD CROPS</u>			
Soybeans	Aphids Bean leaf beetle Green cloverworm Leafhoppers Leafminers Leaf rollers Stink bugs Velvet bean caterpillar	15 to 20	Apply specified dosage per acre.* Repeat as necessary. Do not apply within 45 days of harvest. Do not graze or feed treated vines to livestock.
	Mexican bean beetle	20 to 30	
<u>NUTS</u>			
Walnuts	Aphids Codling moth European red mite	40 to 50	Apply specified dosage per acre.* Repeat as necessary. Do not apply after husks split. Do not graze livestock in treated groves for 21 days after treatment.
<u>VEGETABLES</u>			
Beans (Snap & Dried)	Aphids Bean leaf beetle Green cloverworm Leafhoppers Leaf miners Leaf rollers Mites Stink bugs Velvet bean caterpillar	15 to 20	Apply specified dosage per acre.* Repeat as necessary. Do not apply to snap beans within 7 days of harvest, nor to dry beans within 30 days of harvest. Do not exceed 4 applications on dry beans. Do not feed or ensile treated forage.
	Mexican bean beetle Spotted cucumber beetle Striped cucumber beetle Tarnished plant bug Western-striped cucumber beetle	20	

U.S. LABEL-(S)

Reason to Issue: To propose tolerances for soybeans, walnuts, dry beans, blackeyed peas, and potatoes currently registered on a "no-residue" basis.

Date of Draft: 7/31/69 (L)

USDA Reg. No. 3125-63

GUTHION
3% Dust
Crop Insecticide
Currently Registered Text

ACCEPTED

JUN 16 1972

RECOMMENDED GUTHION APPLICATIONS

CROP	INSECT	Pounds GUTHION 3%	REMARKS
<u>FIELD CROPS</u>			
Soybeans	Aphids Bean leaf beetle Green cloverworm Leafhoppers Leafminers Leaf rollers Stink bugs Velvet bean caterpillar	15 to 20	Apply specified dosage per acre.* Repeat as necessary. Do not apply within 45 days of harvest. Do not graze or feed treated vines to livestock.
	Mexican bean beetle	20 to 30	
<u>NUTS</u>			
Walnuts	Aphids Codling moth European red mite	40 to 50	Apply specified dosage per acre.* Repeat as necessary. Do not apply after husks split. Do not graze livestock in treated groves for 21 days after treatment.
<u>VEGETABLES</u>			
Beans (Snap & Dried)	Aphids Bean leaf beetle Green cloverworm Leafhoppers Leaf miners Leaf rollers Mites Stink bugs Velvet bean caterpillar	15 to 20	Apply specified dosage per acre.* Repeat as necessary. Do not apply to snap beans within 7 days of harvest, nor to dry beans within 30 days of harvest. Do not exceed 4 applications on dry beans. Do not feed or ensile treated forage.
	Mexican bean beetle Spotted cucumber beetle Striped cucumber beetle Tarnished plant bug Western-striped cucumber beetle	20	