C-9302

U.S. LABEL-(S)

Base Reg. (+545)

Reason to Issue: To update in line with PCNB RPAR notice.

Date of Draft: 3/4/82 (Fre-Reg)() Supersedes Draft Dated: 16/16/72

DEXON - TERRACLOR®

35 - 35 WETTABLE POWDER

SEED TREATMENT FUNGICIDE

FOR CONTROL OF PLANT DISEASES

NET WEIGHT

POUNDS

WARNING

KEEP OUT OF REACH OF CHILDREN

(See rear panel for warning statements)

ACTIVE INGREDIENTS:

ACCEPTED

ART 5 1982

EPA Reg. No. 3125-70

STOP - READ THE LABEL BEFORE USE

MOBAY CHEMICAL CORPORATION AGRICULTURAL CHEMICALS DIVISION Box 4913, Kansas City, Mo. 64120

Note: Underlined letter in chemical nomenclature should be italicized when printed, not underlined.

Page 1 of 3



DEMON-TERRACLOR 35-25 WP

DIRECTIONS FOR USE

IMPORTANT: Read these entire Directions and Conditions of Sale, before using DEXON-TERRACLOR 35-35 Wettable Powder.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE THE DIRECTIONS FOR CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED FOR A RANGE OF WEATHER CONDITIONS SIMILAR TO THOSE WEATHER CONDITIONS THAT ARE ORDINARY AND CUSTOMARY IN THE GEOGRAPHIC AREA WHERE THE PRODUCT IS USED. INSUFFICIENT CONTROL OF PESTS MAY RESULT FROM THE OCCURRENCE OF EXTRADRDINARY OR UNUSUAL WEATHER, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO THE CROP TO WHICH THE PRODUCT IS APPLIED, OTHER CROPS, ANIMALS, MAN, OR THE ENVIRONMENT. MOBAY AND THE SELLER OFFER, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL WEATHER, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF MOBAY AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

DEXON-TERRACLOR 35-35 Wettable Powder is recommended as a seed treatment to provide effective control of Pythium, Aphanomyces, and Rhizoctonia which cause damping-off of sugar beet seedlings. The formulation wets readily with water to form a slurry suitable for use in any commercial slurry seed treater. DEXON-TERRACLOR is compatible with all commercial insecticides recommended for seed and soil treatment and may be applied in combination with them.

RECOMMENDED APPLICATIONS

C ROP	DISEASE	Ounces DEXON- TERRACLOR 35-35	REMARKS
Sugar beets	Pythium Aphanomyces Rhizoctonia	4 to 8	Apply specified dosage to 100 pounds of seed as a slurry treatment in sufficient water to provide uniform coating of the seed.

Note: To meet U.S. Federal Seed Act requirements, all seed treated with DEXON should be labeled: "Treated with Sodium[4-(dimethylamino)phenyl] diazinesulfonate (DEXON) and P.C.N.B. (TERRACLOR)".

RESTRICTIONS

Treated seed must not be used for feed, food, or oil purposes. Use only according to label directions. Applications at rates above those shown may result in illegal crop residues.

This product is Loxic to fish and wildlife. Keep out of lakes, streams, or ponds. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife.

CONTAINER AND WASTE DISPOSAL

Rinse equipment and dispose of waste by burying in non-crop lands away from water supplies. Dispose of container by burying with wastes or by burning (keep out of smoke). Never reuse container.

WARNING

May be poisonous if swallowed or inhaled. Do not get in eyes or on skin. Wear natural or synthetic rubber gloves, a mask suitable for use against toxic dusts, and goggles when treating seed or loading seeder. Do not breathe dust. May cause skin irritation.

page 2 of 3

DEXON-TERRACLOR 35-35 WP

WARNING (Cont.)

Do not contaminate feed or food. Keep out of reach of children. Wash hands, arms, and face thoroughly with soap and warm water after use and before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

DEXON is a Reg. TM of the Parent Company of Farbenfabriken Bayer GmbH, Leverkusen.

Terraclor is a Reg. TM of Olin Mathieson Chemical Corporation

TROCK TOTAL