

3125-102

02/15/1973


1/1

U.S. LABEL

Reason to Issue: To add use in Texas
on sugarcane.

Date of Draft: 11/6/72 (Pre-Reg.) (S)

EPA Reg. No. 3125-102-AA

GUTHION 2L
Emulsifiable InsecticideAMENDMENT
To Previously Registered Labeling

AMEND AS FOLLOWS:

RECOMMENDED APPLICATIONS

CROP	INSECT	Pints GUTHION 2L	REMARKS
Sugarcane (Conventional Spray) (Florida, Louisiana, and Texas only)	Sugarcane borer	3	Apply specified dosage per acre by air application using a minimum of 2 gallons of water per acre. For best results applications should be made at approximately cane top level and not more than 10 feet above cane top level. Do not apply more than 5 times per season nor within 30 days of harvest. Bagasse from sugarcane treated with GUTHION may be used for feed of cattle, goats, and sheep. Consult you State Agricultural Extension Service or Experiment Station for specific use information.
Sugarcane(1) (Ultra Low Volume Spray) (Florida, Louisiana, and Texas only)	Sugarcane borer	3	Apply specified dosage per acre undiluted GUTHION 2L may be used undiluted in any aerial spray equipment that has been adapted and calibrated for ultra low volume spraying. Planes must be equipped with accept low volume devices that will produce droplets within the range of 30 to 100 micron in size. For best results applications should be made at approximately cane top level and not more than 10 feet above cane top level. Do not apply more than 5 times per season nor within 30 days of harvest. Bagasse from sugarcane treated with GUTHION may be used for feed of cattle, sheep and goats. Consult your State Agricultural Extension Service or Experiment Station for specific use information.

(1)

Note: This formulation, when used undiluted, may cause spotting of automobile finish if prolonged exposure is permitted. Do not spray directly over automobiles. If a dental exposure does occur, automobiles should be washed immediately.