1/1

U.S. LABEL

Reason to Issue: To expand label to include use on all varieties of pine for coneworm and seedworm control; 5/9/75 Draft: To revise crop designation, rates, and "Remarks".

Date of Draft: 5/9/75 (Pre-Reg.) (S)
Supersedes Pre-Reg. Draft Dated:
2/18/75

EPA Reg. No. 3125-102

JUN 26 1975

Under the Federal Insecticide, The Fungicide, and Historicide Actions a amended, for the pesticide as

R GUTHION 2L

CROP INSECTICIDE

AMENDMENT
To Prevously Registered Labeling

## AMEND AS FOLLOWS:

registered under EPA Reg. No.

CROP	INSECT	Pints GUTHION 2L	REMARKS
Southern Pine Seed Orchards	Coneworm Seedworm	See Remarks	Use 6 pints per 100 gallons of water (0.2% dilution) for high volume sprayers of 3 pints per 10 gallons of water (1% dilution) for low volume sprayers. Apply first application within 30 days following conelet closure, followed by 3 to 5 applications (at least 30 days apart). Apply approximately 5 to 10 gallons of the 0.2% dilution with high volume sprayers or approximately 1 to 2 gallons of the 1% dilution with low volume sprayers per tree. Thorough coverage of cones is necessary for maximum control.