

PM-12

3125-100

p. 1/1

YLD27

U.S. LABEL

Reason to Issue: To redefine cabbage
to include tight heading varieties
of Chinese cabbage.

Date of Draft: 10/2/85 (Pre-Reg) (G)

EPA Reg. No. 3125-102

® GUTHION 2L

Emulsifiable Insecticide

ACTIVE INGREDIENT:

O,O-Dimethyl S-[(4-oxo-1,2,3-benzo-
triazin-3(4H)-yl)methyl] phosphorodithioate 22.2%
Aromatic Petroleum Distillate 59.8%

INERT INGREDIENTS: 18.0%
100.0%

AMENDMENT TO
Previously Registered Labeling

ACCEPTED
JAN 24 1986

Revise as Follows:

RECOMMENDED APPLICATIONS

CROP	INSECT	Pints GUTHION 2L	REMARKS
VEGETABLES Cabbage (includes tight heading varieties of chinese cabbage)	Aphids Cabbage looper Diamondback moth Imported cabbageworm	2 to 3	Apply specified dosage per acre in sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary. Do not apply within 7 days of harvest for Brussels sprouts, 15 days of harvest for broccoli and cauliflower, nor within 21 days of harvest for cabbage.
	Cabbage maggot	1/2	Mix specified dosage in 50 gallons of water. Apply 4 to 6 ounces of this emulsion per plant immediately after transplanting.
	Cabbage maggot (Transplant Fields in California only)	3	Apply specified dosage in 300 to 400 gallons of water per acre as a soil drench in the rows when damage first appears. Additional applications may be necessary. Do not apply within 21 days of harvest.
	Cabbage maggot (Direct Seeded Fields in California only)	3	Apply specified dosage per acre in sufficient water for uniform distribution. Mix in the upper 2 inches of soil prior to seeding or spray in the seed row at planting time. Usually 2 to 3 additional sprays are necessary during the growing season depending upon time of year and maggot population. Do not apply within 21 days of harvest.