

Best Available Copy

3125-111

11-9-1976

11

ACCEPTED
NOV 9 1976
UNDER NO. 3125-111
FUNGICIDE AND INSECTICIDE ACT
FOR ECONOMIC REGISTER

Metasystox-R Spray Concentrate
EPA Reg. No. 3125-111-A4

CROP	INSECT	INSECT	REMARKS
FIELD CROPS	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.
VEGETABLES	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.
FRUIT TREES	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.

RECOMMENDED APPLICATIONS

CROP	INSECT	INSECT	REMARKS
FIELD CROPS	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.
VEGETABLES	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.
FRUIT TREES	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.

RECOMMENDED APPLICATIONS

CROP	INSECT	INSECT	REMARKS
FIELD CROPS	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.
VEGETABLES	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.
FRUIT TREES	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.

DIRECTIONS FOR USE

Metasystox-R is a systemic insecticide for control of sucking insects on a wide range of crops. It is applied as a soil drench or as a foliar spray. The active ingredient is dimethoate, which is absorbed by the plant and moves to the feeding site of the insect. Insects that feed on the plant ingest the insecticide and die. Metasystox-R is effective against a wide range of insects, including aphids, beet beetles, and root-knot. It is also effective against some mites and nematodes. Metasystox-R is registered for use on a wide range of crops, including corn, soybeans, cotton, and various fruits and vegetables. It is also registered for use on ornamental plants and lawns. Metasystox-R is a highly effective insecticide that provides long-term control of sucking insects. It is easy to apply and does not require special equipment. Metasystox-R is a valuable tool for the farmer and the gardener alike.

111113



METASYSTOX-R

SPRAY CONCENTRATE

SYSTEMIC INSECTICIDE

FOR EFFECTIVE SYSTEMIC CROP PROTECTION

NET CONTENTS 2 U.S. GALS.
WARNING
KEEP OUT OF REACH OF CHILDREN
(See rear panel for handling instructions)

WARNING - FLAMMABLE! KEEP AWAY FROM HEAT AND OPEN FLAME
STOP - READ THE LABEL BEFORE USE

Chemagro
Agricultural Division
Bayer Corporation

RECOMMENDED APPLICATIONS

CROP	INSECT	INSECT	REMARKS
FIELD CROPS	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.
VEGETABLES	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.
FRUIT TREES	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.

RECOMMENDED APPLICATIONS

CROP	INSECT	INSECT	REMARKS
FIELD CROPS	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.
VEGETABLES	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.
FRUIT TREES	Root-knot	1%	Apply 10-15 days before harvest. Do not apply more than 1.0 gals. per acre.

111113