

E-300
Ref. 3

1968-1969 (600)

For use on roses, pyracantha, spirea, birch, and other flowers and shrubs
from damage by aphids, spider mites, thrips, leaf miners,
whiteflies, lace bugs, leafhoppers, and certain other pests.

ACCEPTED
NOV 8 1968
UNDER THE FEDERAL INSECTICIDE
AND FUNGICIDE ACT
ECONOMIC POISON REG. ST.
NO. 572-116

FOR ROSES, SPIREA, BIRCH, AND OTHER
FLOWERS AND SHRUBS

Protects plants, such as roses, pyracantha, spirea, birch, and other flowers and shrubs from damage by aphids, spider mites, thrips, leaf miners, whiteflies, lace bugs, leafhoppers, and certain other pests.

WARNING
KEEP OUT OF REACH OF CHILDREN
(See side panel for warning statements)

ACTIVE INGREDIENTS:
2,2-Diethyl 2-[2-(ethylthio)ethyl] phosphorothioate 4.25
INERT INGREDIENTS: 100
100
U.S. Patent No. 2,795,610 Canadian Patent No. 518,967

EPA Reg. No. 3209-116

How to Use: Spray thoroughly on the undersides of leaves and on the stems of plants. Repeat every 7-10 days. Do not spray on plants that are stressed or in bloom. Do not spray on plants that are in contact with water. Do not spray on plants that are in contact with children or animals. Do not spray on plants that are in contact with food or feed.

Note: Underline letters in boldface and should be underlined when printed, not underlined.

DI-SYSTON Granules (3 1/2 lb.)

HOW TO USE:

Apply the ready-to-use granules uniformly (one lb. per 100 sq. ft. of application) to the soil around the base of the plants. Use the granules in the soil, then water. The chemical, released from the granules, will be carried into the soil and will be taken into the plants. The granules are also effective against insects on the soil surface. Use the top soil before sowing seed or setting out plants. Insect control applications of DI-SYSTON Systemic Insecticide can be made as needed to maintain insect control after the plants start to grow.

NOTE: THE CAP OF THIS CONTAINER, WHEN FULL, HOLDS APPROXIMATELY ONE ROUNDED TEASPOONFUL OF DI-SYSTON GRANULAR. *(Printer see below).

RECOMMENDED PLANTS TO USE

FOR FLOWERS

- Ageratum
- Calendula
- Carnation
- Chrysanthemum
- Delphinium
- Petunia
- & other flowers

TO CONTROL

- Aphids (plant lice)
- Lace bug
- Leafhoppers
- Spider mites
- Trips
- Whiteflies

BRANCH PLANTING:

Apply 10 ounces evenly over 200 square feet (10 X 20 ft.) of bed area and work into the top 2 to 3 inches of soil. Sow seed or set plants and water thoroughly.

GROWING PLANTS:

New Newly Set Plants: Apply one rounded teaspoonful around base of plant, work into soil and water thoroughly.


New Plants in Row: Apply one rounded teaspoonful per foot of row in a narrow band near base of plants. Work into top soil and water thoroughly. Repeat application at 4- to 6-week intervals if insect damage occurs.

Established Plants:

Apply one rounded teaspoonful per 6-inch pot. Mix thoroughly in to soil and water with water.

Shrub Planting:

Apply 2 to 4 rounded teaspoonfuls per foot of shrub height. Use the lower rate for shrubs up to 3 ft. tall. The higher rate for shrubs 3 to 6 ft. tall. For shrubs over 6 ft. in height, use 1/2 lb. of DI-SYSTON per 100 sq. ft. of ground. Apply in all directions. Distribute the granules around evenly from the base of shrub to the drip line on all sides of the shrub. Water thoroughly. Repeat application at 4- to 6-week intervals if required.



FOR SHRUBS

- Azalea
- Birch
- Camellia
- Fuquias
- Moraine locust
- Pine
- Phododendron
- Roses
- Yew
- & other shrubs

TO CONTROL

- Aphids (plant lice)
- Bark leaf miner
- Curculionid beetle
- Leafhopper
- Lace bug
- Leafhoppers
- Phytophagous
- Pine saw fly
- Salt cedar (gambel)
- Whiteflies

Recommendation for use of this cap for the purpose of measuring is included on the printed label unless a handle is provided on the cap.

RESTRICTIONS

Use this product only in accordance with label directions.
Do not re-use container for any purpose. Dispose of empty container properly or burning (keep cup of ashes).

Do not apply broadcast applications to lawns. Use measuring cup of a teaspoon or measuring cup should not be used for food purposes after use with insecticides.

WARNING

Poisoning may occur as a result of swallowing. Do not eat or swallow. If swallowed, do not induce vomiting. If swallowed, do not induce vomiting. Do not breathe dust. Do not contaminate feed or food. Wash, dry, and store in a safe place. Keep children and pets away from treated area until product is removed. Do not water into soil and treated area is completely dry. Wash hands thoroughly after handling. Ingredients: Azoxystrobin sulfate and/or 2-P.A.

DI-SYSTON is Reg. in U.S. Pat. & Tm. Office. U.S. Pat. 4,811,111. Bayer A.G., Chemagro Corporation, licensed.

FOR FURTHER INFORMATION CONTACT THE MANUFACTURER OR THE DISTRIBUTOR IMMEDIATELY.

There is no known hazard to humans from the use of this product. (EPA Reg. No. 101-101) previously submitted.

3125-11
ACCEPTED
MAY 9 1968
UNDER THE FEDERAL INSECTICIDE
ACT OF 1947, THIS PRODUCT HAS
BEEN LISTED IN THE REGISTER.
EPA REG. NO. 101-101
NO OTHER COMMENTS.