

FOR EASY OPENING,



CUT ALONG THIS LINE

ACCEPTED

APR 23 1976

Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered for the pesticide use stated under EPA Reg. No.



# MORESTAN

25% WETTABLE POWDER  
MITICIDE, FUNGICIDE

**FOR EFFECTIVE CONTROL OF MITES AND POWDERY MILDEW**

NET WEIGHT 2 POUNDS

## CAUTION

See label for use and other instructions.

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT 25%  
INERT INGREDIENTS 75%  
NET WEIGHT 100

*Chemicals for Agriculture*

## CAUTION

Morestan is a highly effective miticide and fungicide. It is highly toxic to fish and aquatic life. Do not use in or near water.

Morestan is a highly effective miticide and fungicide. It is highly toxic to fish and aquatic life. Do not use in or near water.

MORESTAN is Reg. TM of the Pesticide Industry. For more information, contact Chemagro Corp., P.O. Box 1000, Kansas City, MO 64101.

## DIRECTIONS FOR USE

Apply Morestan to the foliage of plants and trees. Use a spray nozzle that produces a fine mist. Apply during the early morning or late afternoon. Do not apply when the temperature is below 50°F or above 90°F. Do not apply to plants that are stressed or damaged.

## RECOMMENDED APPLICATIONS

| CROP       | RATE                   | INTERVAL  |
|------------|------------------------|-----------|
| Apple      | 1/2 to 1 lb/1000 sq ft | 7-14 days |
| Blueberry  | 1/2 to 1 lb/1000 sq ft | 7-14 days |
| Cherry     | 1/2 to 1 lb/1000 sq ft | 7-14 days |
| Orange     | 1/2 to 1 lb/1000 sq ft | 7-14 days |
| Peach      | 1/2 to 1 lb/1000 sq ft | 7-14 days |
| Pineapple  | 1/2 to 1 lb/1000 sq ft | 7-14 days |
| Strawberry | 1/2 to 1 lb/1000 sq ft | 7-14 days |
| Tomato     | 1/2 to 1 lb/1000 sq ft | 7-14 days |
| Wine grape | 1/2 to 1 lb/1000 sq ft | 7-14 days |

Chemagro

Chemagro Corp. P.O. Box 1000  
Kansas City, MO 64101

Chemagro

FOR EASY OPENING,



CUT ALONG THIS LINE

ACCEPTED  
APR 23 1976  
The Federal Insecticide,  
Fungicide and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA No.



# MORESTAN

25% WETTABLE POWDER  
MITICIDE, FUNGICIDE


**FOR EFFECTIVE CONTROL OF MITES AND POWDERY MILDEW**

NET WEIGHT 2 POUNDS

### CAUTION

See side panel for caution statements

KEEP OUT OF REACH OF CHILDREN



*Chemicals for Agriculture*

### CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing of spray mist. Wash thoroughly with soap and warm water after handling.

Under conditions of tight-fitting clothing, spray mist may present a respiratory hazard. MORESTAN may irritate the skin, eyes, nose and throat. When applied to crops, it may injure plants. When applied to trees, it may injure bark. Do not apply to plants that are stressed or that are in bloom. Do not apply to plants that are in fruit. Do not apply to plants that are in flower. Do not apply to plants that are in seed. Do not apply to plants that are in fruit. Do not apply to plants that are in flower. Do not apply to plants that are in seed.

Do not use indoors. Do not use in food. Do not use in feed. Do not use in water.

MORESTAN is a Reg. TM of the Bayer Corporation, 501 Main Street, New York, NY 10017.

EPA Reg. No. 100-100000

### DIRECTIONS FOR USE

Apply MORESTAN to plants and soil in the following manner: ...

### RECOMMENDED APPLICATIONS

| Crop                           | Application Rate | Frequency |
|--------------------------------|------------------|-----------|
| Apples                         | 1/2 lb per acre  | 2-3 times |
| Oranges                        | 1/2 lb per acre  | 2-3 times |
| Pears                          | 1/2 lb per acre  | 2-3 times |
| Plums                          | 1/2 lb per acre  | 2-3 times |
| Prunes                         | 1/2 lb per acre  | 2-3 times |
| Almonds                        | 1/2 lb per acre  | 2-3 times |
| Walnuts                        | 1/2 lb per acre  | 2-3 times |
| Pistachios                     | 1/2 lb per acre  | 2-3 times |
| Cherries                       | 1/2 lb per acre  | 2-3 times |
| Peaches                        | 1/2 lb per acre  | 2-3 times |
| Apricots                       | 1/2 lb per acre  | 2-3 times |
| Quinces                        | 1/2 lb per acre  | 2-3 times |
| Small Fruits                   | 1/2 lb per acre  | 2-3 times |
| Strawberries                   | 1/2 lb per acre  | 2-3 times |
| Raspberries                    | 1/2 lb per acre  | 2-3 times |
| Blackberries                   | 1/2 lb per acre  | 2-3 times |
| Blueberries                    | 1/2 lb per acre  | 2-3 times |
| Cranberries                    | 1/2 lb per acre  | 2-3 times |
| Gooseberries                   | 1/2 lb per acre  | 2-3 times |
| Currents                       | 1/2 lb per acre  | 2-3 times |
| Unguarded Berries              | 1/2 lb per acre  | 2-3 times |
| Black Currants                 | 1/2 lb per acre  | 2-3 times |
| Red Currants                   | 1/2 lb per acre  | 2-3 times |
| White Currants                 | 1/2 lb per acre  | 2-3 times |
| Goose Currants                 | 1/2 lb per acre  | 2-3 times |
| Common Huckleberries           | 1/2 lb per acre  | 2-3 times |
| Salmonberries                  | 1/2 lb per acre  | 2-3 times |
| Loganberries                   | 1/2 lb per acre  | 2-3 times |
| Marionberries                  | 1/2 lb per acre  | 2-3 times |
| Thornless Raspberries          | 1/2 lb per acre  | 2-3 times |
| Thornless Blackberries         | 1/2 lb per acre  | 2-3 times |
| Thornless Blueberries          | 1/2 lb per acre  | 2-3 times |
| Thornless Cranberries          | 1/2 lb per acre  | 2-3 times |
| Thornless Gooseberries         | 1/2 lb per acre  | 2-3 times |
| Thornless Currents             | 1/2 lb per acre  | 2-3 times |
| Thornless Unguarded Berries    | 1/2 lb per acre  | 2-3 times |
| Thornless Black Currants       | 1/2 lb per acre  | 2-3 times |
| Thornless Red Currants         | 1/2 lb per acre  | 2-3 times |
| Thornless White Currants       | 1/2 lb per acre  | 2-3 times |
| Thornless Goose Currants       | 1/2 lb per acre  | 2-3 times |
| Thornless Common Huckleberries | 1/2 lb per acre  | 2-3 times |
| Thornless Salmonberries        | 1/2 lb per acre  | 2-3 times |
| Thornless Loganberries         | 1/2 lb per acre  | 2-3 times |
| Thornless Marionberries        | 1/2 lb per acre  | 2-3 times |

emagro

CHEMAGRO AGRICULTURAL DIVISION  
MAYTAG CHEMICAL CORPORATION  
P.O. BOX 1000, MARIETTA, GA 30067

Chemagro

