U.S. LABEL

<u>Reason to Issue</u>: Under citrus, (1) propose rate range for Florida and Texas citrus, and to show maximum of 5 pounds of product per acre and (2) add summer oil precaution; modify spray adjuvant statement under RESTRICTIONS.

<u>Date of Draft</u>: <u>3/27/90 (Pre-Reg) (S)</u>

EPA Reg. No. 3125-117

MORESTAN®
25% Wettable Powder

ACTIVE INGREDIENT:
6-methyl-1,3-dithiolo[4,5-<u>b</u>]
quinoxalin-2-one 25%

INERT INGREDIENTS: ....<u>75%</u>
100%

AMENDMENT

To Previously Registered Labeling

ACCEPTED

JAN 2 5 1991

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under
EPA Reg. No.   117

<u>Revise as Follows</u>:

RECOMMENDED APPLICATIONS

CROP	PEST	Pounds MORESTAN 25% WP	REMARKS
BEARING CITRUS TREES Citrus (Arizona, and California Only)	Citrus red mite Citrus rust mite Texas citrus mite Mite eggs	5	Apply specified dosage per acre by concentrate or dilute ground spray using 20 to 500 gallons of water per acre. Schedule applications no closer than 90 days apart in Texas and Florida, or 180 days apart in Arizona and California. Allow at least 21 days between last application and harvest.
Citrus (Florida and Texas Only)	Citrus rust mite Texas citrus mite Mite eggs	<u>2-1/2 to 5</u>	<u>Do not use MORESTAN 25% Wettable Powder within 3 weeks of summer oil sprays.</u> NOTE: <u>A maximum of 5 pounds of MORESTAN 25% Wettable Powder may be used per acre per year on bearing fruit</u>.

RESTRICTIONS

Under certain adverse weather conditions, particularly those that create slow drying of spray deposits, injury may occur to foliage of strawberries, and deciduous fruit trees having new and/or tender growth.

Do not use summer oils if MORESTAN 25% Wettable Powder is used in the cover spray program.

Do not use petroleum based spray adjuvants with MORESTAN 25% Wettable Powder. Consult your supplier for approved adjuvants.

Do not allow sprays to drift onto other crops, including pastures. Do not graze treated areas.

Do not use on other crops used for food and forage. Use only according to label directions. Applications at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse.

All copy, except that which is underlined, has been previously accepted for registration.