

BEST DOCUMENT AVAILABLE

000000
000000

335-124

Container and waste disposal.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Caution.

May be harmful if swallowed or absorbed through the skin. Do not contaminate feed or food. Avoid prolonged skin contact. Wash thoroughly after using.

Symptoms of poisoning: A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea.

Treatment: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, administer milk or water freely and induce vomiting by giving one dose (1/2 oz. or 15 ml.) of syrup of ipecac. If vomiting does not occur within 30 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Professional medical assistance should be secured immediately. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes.

To physician.

Antidote—Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance.

BAYGON is a Reg. TM of the Parent Company of Farbwerke Bayer GmbH, Leverkusen

777270

Baygon

2% Bait Insecticide

**for effective control
of cockroaches
and crickets.**

ACTIVE INGREDIENT

2-(1-Methylethoxy) phenol methyl-
carbamate 2%

INERT INGREDIENTS 98%

U.S. Patent No. 3,111,539 100%

Canadian Patent No. 725,204

EPA Reg. No. 3125-121-ZA

EPA Est. 3125-MO-1

STOP—Read the label before use

CAUTION.

Keep out of reach of children.

(See rear panel for caution statements)

NET WEIGHT 5 POUNDS



Chemagro Agricultural Division
Mobay Chemical Corporation
Box 4913, Kansas City, Mo. 64120

BEST DOCUMENT AVAILABLE

3125-121

Aviso.

AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

Directions for use. Important.

Read these entire Directions and Conditions of Sale before using BAYGON 2% Bait.

Conditions of sale.

The directions on this label were determined through research to be the directions for correct use of this product. This product has been tested for a range of weather conditions similar to those weather conditions that are ordinary and customary in the geographic area where the product is used. Insufficient control of pests may result from the occurrence of extraordinary or unusual weather, or from failure to follow label directions. In addition, failure to follow label directions may cause injury to the crop to which the product is applied, other crops, animals, man, or the environment. Chemagro and the seller offer, and the buyer accepts and uses, this product subject to the conditions that extraordinary or unusual weather, or failure to follow label directions are beyond the control of Chemagro and are, therefore, the responsibility of the buyer.

BAYGON 2% Bait insecticide, a ready-to-use product, is very effective for the control of crickets and the following cockroaches listed below. It gives rapid kill of these insects including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. Crickets and roaches die quickly after feeding on the bait.

Cockroaches: American, Brown, Brown banded, German, and oriental roaches.
Crickets.

For use in and around homes, apartment buildings, stores, restaurants (non-food areas) and warehouses. Apply bait lightly to floors near baseboards, in closets, under sinks and refrigerators, around and inside garbage cans, and in other areas where insects are found but which are inaccessible to children such as cracks and crevices. Do not apply to animals. Do not treat portions of buildings where animals can feed on bait. Use approximately 4 ounces of bait per 1,000 sq. ft. of space and scatter lightly so that bait is barely visible. Repeat as necessary. Keep out of reach of children.

For use by Pest Control Operators in restaurants (food areas), poultry-packing, red meat packing and food processing plants, including plants that are federally inspected. Apply bait on paper, pasteboard, or other material that will permit removal. Locate these bait stations on window sills, on floor near walls, in storage areas and other areas where insects have been observed. Maintain strict count of location and number of bait stations. Do not apply bait

in edible product areas until operations have ceased for the day. Remove all bait stations from the edible products areas before resuming operations.

For use in research facilities. The use of BAYGON in and around research animal quarters for pest control will not induce changes in liver microsomal enzyme activity in these animals. BAYGON 2% Bait should not be applied directly on animals or applied where animals can feed on the bait.

For outdoor use around flower gardens, patios, driveways, sidewalks and foundations. Apply bait evenly in a 2 to 3 feet wide band at the rate of 4-oz./1,000 sq. ft. around foundations, patios, driveways or sidewalks, taking care to treat any cracks or crevices. For maximum effectiveness treat beneath concrete drain splashpans. For applications in or around flower gardens scatter bait at the rate of 4-oz./1,000 sq. ft. as a broadcast or apply a protective barrier around the garden as a 1 foot wide band.

Fish and wildlife cautions.

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Restrictions.

Do not apply to floors or other surfaces likely to be contacted by children or pets.