

3125-123

8-9-1976

4/3

RECOMMENDED APPLICATIONS

Table with columns: CROP, INSECT, GUTHION 2S, and REMARKS. Rows include Vegetables (Leaf miners, Flea beetles), Onions (Thrips), Peppers (Colorado potato beetle, European corn borer), Potatoes (Colorado potato beetle, European corn borer), Spinach (Aphids, Leaf miners), Tomatoes (Colorado potato beetle, European corn borer), Melons (Colorado potato beetle, European corn borer), and Ornamentals (Aphids, Colorado potato beetle, European corn borer).



AUG 9 1976

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended and registered with EPA Reg. No. 3125-123-ZC

GUTHION 2S EMULSIFIABLE INSECTICIDE


FOR EFFECTIVE ECONOMICAL INSECT CONTROL

DO NOT STORE BELOW 25° F. PROTECT FROM HEAT KEEP AWAY FROM OPEN FLAME DO NOT HEAT

Table with columns: ACTIVE INGREDIENTS and INERT INGREDIENTS. Active ingredients include O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate (27%), Aromatic Petroleum Distillate (52%), and Inert ingredients (26%, 100%).

Contains 2 lbs. O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl] phosphorodithioate per gallon.

NET CONTENTS 5 GALLONS



POISON DANGER

KEEP OUT OF REACH OF CHILDREN

(See rear panel for antidotes and danger statements)

EPA Reg. No. 3125-123-ZC

Chemicals for Agriculture

Chemagro Agricultural Division Moberly Chemical Corporation Box 4913, Kansas City, Missouri 64120

110 135

\*This concentration is calculated for conventional hydraulic-type sprayers. When lower volumes of spray are applied per acre with cone-type sprayers, increase the concentration of GUTHION 2S in the spray mixture in order to apply amount of GUTHION 2S per acre equivalent to a full coverage spray. Where conditions dictate an air application, apply dosage per acre equivalent to a full coverage ground spray in not less than one gallon of water per acre.



® GUTHION 2S  
EPA Reg. No. 3125-123

Since the labeling on the attached container was printed GUTHION 2S was accepted for registration by the U.S. Environmental Protection Agency for use in undiluted ultra-low-volume applications for pest control on apples, crab apples, and quinces. Full text of the accepted labeling follows:

RECOMMENDED APPLICATIONS

CROP	INSECT	Pints GUTHION 2S	REMARKS
Apples Crab apples Quinces (Washington, Oregon, Idaho, Utah, & Northern California)	Apple maggot Codling moth European apple sawfly Eye-spotted bud moth Fruit tree leaf roller Green fruitworm Orange tortrix Plum curculio Red-banded leaf roller	2 to 4	Apply specified dosage per acre undiluted by helicopter or fixed wing aircraft using equipment capable of applying ultra-low-volume sprays. Use Beecomist nozzles with 20 micron sleeves. Do not apply more than 5 times per season nor within 7 days of harvest.

NOTE: This formulation, when used undiluted, may cause spotting of automobile finishes, if prolonged exposure is permitted. Do not spray directly over automobiles. If accidental exposure does occur, automobiles should be washed immediately.

BEFORE USE, BE SURE TO FOLLOW THE OTHER USE AND HANDLING DIRECTIONS AND PRECAUTIONS FOUND ON THIS CONTAINER.

W/C