

ORTHO
Lawn Fungicide
Contains 8 DYNAMS

Controls 8 DISEASES SUCH AS BROWN PATCH, RUST, SNOW MOLD, COLLAR ROT, CORNER SPOT, RED LEAF SPOT OF BENTGRASSES, LEAF BLOTCH OF BARNYARD GRASS, HEAVY BROWN BLIGHT OF FESCUES, BROWN BLIGHT OF RYE GRASSES, ZONATE EYE SPOT, AND HEAVY BROWN LEAF SPOT OF BLUEGRASSES, BENTGRASSES, AND FESCUES.

Active Ingredient

1,4-dichloro-6-(o-chlorophenyl)-o-anisidine

Inert Ingredients.....

U.S. Pat. Nos. 2,770,480 - 3,074,481 - This material used under License from Chemagro Corporation.

8 DYNAMS is a registered trademark of Chemagro Corporation.

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE BACK PANEL FOR ADDITIONAL CAUTIONS.
(18 pt.) (12 point type)

BACK PANEL

ORTHO
Lawn Fungicide
Contains 8 DYNAMS

READ THE LABEL. USE SPECIFICALLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS. ALWAYS OBSERVE FEDERAL AND STATE REGULATIONS.

ORTHO Lawn Fungicide gives effective control of Helicostoma brown patch, Sclerotinia collar rot, corner spot, rust, snow mold (Typharia), and Helminthosporium diseases, such as leaf blight of Kentucky bluegrass, red leaf spot of bentgrasses, leaf blotch of Barnyard grass, heavy brown blight of fescues, brown blight of rye grasses, zonate eye spot, and heavy brown leaf spot of bluegrasses, bentgrasses, and fescues.

HOW TO USE:

Hand Sprayers: Put ORTHO Lawn Fungicide into sprayer jar, fill with water and stir until a uniform mixture is obtained. Apply spray mixture uniformly to lawn area. For disease prevention, use 4 ounces (approximately 1 cup) ORTHO Lawn Fungicide and spray to cover 1,000 square feet. For spot treatment of diseased areas, use 6 ounces (approximately 1 1/2 cups) ORTHO Lawn Fungicide and thoroughly wet these areas.

Back Sprayer or Sprayers: For use in hand or backpack-type sprayers, add 4 ounces (approximately 1 cup) of ORTHO Lawn Fungicide to 1 1/2 gallons of water and apply to 1,000 square feet (30 ft. x 30 ft.) of lawn or other turf area. For spot treatment of heavily diseased areas, add 6 ounces (approximately 1 1/2 cups) to 2 1/2 gallons of water and thoroughly wet these areas.

WHEN TO APPLY:

For Collar Rot and Rust Diseases: Begin applications in the spring before disease starts and repeat at 7 to 14-day intervals. During extended periods of wet weather, applications every 7 days may be required. Continue treatments throughout the spring and summer months. If necessary, water the lawn before applying ORTHO Lawn Fungicide.

For Control of Snow Mold: Apply after the first melting snow in the spring and repeat applications should be made when the lawn or other turfgrass area is free of snow in the spring. Apply a final application after the final melting of snow in the spring.

ORTHO Lawn Fungicide - 1000

Instructions

Keep out of hands of children. If swallowed or inhaled, avoid breathing fumes. Avoid breathing of dust or spray mist. Wash thoroughly with soap and water after handling. Wash contaminated clothing with soap and hot water before reuse. Empty container promptly when empty.

Do not contaminate feed, foodstuffs, water supplies, and fish ponds.

This product is toxic to fish. Keep out of any body of water.

NOTICE: Bayer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Chevron

Distributed by

CHEVRON CHEMICAL COMPANY
ORTHO DIVISION - San Francisco, California

Richmond, California

Atlanta, Georgia

Haddonfield, New Jersey

Form

USDA Reg. No. 3125-127

Product No. 1000

Made in U. S. A.

