

# ANTHON HORSE WORMER

For the Removal of Bots (*Gasterophilus spp.*),  
ascarids (*Parascaris equorum*) and pinworms  
(*Oxyuris equi*) from horses.

## DOSAGE

Give one packet (5 grams) per 250 pounds of  
body weight. For most effective control, treat-  
ment should be repeated at 3 to 4 month in-  
tervals. ANTHON Horse Wormer can be used at  
any time but do not repeat more frequently than  
every 30 days.

## DIRECTIONS

Use required dosage dry or dissolved in water  
and mix with the amount of feed to be consumed  
at one feeding.

Withdrawal of feed 12 to 18 hours prior to treat-  
ment will assure complete ingestion. Allow  
horse to consume all of the medicated feed be-  
fore additional feed is given. Remove and de-  
stroy any medicated feed not consumed within  
12 hours.

## ACTIVE INGREDIENT:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl)  
phosphonate\*† 90%

## INERT INGREDIENTS:

(Colloidal Silica and Dextrose) 10%  
100%

\*U.S. Patent No. 2,701,225; Canadian Patent No. 529,379  
†Trichlorfon

USDA Reg. No. 3125-131

SEE INSIDE FLAP FOR WARNING STATEMENTS



**CHEMAGRO  
CORPORATION**

KANSAS CITY, MISSOURI 64120

NET CONTENTS 5 GRAMS

**WARNING: KEEP OUT OF REACH OF CHILDREN**  
**NET CONTENTS 5 GRAMS**



USE BY DECEMBER, 1972

1 (ct. 2)

## ANTHON HORSE WORMER

For the Removal of Bots (*Gasterophilus* spp.), ascarids (*Parascaris equorum*) and pinworms (*Oxyuris equi*) from horses.

### DOSAGE

Give one packet (5 grams) per 250 pounds of body weight. For most effective control, treatment should be repeated at 3 to 4 month intervals. ANTHON Horse Wormer can be used at any time but do not repeat more frequently than every 30 days.

### DIRECTIONS

Use required dosage dry or dissolved in water and mix with the amount of feed to be consumed at one feeding.

Withdrawal of feed 12 to 18 hours prior to treatment will assure complete ingestion. Allow horse to consume all of the medicated feed before additional feed is given. Remove and destroy any medicated feed not consumed within 12 hours.

### ACTIVE INGREDIENT:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate\*† 90%

### INERT INGREDIENTS:

(Colloidal Silica and Dextrose) 10%  
100%

\*U.S. Patent No. 2,701,225; Canadian Patent No. 529,379

†Trichlorfon

USDA Reg. No. 3125-131

### RESTRICTIONS:

Use only as directed. Do not treat horses to be used for food. Do not treat sick or debilitated horses, colts under 4 months of age, mares in the last month of pregnancy, or animals other than horses. Do not exceed recommended dose. Do not administer in conjunction with or for 10 days after treatment with other organic phosphates or cholinesterase inhibitors. Do not administer intravenous anesthetics, especially muscle relaxants, for a period of 2 weeks after using this product.

### NOTICE TO VETERINARIANS:

Symptoms of over dosage in horses are ataxia and colic. Atropine is antidotal. Give 100 milligrams or 1½ grains atropine per 1,000 pounds body weight.

### WARNING:

Not for human use. Keep out of reach of children. Avoid prolonged or repeated contact with the skin. Wash hands with soap and water after use. If swallowed by humans, induce vomiting by giving 2 tablespoonfuls of salt in a glass of water. Call physician immediately.

### TO PHYSICIAN:

This material is a cholinesterase inhibitor. Atropine and 2-PAM are antidotal. Give supportive treatment. Do not walk patient nor give morphine.

964111