

C-228471

Base Pre-Reg. (2131)

U.S. LABEL (S)

Reason to Issue: To revise in accordance with cancellation notice published in Federal Register of 7/17/73 showing non-food use only on deciduous fruit (pre-bloom, post-harvest), non-bearing fruit and ornamentals. 2/25/74 delete use on bearing deciduous fruit.

Date of Draft: 2/25/74  
Supersedes Pre-Reg. Draft of 1/1/74

Ⓜ  
MORESTAN

2% DUST

MITICIDE

FOR EFFECTIVE CONTROL OF MITES  
AND MITE EGGS

NET WEIGHT                      POUNDS

CAUTION  
KEEP OUT OF REACH OF CHILDREN

(See rear panel for caution  
statements)

GUARANTEE

ACTIVE INGREDIENT:

6-Methyl-1,3-dithioleptan-2-one  
quinoxalin-2-one \*

INERT INGREDIENTS: ..... 98  
100

\*U.S. Patent No. 3,091,612  
Canadian Patent No. 659,270

EPA Reg. No. 3125-112

ACCEPTED


STOP - READ THE LABEL BEFORE USE

MANUFACTURED BY

CHEMACRO  
Division of Bionex Corporation  
Box 4913, Kansas City, Mo. 64120

Note: Underlined letter in chemical nomenclature should be italicized when printed, not underlined.

11-1-74



MORESTAN 2 Dust

DIRECTIONS FOR USE

IMPORTANT: Read these entire Directions and Conditions of Sale, including the Warranties and Limitation of Damages provision, before using MORESTAN 2 Dust.

CONDITIONS OF SALE: These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use under a variety of abnormal conditions beyond the control of the seller. Therefore, the seller, distributor, and the Buyer uses, this product subject to the understanding that the Buyer assumes all such risks.

MORESTAN Miticide-Fungicide is recommended for residual control of mites and/or mite eggs on deciduous fruits, strawberries, and ornamentals. For best results, apply a thorough coverage before mites or mite eggs are present in large numbers and before mites cause bronzing of the foliage. Apply during good drying conditions. (See Restrictions.) Consult your State Agricultural Extension Service or Imperial Station for detailed information on timing of applications for most effective control.

RECOMMENDED APPLICATIONS

CROP	PEST	Per acre MORESTAN 2 Dust	REMARKS
<u>NON-BEARING FRUIT</u>			
Apples			
Apricots			
Cherries			
Nectarines	Mites	30 to 50	Apply specified rate per acre.
Peaches	Mite eggs		Repeat as necessary. Do not apply to fruit trees that will be cut within one year of application.
Pears			
Plums			
Prunes			
Strawberries	Mites (except cyclamen mite) Mite eggs	30 to 50	Apply specified rate per acre. Repeat as necessary.
<u>ORNAMENTALS</u>			
Arborytae			
Juniper	Mites	30 to 50	Apply specified rate per acre.
Roses	Mite eggs		Repeat as necessary.
Yew			

RESTRICTIONS

Under certain adverse weather conditions such as high humidity it may cause injury to foliage of strawberries or damaged fruit. Do not use summer fall if MORISTAN 21 is used in the summer program.

MORISTAN Miticide-Fungicide is toxic to fish.  
Do not contaminate streams, lakes, or ponds with this material.

Do not allow dust to drift onto other crops, including pastures. Do not graze treated areas.

Do not use on other crops used for food or forage. Use only according to label directions. Applications at rates above those shown may result in illegal drug residues. Do not treat food crops grown in the greenhouse.

Phytotoxicity has occurred on the foliage and flowers of several varieties of grapes, especially during periods of high temperatures. Treat roses only after flowering and before new growth appears. Before treating large plantings for the first time, treat a few plants and observe for 4 or 5 days. Under certain conditions, damage to other plants may occur.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by turning with a pick or burning (stay away from leaks).

CAUTION

Harmful if swallowed or inhaled. MORISTAN 21 Dust may cause mild irritation of the respiratory tract and, therefore, contact with skin, eyes, or clothing should be avoided. Avoid breathing of dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate feed or food. Keep out of reach of children.

WARRANTY AND LIMITATION OF REMEDY

Chemagro warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the label as for the conditions of sale, subject to the sales referred to here in. CHEMAGRO MAKES NO EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPLICIT OR IMPLIED WARRANTY, OF MERCHANTABILITY, and no agent or Contractor is authorized to do so. This warranty with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential or special damages such as loss of profits or values, etc.

MORISTAN is a Reg. U.S. Pat. & Tm. Off. of the Par. at Company of Carbon-Fiberized Silver Dust, Inc.

6/10/74  
Pesticides Manual