

Reason to Issue: Note alfalfa and clover
to include mixed stands with grasses,
add new use on grasses; add certain
new insects to alfalfa and clover;
increase number of applications on
alfalfa, clover and corn; delete
PHI on alfalfa, clover, cereal, and
corn forage. 6/29/72 draft: to
recommend minimum of 1 pound active
for armyworms in alfalfa and clover,
and for armyworms, fall armyworms, and
plant bugs in grasses.

U.S. LABEL

Date of Draft: 6/29/72 (Pre-Reg.) (S)

Supersedes Pre-Reg. Draft: 1/28/71

EPA Reg. No. 3125-143


®DYLOX

Liquid Solution

INSECTICIDE

AMENDMENT

To Previously Registered Label



ADD THE FOLLOWING:

RECOMMENDED APPLICATIONS

CROP	INSECT	Pints DYLOX LS	REMARKS
Alfalfa clover (including mixed stands with grasses)	Alfalfa caterpillar	3/4 to 1	Apply specified dosage per acre using sufficient water for complete cover- age but not less than 1 gallon per acre. Three applications may be made per season with the first ap- plication at the first cutting.
	Alfalfa webworm	1/2 to 2	
	Western yellow- striped armyworm	1	
	Feet armyworm		
	Leafhoppers	1 to 2	
	Variiegated cutworm		
	Armyworm		
Alfalfa Plant bug			
	Lygus bugs	2	
	Stink bugs		
	Tarnished plant bug		

RECOMMENDED APPLICATIONS
(Continued)

CROP	INSECT	Pints DYLOX LS	REMARKS
FIELD CROPS (Cont'd)			
<u>Corn</u>	<u>Armyworms</u> <u>Cutworms</u>	1 to 2	Apply specified dosage per acre in sufficient water for complete coverage but not less than 1 gallon per acre. Three applications may be made per season. For early application, when plants are 3 to 12 inches high, direct the spray to the lower portions of plants and to the soil around the base of the plants. Later applications may be made as full coverage spray up to day of harvest.
<u>Barley</u>	<u>Armyworms</u> <u>Beet webworm</u>	1 to 2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Three applications may be made per season but not within 21 days of harvest.
<u>Flax</u>	<u>Variegated cutworm</u>		
<u>Oats</u>	<u>Bertha armyworm</u>	2	
<u>Wheat</u>	<u>Diamondback moth</u>		
			NOTE: Applications may be made without removal of grazing livestock.
<u>Grasses</u> (Including <u>Rangeland</u> & <u>Pasture</u>)	<u>Range Caterpillar</u> <u>Leafhoppers</u> <u>Armyworm</u> <u>Hill Armyworm</u> <u>Plant bugs</u>	1 to 2 2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Three applications may be made per cutting with the last application up to the day of cutting. Do not apply more than 3 times per season if crop is not cut for hay. NOTE: Applications may be made without removal of grazing livestock.

ALL LABEL DATA ABOVE, EXCEPT THAT WHICH IS UNDERLINED, HAS HAD PREVIOUS REGISTRATION ACCORDANCE.