

3125-148

05/02/1966

1/3

®
Baytex

**8 CONCENTRATE
 INSECTICIDE**

®

ACCEPTED

MAY 2 1966

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 3125-148 SUBJECT
 TO ATTACHED

FOR EFFECTIVE CONTROL OF MOSQUITOES AND FLIES

GUARANTEE: Contains 8 lbs. *O,O*-Di-
 methyl *O*-[4-(methylthio)-*m*-tolyl] phos-
 phorothioate per gallon.

NET CONTENTS GALLONS

WARNING

(See rear panel for warning statements.)

**KEEP OUT OF REACH OF
 CHILDREN.**

ACTIVE INGREDIENTS:

<i>O,O</i> -Dimethyl <i>O</i> -[4-(methylthio)- <i>m</i> -tolyl] phosphorothioate †	81%
Aromatic Petroleum Distillate	19%
	100%

† U.S. Patent No. 3,042,703
 Canadian Patent No. 631,472

USDA Reg. No. 3125-148



**CHEMAGRO
 CORPORATION**
 KANSAS CITY, MISSOURI



DIRECTIONS FOR USE

AREA MOSQUITO APPLICATIONS TO POPULATED AND RURAL AREAS ARE TO BE MADE ONLY BY MOSQUITO ABATEMENT DISTRICTS, PUBLIC HEALTH OFFICIALS, PUBLIC MOSQUITO CONTROL OFFICIALS, AND COMMERCIAL PEST CONTROL OPERATORS.

BAYTEX 8 Concentrate is recommended for use wherever mosquitoes and flies are a problem. It provides long residual control and is as effective against DDT-resistant strains as non-resistant strains. The spray can be applied to most surfaces, including freshly whitewashed areas, without loss of biological effectiveness.

MIXING: BAYTEX 8 Concentrate contains no emulsifying agents and should be diluted only with oil carriers such as No. 2 fuel oil (diesel) or kerosene. Should it be desirable to use water as a carrier, sufficient quantities of suitable emulsifiers should be added to the BAYTEX 8 Concentrate prior to mixing with water. These water or oil mixtures are suitable for use in hand or power-operated sprayers or foggers as well as with aircraft sprayers or in mist-type applicators. To mix with oil, pour required amount of BAYTEX 8 Concentrate into full amount of oil and then agitate.

If the mixture of BAYTEX and oil, prepared according to label directions, is to be held in storage for longer than a few hours, it is necessary to add an anti-sludge agent to the mixture. Request information from your dealer on the type and amount of anti-sludge agent to use.

MIXING: Use specified dosage of BAYTEX 8 Concentrate in the amount of water or oil necessary to give uniform coverage. The type of equipment used will determine the concentration required.

SPRAYING: For mosquito larvae control, applications should be made to moist areas, standing water pools, marshes, swamps, and tidal areas. Also spray around and in catch basins and similar areas where mosquitoes breed. Do not apply to ponds, lakes, streams, or other bodies of water containing valuable fish. Allow at least three weeks between applications.

For large area mosquito control, applications should be made to wet grassy, bushy, or woody areas.

For localized mosquito and fly control, applications should be made to outside surfaces of buildings, screen doors, window frames, around outdoor light fixtures, and in garages.

FOG AND MIST APPLICATIONS: Outdoor applications for adult mosquito control may be made by thermal fogging, mist spraying, airplane spraying, or residual spraying. Do not apply to ponds, lakes, streams, or other bodies of water containing valuable fish. Allow at least 3 weeks between mist applications. Fog applications may be repeated as necessary.

RESTRICTIONS

1. Do not allow spray treatment to drift on pasture land, crop land, poultry ranges, or water supplies.
2. Do not use inside homes, in dairy barns, poultry houses, or buildings where food is processed. Do not apply directly to animals.
3. Do not use on crops used for food, forage, or pasture.
4. Do not apply oil sprays directly to flowers or other ornamental plants which may be damaged by the oil. Fog application will not cause plant injury.

WARNING

May be fatal if swallowed, inhaled, or absorbed through the skin. Rapidly absorbed through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe spray mist. Wash thoroughly with soap and warm water after handling. Wash clothing with soap and hot water before re-use.

Avoid contamination of feed or food. Keep out of reach of children.

TO PHYSICIAN:

ANTIDOTE—Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

CHEMAGRO

RECOMMENDED



APPLICATIONS

INSECT	Fluid Ounces BAYTEX Conc.	REMARKS
Mosquito Larvae	$\frac{3}{4}$	SPRAY: Apply specified dosage (0.05 lb. active) per acre in sufficient water* or oil** to obtain uniform coverage, or per $1\frac{1}{2}$ gallons of water* or oil** for localized applications. Apply as a light, uniform spray. Allow at least 3 weeks between treatments.
Mosquitoes	$\frac{1}{2}$ to $1\frac{1}{4}$	THERMAL FOG: Mix specified dosage per gallon of oil.** This is equivalent to $\frac{1}{2}$ to 1 gallon of concentrate per 100 gallons of oil. Apply with standard fogging machines calibrated to deliver 40 gallons per hour at a machine speed of 5 m.p.h. to cover a swath width of up to 350 feet. Repeat as necessary.
Mosquitoes	$\frac{3}{4}$ to $1\frac{1}{2}$	MIST SPRAY: Apply specified dosage (0.05 to 0.1 lb. active) per acre. For mist-blower machines calibrated to deliver 100 gallons per hour traveling at a speed of 4 m.p.h. to cover a swath width of up to 350 feet, use 1 to 2½ gallons of concentrate per 100 gallons of water* or oil.** Allow at least 3 weeks between treatments.
Mosquitoes	(See Remarks)	AIRPLANE APPLICATION: Use BAYTEX 8 Concentrate at dosages of 0.05 to 0.1 lb. active ingredient per acre in water* or oil.** Mix 5 to 10 pints of concentrate in 100 gallons of water* or fuel oil** and apply 1 gallon of this mixture per acre. Allow at least 3 weeks between treatments.
Mosquitoes Houseflies	1 to 2	RESIDUAL SPRAY: For long residual control of mosquitoes and for residual fly control, apply specified dosage in 1 gallon of water* for application per 500 square feet or to run-off in localized areas. Residual control lasts 7 to 8 weeks for the lower rate and even longer at the higher rate. Repeat as necessary. Do not apply as a space spray.

*When using water as a carrier add sufficient quantities of suitable emulsifying agents to BAYTEX 8 Concentrate prior to mixing with water.

**Kerosene-type oils, fuel oils, diesel oils, and other base oils suitable for insecticide use.

NOTICE

Chemagro Corporation and the seller make no warranty of any kind, express or implied, on the use of this material and assume no responsibility whatsoever for any personal injury, property damage, or other type of loss resulting from the use or handling of this material. The user assumes all risks for use and handling whether in accordance with directions or not.

PATENT LICENSE

The principal active ingredient of this product is patented under U.S. Patent No. 3,042,703 and Canadian Patent No. 631,472 of Chemagro Corporation, Kansas City, Missouri, and the purchase of this product includes a limited license under said patents to use or reformulate the product only in the manner provided by the recommended applications on the present label. The purchaser obligates itself to notify Chemagro Corporation of any proposed use or reformulation of such licensed product not provided by said recommended applications, and to obtain Chemagro's approval in writing before making any such use or reformulation of the licensed product. Any use or reformulation intended for use of the licensed product in the home or garden field is specifically prohibited by said license.

BAYTEX is Reg. TM in U.S. and Canadian Pat. Offs. by Farbenfabriken Bayer A.G. Chemagro Corporation Licensee 563021

CHEMAGRO