

PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

3125-172

9-25-1972

1/2

U.S. LABEL

Reason to Issue: To recommend 3 post-planting applications into the sorghum whorl following an at-planting application.

Date of Draft: 6/7/72 (Pre-Reg.) (S)

EPA Reg. No. 3125-172-AA

Ⓡ

DI-SYSTON
15% Granular

SYSTEMIC INSECTICIDE

AMENDMENT

To Previously Registered Labeling

RECOMMENDED APPLICATIONS

CROP	INSECT	DOSAGE DI-SYSTON 15% Granular	REMARKS
Sorghum	Aphids (Greenbugs)	6 to 8 ozs. per 1000 ft. of row (for any row spac- ing*) -or- 5 to 6.7 pounds per acre (40-inch row spacing)	<u>APPLICATION AT PLANTING: IN-FURROW</u> Place the granules in the seed furrow approximately 1/2 inch above the seed at planting. Do not apply directly on the seed. See "NOTE" below.
		8 ozs. per 1000 ft. of row (for any row spacing*) or 6.7 pounds per acre (40-inch row spacing)	<u>APPLICATION AT PLANTING: BAND</u> Place the granules in a 4 to 6 inch band directly behind the planter shoe in front of the press wheel. Do not apply directly on the seed. See "NOTE" below.
			<u>POST-PLANTING APPLICATION: WHORL</u> Apply specified dosage by broadcasting into the whorl. Do not apply more than 3 times per season nor within 30 days of harvest of grain or 14 days of harvest for forage or fodder. See "NOTE" below.
		<u>NOTE:</u> Post-planting broadcast applications into the whorl may be made following an at-planting application.	

*ON ROW CROPS WHERE ROW SPACING IS EXTREMELY NARROW, DO NOT PLACE TREATED ZONES (BAND OR FURROW) CLOSER TOGETHER THAN 6 INCHES.

RECEIVED
SEP 25 1972

ACCEPTED
SEP 25 1972
UNDER THE FEDERAL BUREAU OF INVESTIGATION
FUNCTIONS AND RESPONSIBILITIES
FOR ECONOMIC COMMUNITIES
ED ATTACHED TO

3/25-172

NEXT

LABEL

3125-172

9-25-1972

1/2

U.S. LABEL

Reason to Issue: To propose use on
non-bearing raspberries for aphid
control.

Date of Draft: 6/28/72 (Pre-Reg.) (S)

EPA Reg. No. 3125-172-AA

©DI-SYSTON
15% Granular

SYSTEMIC INSECTICIDE

AMENDMENT
To Previously Registered Labeling

ADD THE FOLLOWING:

RECOMMENDED APPLICATIONS

CROP	INSECT	DOSAGE-DI-SYSTON 15% Granular	REMARKS
<u>NON-BEARING FRUIT</u>			
Raspberries (Nursery stock only) (Northeast States Only)	Aphids	1.2 lb./1000 feet of row (for any row spacing*)	<u>BAND APPLICATION:</u> Apply specified dosage in a 24 inch band over the row to new or established plantings. Work granules 2 to 3 inches into the soil using a cultivator or other suitable equipment. A second application of granules may be applied to the soil later in the season. Consult your State Agricultural Experiment Station or Extension Service for proper timing of applications.
		53.3 lbs. per acre	<u>BROADCAST APPLICATION:</u> Apply specified dosage per acre as a broadcast application over new or established plantings. Cultivate or otherwise work granules into top 2 to 3 inches of soil. A second broadcast application may be made later in the season. Consult your State Agricultural Experiment Station or Extension Service for proper timing of applications.

*ON ROW CROPS WHERE ROW SPACING IS EXTREMELY NARROW, DO NOT PLACE TREATED ZONES
(BAND OR FURROW) CLOSER TOGETHER THAN 6 INCHES.

SEP 25 1972

ONE OF THE FEDERAL FUNGICIDE AND RTO FOR ECONOMIC ED UNDER NO. TO ATTACHED COMMENTS.

3185-172