

3125-172

10-30-1972

U.S. LABEL

Reason to Issue: To propose use on
non-bearing raspberries for aphid
control; 10/11/72 draft to increase
band application rate as per EPA.

Date of Draft: 10/11/72 (Pre-Reg.) (S)

Supersedes Pre-Reg. Draft: 6/28/72

EPA Reg. No. 3125-172-AA

©DI-SYSTON

15% Granular

SYSTEMIC INSECTICIDE

AMENDMENT

To Previously Registered Labeling

ACCEPTED

OCT 30 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

3125-172

ADD THE FOLLOWING:

RECOMMENDED APPLICATIONS

| CROP | INSECT | DOSAGE - DI-SYSTON 15% Granular | REMARKS |
|---|--------|--|---|
| NON-BEARING FRUIT | | | |
| Raspberries (Nursery stock only) (Northeast States Only) | Aphids | 2.5 lb./1000 feet of row (for any row spacing*) | BAND APPLICATION: Apply specified dosage in a 24-inch band over the row to new or established plantings. Work granules 2 to 3 inches into the soil using a cultivator or other suitable equipment. A second appli- cation of granules may be applied to the soil later in the season. Consult your State Agricultural Experiment Station or Extension Service for proper timing of appli- cations. |
| | | 53.3 lbs. per acre | BROADCAST APPLICATION: Apply speci- fied dosage per acre as a broadcast application over new or established plantings. Cultivate or otherwise work granules into top 2 to 3 inches of soil. A second broadcast appli- cation may be made later in the sea- son. Consult your State Agricultural Experiment Station or Extension Ser- vice for proper timing of applications. |

*ON ROW CROPS WHERE ROW SPACING IS EXTREMELY NARROW, DO NOT PLACE TREATED ZONES (BAND
OR FURROW) CLOSER TOGETHER THAN 6 INCHES.