

3125-172

7-9-1973

U.S. LABEL

Label to Issue: To recommend applica-
tions to "Hill-dropped Cotton".

Date of Draft: 1/25/73 (Pre-Reg.) (S)

EPA Reg. No. 3125-172

ACCEPTED**JUL 9 1973**

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT-GIDE ACT
FOR ECONOMIC POISON REG'STER-
ED UNDER NO. 3125-172 SUBJECT
TO ATTACHED COMMENTS.

Ⓡ

DI-SYSTON

15% Granular

SYSTEMIC INSECTICIDE

AMENDMENT

To Previously Registered Labeling

AMEND AS FOLLOWS:

RECOMMENDED APPLICATIONS

CROP	INSECT	DOSAGE - DI-SYSTON 15% Granular	REMARKS
Cotton	Aphids Mites Thrips	5 to 8 ounces per 1000 feet of row (for any row spac- ing*) 4 to 6.4 pounds per acre (40-inch row spacing)	At Planting: Place the granules in the seed furrow or in a band on each side of the fur- row. RATE CHART FOR PLANTING TIME APPLICATION
			Heavy Soils Medium Soils Light Soils
			8 oz. 15% 6.6 oz. 15% 5 oz. 15%
			Gran./1000 Gran./1000 Gran./1000
Cotton	Aphids Mites Thrips	5 to 8 ounces per 1000 feet of row (for any row spac- ing*) 4 to 6.4 pounds per acre (40-inch row spacing)	ft. of row ft. of row ft. of row
			6.7 lbs. 15% 5.3 lbs. 15% 4 lbs. 15%
			Gran./acre. Gran./acre. Gran./acre
			Post-Plant: (Irrigated Cotton Only) If an additional treatment is necessary, make one side-dress application or a single broad- cast application over the foliage. Rates up to 13.3 pounds per acre can be applied to first squaring cotton (90 days before harvest) and rates up to 6.7 pounds per acre can be applied to within 28 days of harvest. Do not apply more than twice per season. Allow a minimum of 21 days between applications. Read "IMPORTANT" statement under "Recommended Applications" for additional information before use on this crop.
Cotton	Aphids Mites Thrips	2.5 to 3.3 ounces per 1000 feet of row (for any row spac- ing*) -or- 2 to 2.7 pounds per acre (40-inch row spacing)	HILL DROPPED COTTON: Apply specified dosage with the seed at planting. Use the lower rate where hill spacing is greater than 14 inches apart or where applications are made to light, sandy soils. Read "IMPORTANT" statement under "Recommended Applications" for additional information before use on this crop.

*ON ROW CROPS WHERE ROW SPACING IS EXTREMELY NARROW, DO NOT PLACE TREATED ZONES
(BAND OR FURROW) CLOSER TOGETHER THAN 6 INCHES.

NOTE: All copy except that which is underlined has previously been granted registra-
tion.